

# TROPICALIZATION, SYMMETRIC POLYNOMIALS, AND COMPLEXITY

ALEXANDER WOO AND ALEXANDER YONG

ABSTRACT. D. Grigoriev-G. Koshevoy recently proved that tropical Schur polynomials have (at worst) polynomial tropical semiring complexity. They also conjectured tropical skew Schur polynomials have at least exponential complexity; we establish a polynomial complexity upper bound. Our proof uses results about (stable) Schubert polynomials, due to R. P. Stanley and S. Billey-W. Jockusch-R. P. Stanley, together with a sufficient condition for polynomial complexity that is connected to the saturated Newton polytope property.

## 1. INTRODUCTION

The **tropicalization** of a polynomial

$$f = \sum_{(i_1, i_2, \dots, i_n) \in \mathbb{Z}_{\geq 0}^n} c_{i_1, \dots, i_n} x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n} \in \mathbb{C}[x_1, x_2, \dots, x_n]$$

(with respect to the trivial valuation  $\text{val}(a) = 0$  for all  $a \in \mathbb{C}^*$ ) is defined to be

$$(1) \quad \text{Trop}(f) := \max_{i_1, \dots, i_n} \{i_1 x_1 + \cdots + i_n x_n\}.$$

This is a polynomial over the tropical semiring  $(\mathbb{R}, \oplus, \odot)$ , where

$$a \oplus b = \max(a, b) \text{ and } a \odot b = a + b$$

respectively denote tropical addition and multiplication, respectively. We refer to the books [ItMiSh09, MaSt15] for more about tropical mathematics.

Let  $\text{Sym}_n$  denote the ring of symmetric polynomials in  $x_1, \dots, x_n$ . A linear basis of  $\text{Sym}_n$  is given by the *Schur polynomials*. These polynomials are indexed by partitions  $\lambda$  (identified with their Ferrers/Young diagrams). They are generating series over semistandard Young tableaux  $T$  of shape  $\lambda$  with entries from  $[n] := \{1, 2, \dots, n\}$ :

$$s_\lambda(x_1, \dots, x_n) := \sum_T x^T \quad \text{where } x^T := \prod_i x_i^{\#i\text{'s in } T}.$$

The importance of this basis stems from its applications to, for example, enumerative and algebraic combinatorics, the representation theory of symmetric groups and general linear groups, and Schubert calculus on Grassmannians; see, for example, [Fu97, St99].

D. Grigoriev and G. Koshevoy [GrKo16] studied the complexity of the tropical polynomial  $\text{Trop}(s_\lambda)$  over  $(\mathbb{R}, \oplus, \odot)$ . An *arithmetic circuit* is a circuit where inputs are each labelled by a single variable  $x_i$  or a fixed constant, each gate performs a single  $\oplus$  or  $\odot$  operation, and there is one output. An arithmetic circuit  $C$  naturally gives an expression  $\text{res}(C)$ , the tropical polynomial in the variables  $x_1, \dots, x_n$  that it computes. The circuit  $C$  *evaluates*  $f$  if  $\text{res}(C) = f$  as tropical polynomials, meaning that one can show  $\text{res}(C) = f$  using the *tropical semiring axioms*, by which we mean the semiring axioms along with the idempotence

---

Date: October 9, 2017.

property  $a \oplus a = a$ . The *tropical semiring complexity* of  $f$  is the smallest number of gates in a circuit  $C$  evaluating  $f$ ; see [JeSn82, Section 2].

The following is [GrKo16, Theorem 2.1]:

**Theorem 1.1** (D. Grigoriev-G. Koshevoy). *The tropical semiring complexity of  $\text{Trop}(s_\lambda)$  is at most  $O(n^2 \cdot \lambda_1)$ .*

The *skew-Schur polynomial*  $s_{\lambda/\mu}(x_1, \dots, x_n) = \sum_T x^T$  is the generating series for semistandard tableau of skew shape  $\lambda/\mu$  with entries from  $[n]$ . When  $\mu = \emptyset$  then  $s_{\lambda/\emptyset} = s_\lambda$ ; hence skew-Schur polynomials generalize Schur polynomials. Also,

$$(2) \quad s_{\lambda/\mu} = \sum_{\nu} c_{\mu,\nu}^\lambda s_\nu,$$

where  $c_{\lambda,\mu}^\nu \in \mathbb{Z}_{\geq 0}$  is the *Littlewood-Richardson coefficient*. The next statement is from [GrKo16, Section 5]:

**Conjecture 1.2** (D. Grigoriev-G. Koshevoy). *The tropical semiring complexity of  $\text{Trop}(s_{\lambda/\mu})$  is at least exponential.*

We will show the following:

**Theorem 1.3.** *There is an explicitly described  $\beta$ , depending on  $\lambda/\mu$ , with  $\beta_1 = \lambda_1$ , such that*

$$\text{Trop}(s_{\lambda/\mu}(x_1, \dots, x_n)) = \text{Trop}(s_\beta(x_1, \dots, x_n))$$

over the tropical semiring  $(\mathbb{R}, \oplus, \odot)$ .

*Example 1.4.* Let  $\lambda = (2, 1)$  and  $\mu = (1)$ . Then the tableaux contributing to  $s_{\lambda/\mu}$  are:

$$\begin{array}{|c|} \hline 1 \\ \hline \end{array}, \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array}, \begin{array}{|c|} \hline 2 \\ \hline 1 \\ \hline \end{array}, \begin{array}{|c|} \hline 2 \\ \hline 2 \\ \hline \end{array}.$$

Hence  $s_{\lambda/\mu} = x_1^2 + 2x_1x_2 + x_2^2$ . On the other hand, (2) in this case is:

$$s_{\lambda/\mu} = s_{1,1} + s_2 = (x_1x_2) + (x_1x_2 + x_1^2 + x_2^2).$$

By definition,

$$\begin{aligned} \text{Trop}(s_{\lambda/\mu}) &= \max\{x_1 + x_2, x_1 + x_2, 2x_1, 2x_2\} \\ &= x_1 \odot x_2 \oplus x_1 \odot x_2 \oplus x_1^{\odot 2} \oplus x_2^{\odot 2} \\ &= x_1 \odot x_2 \oplus x_1^{\odot 2} \oplus x_2^{\odot 2} \quad (\text{idempotence}) \\ &= \text{Trop}(s_2), \end{aligned}$$

in agreement with Theorem 1.3. □

The following addresses Conjecture 1.2:

**Corollary 1.5** (of Theorems 1.1 and 1.3).  *$\text{Trop}(s_{\lambda/\mu})$  has at most  $O(n^2 \cdot \lambda_1)$  tropical semiring complexity.*

In Section 2, we describe a sufficient condition for polynomial complexity. This is explained in terms of *saturated Newton polytopes* [MoToYo17]. Section 3 applies this condition to *Stanley symmetric polynomials* [St84]. Since skew-Schur polynomials are a special case of Stanley symmetric polynomials, we thereby deduce Theorem 1.3. In Section 4, we remark on how the condition applies to other families of symmetric polynomials.

## 2. DOMINANCE ORDER, NEWTON POLYTOPES AND SATURATION

Let  $\text{Par}(d) = \{\lambda : \lambda \vdash d\}$  be the set of partitions of size  $d$ . *Dominance order*  $\leq_D$  on  $\text{Par}(d)$  is defined by

$$(3) \quad \mu \leq_D \lambda \quad \text{if} \quad \sum_{i=1}^k \mu_i \leq \sum_{i=1}^k \lambda_i \quad \text{for all } k \geq 1.$$

*Definition 2.1.* Suppose  $f \in \text{Sym}_n$  is homogeneous of degree  $d$  such that

$$f = \sum_{\mu \in \text{Par}(d)} c_\mu s_\mu$$

with  $c_\mu \geq 0$  for all  $\mu$ . Moreover, assume there exists  $\lambda$  with  $c_\lambda \neq 0$  such that  $c_\mu \neq 0$  only if  $\mu \leq_D \lambda$ . Then we say  $f$  is *dominated* by  $s_\lambda$ .  $\square$

The *Newton polytope* of a polynomial  $f$  is the convex hull of its exponent vectors, so

$$\text{Newton}(f) = \text{conv}(\{(i_1, i_2, \dots, i_n) : c_{i_1, i_2, \dots, i_n} \neq 0\}) \subseteq \mathbb{R}^n.$$

C. Monical, N. Tokcan and the second author [MoToYo17] define  $f$  to have *saturated Newton polytope* (SNP) if  $c_{i_1, \dots, i_n} \neq 0$  whenever  $(i_1, \dots, i_n) \in \text{Newton}(f)$ .

The *permutahedron* of  $\lambda = (\lambda_1, \dots, \lambda_n)$ , denoted  $\mathcal{P}_\lambda$ , is the convex hull of the  $S_n$ -orbit of  $\lambda$  in  $\mathbb{R}^n$ . It follows from R. Rado's theorem [Ra52] that

$$\text{Newton}(s_\lambda) = \mathcal{P}_\lambda \quad \text{and} \quad \text{Newton}(s_\mu) \subseteq \text{Newton}(s_\lambda) \quad \text{if } \mu \leq_D \lambda.$$

A consequence of R. Rado's theorem [MoToYo17, Proposition 2.5] is therefore:

**Proposition 2.2.** *If  $f \in \text{Sym}_n$  is dominated by  $s_\lambda$ , then  $f$  is SNP, and*

$$\text{Newton}(f) = \text{Newton}(s_\lambda) = \mathcal{P}_\lambda.$$

We give a technical strengthening of [GrKo16, Theorem 2.5]:

**Proposition 2.3** (Sufficient condition for polynomial complexity). *Suppose  $f \in \text{Sym}_n$  is dominated by  $s_\lambda(x_1, \dots, x_n)$ . Then  $\text{Trop}(f) = \text{Trop}(s_\lambda)$  over the tropical semiring  $(\mathbb{R}, \oplus, \odot)$ . Therefore,  $f$  has at most  $O(n^2 \cdot \lambda_1)$  tropical semiring complexity.*

*Proof.* By Proposition 2.2,

$$(4) \quad \text{Newton}(f) = \text{Newton}(s_\lambda),$$

and  $f$  is SNP. This proves the first statement.

At this point, we can appeal to Theorem 1.1 to obtain the second claim. However, for convenience, we recall the ideas from [GrKo16, Theorem 2.5], thus indicating the underlying circuit. There it is shown that

$$(5) \quad \text{Newton}(s_\lambda)[\mathbb{Z}] = \sum_{1 \leq k \leq \lambda_1} \text{Newton}(e_{\lambda'_k})[\mathbb{Z}].$$

In the Minkowski sum of (5),

$$e_k = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} x_{j_1} \cdots x_{j_k}$$

is the *elementary symmetric polynomial* of degree  $k$ . Also,  $\lambda'$  is the conjugate partition of  $\lambda$ , obtained by transposing the Young diagram for  $\lambda$ . Finally, for a polytope  $\mathcal{P} \subseteq \mathbb{R}^n$ ,  $\mathcal{P}[\mathbb{Z}]$  denotes the set of integer lattice points of  $\mathcal{P}$ .

Combining (4) and (5), we see

$$(6) \quad \text{Newton}(f)[\mathbb{Z}] = \sum_{1 \leq k \leq \lambda_1} \text{Newton}(e_{\lambda'_k})[\mathbb{Z}].$$

By Proposition 2.2,  $f$  is SNP. This property of  $f$ , together with (6), implies

$$(7) \quad \text{Trop}(f) = \bigodot_{1 \leq k \leq \lambda_1} \text{Trop}(e_{\lambda'_k}),$$

as tropical polynomials.

Therefore, following *loc. cit.*, to calculate  $\text{Trop}(f)$  it suffices to compute  $\text{Trop}(e_{\lambda'_k})$  for  $1 \leq k \leq \lambda_1$ . The latter has at worst  $O(n^2)$  complexity, using the (tropicalization) of the Pascal-type recurrence

$$e_k(x_1, \dots, x_n) = e_k(x_1, \dots, x_{n-1}) + x_n e_{k-1}(x_1, \dots, x_{n-1}).$$

This proves the second claim. □

*Remark 2.4.* The assumption in Definition 2.1 that  $f$  be Schur-positive ( $c_\mu \geq 0$  for each  $\mu$ ) is needed for Proposition 2.2. Consider the *monomial symmetric polynomial*  $m_\lambda := \sum_{\theta} x_1^{\theta_1} \cdots x_n^{\theta_n}$ , where the sum is over distinct rearrangements of  $\lambda$ . It is true that  $m_\lambda = \sum_{\mu \leq_{\mathcal{D}} \lambda} I_{\lambda, \mu} s_\mu$ , where  $I_{\lambda, \lambda} = 1$ . Yet,  $\{m_\lambda\}$  has exponential complexity, by [GrKo16]. □

### 3. STANLEY SYMMETRIC POLYNOMIALS AND THE PROOF OF THEOREM 1.3

For any permutation  $w$ , R. P. Stanley [St84] defined the symmetric power series

$$F_w = \sum_{\mathbf{a} \in \text{Red}(w)} \sum_{\mathbf{b} \in C(\mathbf{a})} x_{\mathbf{b}}.$$

Here  $\text{Red}(w)$  is the set of reduced words for  $w$  in the simple transpositions  $s_i = (i \ i+1)$ . This means  $\mathbf{a} = (a_1, a_2, \dots, a_\ell)$ , where  $s_{a_1} s_{a_2} \cdots s_{a_\ell} = w$  and  $\ell = \ell(w)$  is the number of inversions of  $w$ . Now, if  $\mathbf{b} = (b_1, \dots, b_\ell)$ , then  $\mathbf{b} \in C(\mathbf{a})$  if

- $1 \leq b_1 \leq b_2 \leq \cdots \leq b_\ell$ ; and
- $a_i < a_{i+1} \implies b_i < b_{i+1}$ .

Set  $x_{\mathbf{b}} := x_{b_1} x_{b_2} \cdots x_{b_\ell}$ . (This actually defines  $F_{w^{-1}}$  in [St84]. Thus we use the results of *loc. cit.* with this swap of convention.)

*Remark 3.1.* The original motivation for  $F_w$  is that  $\#\text{Red}(w) = [x_1 x_2 \cdots x_\ell] F_w$ . If we define  $a_{w\lambda}$  as the coefficients in

$$(8) \quad F_w = \sum_{\lambda} a_{w\lambda} s_\lambda,$$

then  $a_{w\lambda} \in \mathbb{Z}_{\geq 0}$ . This nonnegativity is proved by work of [EdGr87] (see also [LaSc82]). In fact,  $a_{w\lambda}$  is a generalization of the Littlewood-Richardson coefficient. A theorem of

H. Narayanan [Na06] states that computation of  $c_{\lambda,\mu}^\nu$  is #P-complete in L. Valiant's complexity theory for counting problems [Va79]. Hence  $a_{w\lambda}$  is a #P-complete counting problem. In particular, this means that there is no polynomial time algorithm for computing either  $c_{\lambda,\mu}^\nu$  or  $a_{w\lambda}$  unless  $P = NP$ .

Now,  $[x_1 \dots x_\ell]_{s_\lambda} = f^\lambda$  counts standard Young tableaux of shape  $\lambda$ . These numbers are computed by the famous *hook-length formula*. The resulting enumeration  $\#\text{Red}(w) = \sum_\lambda a_{w\lambda} f^\lambda$  establishes that  $\#\text{Red}(w)$  is a #P counting problem. Is it #P-complete?  $\square$

Recall that the *Rothe diagram* of  $w$  is given by


$$D(w) = \{(i, j) : 1 \leq i, j \leq n, j < w(i), i < w^{-1}(j)\}.$$

Pictorially, this is described by placing  $\bullet$ 's in positions  $(i, w(i))$  (in matrix notation), striking out boxes below and to the right of each  $\bullet$ . Then  $D(w)$  consists of the remaining boxes.

For example, if

$$(9) \quad w = 4\ 1\ 5\ 2\ 7\ 3\ 9\ 6\ 10\ 8 \in S_{10} \quad (\text{in one line notation}),$$

then  $D(w)$  is depicted by:



For  $w \in S_m$ , set  $q_i$  to be the number of boxes of  $D(w)$  in column  $i$  (counting from the left) for  $1 \leq i \leq m$ . Then  $(q_1, q_2, \dots, q_m)$  is the *code* of  $w^{-1}$ . Let  $\beta_{\max}(w)$  be the partition obtained by sorting  $(q_1, q_2, \dots, q_m)$  in decreasing order and taking the conjugate shape.

**Theorem 3.2** (Complexity of tropical Stanley polynomials). *Let  $w \in S_m$ . Then the tropical semiring complexity of  $\text{Trop}(F_w(x_1, \dots, x_n))$  is at most  $O(n^2 \cdot \beta_{\max}(w)_1)$ .*

*Proof.* By [St84, Theorem 4.1] (up to convention), if  $a_{w,\lambda} \neq 0$ , then

$$\lambda \leq_D \beta_{\max}(w).$$

Since the  $a_{w\lambda}$  in (8) are positive, if  $F_w(x_1, \dots, x_n)$  is nonzero, then  $a_{w,\beta_{\max}(w)} \neq 0$  and  $F_w$  is dominated by  $s_{\beta_{\max}(w)}$ . Now use Proposition 2.3.  $\square$

*Proof of Theorem 1.3 (and Corollary 1.5):* We show that  $s_{\lambda/\mu}(x_1, \dots, x_n)$  is dominated by  $s_\beta(x_1, \dots, x_n)$  for some shape  $\beta$  (to be determined) with  $\beta_1 = \lambda_1$ .

Given  $\lambda/\mu$ , construct a permutation  $w_{\lambda/\mu}$  by filling all boxes in the same northwest-southeast diagonal with the same entry, starting with 1 on the northeastmost diagonal and increasing consecutively as one moves southwest. Call this filling  $T_{\lambda/\mu}$ .

For instance, if  $\lambda/\mu = (5, 4, 3, 2, 1)/(2, 2, 1, 0, 0)$  then

$$T_{\lambda/\mu} = \begin{array}{cccc} & & 3 & 2 & 1 \\ & & 4 & 3 & \\ & 6 & 5 & & \\ 8 & 7 & & & \\ 9 & & & & \end{array}$$

Let  $(r_1, r_2, \dots, r_{|\lambda/\mu|})$  be the left-to-right, top-to-bottom, row reading word of  $T_{\lambda/\mu}$ . In our example, this is  $(3, 2, 1, 4, 3, 6, 5, 8, 7, 9)$ .

Define  $w_{\lambda/\mu} = s_{r_1} s_{r_2} \cdots s_{r_{|\lambda/\mu|}}$ . By [BiJoSt93, Corollary 2.4],

$$F_{w_{\lambda/\mu}}(x_1, \dots, x_n) = s_{\lambda/\mu}(x_1, \dots, x_n) \in \text{Sym}_n.$$

By [BiJoSt93, Section 2],  $\lambda/\mu$  is obtained by removing empty rows and columns of  $D(w_{\lambda/\mu})$  and reflecting across a vertical line. In our example,  $w_{\lambda/\mu}$  is the permutation  $(9)$ . The reader can check from the Rothe diagram that this process gives  $\lambda/\mu$ .

By definition,  $\beta_{\max}(w_{\lambda/\mu})$  is the conjugate of the decreasing rearrangement of the code of  $w_{\lambda/\mu}^{-1}$ . Hence, in our example, the code of  $w_{\lambda/\mu}^{-1}$  is  $(1, 2, 3, 0, 0, 2, 0, 2, 0, 0)$ , which rearranges to  $(3, 2, 2, 2, 1, 0, 0, 0, 0, 0)$ . Therefore,

$$\beta_{\max}(w_{\lambda/\mu}) = \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & \square & \square \end{array}' = \begin{array}{cccc} \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \\ \square & \square & \square & \square \end{array}.$$

(Thus,  $\beta_{\max}(w_{\lambda/\mu})$  is obtained from  $\lambda/\mu$  by first pushing the boxes in each column north and left-justifying the result.)

Since the coefficients  $c_{\mu, \nu}^\lambda$  in the Schur expansion (2) are positive,  $s_{\lambda/\mu}(x_1, \dots, x_n)$  is dominated by  $s_\beta$ , where  $\beta = \beta_{\max}(w_{\lambda/\mu})$ . By the above process from [BiJoSt93] relating  $D(w_{\lambda/\mu})$  and  $\lambda/\mu$ ,

$$\beta_1 = \beta_{\max}(w_{\lambda/\mu})_1 = \lambda_1,$$

as desired. Theorem 1.3 now holds by the first conclusion of Proposition 2.3.

To conclude Corollary 1.5, we may now either apply Theorem 1.1 or Theorem 3.2.  $\square$

#### 4. SOME OTHER SYMMETRIC POLYNOMIALS

In [MoToYo17, Sections 2 and 3], some symmetric polynomials are observed to be SNP because they are dominated by  $s_\lambda$  (for some  $\lambda$ ), or for other reasons. These include:

- (1) J. R. Stembridge's polynomial  $F_M$  for a totally nonnegative matrix  $M$  [St91];
- (2) the cycle index polynomial  $c_G$  of a subgroup  $G \leq S_n$  (from Redfield-Pólya theory);
- (3) C. Reutenauer's  $q_\lambda$  basis of  $\text{Sym}_n$  [Re95];
- (4) the symmetric Macdonald polynomial (where  $(q, t) \in \mathbb{C}^2$  is generic);
- (5) the Hall-Littlewood polynomial (for any positive evaluation of  $t$ ); and
- (6) Schur  $P$ - and Schur  $Q$ - polynomials.

Consequently, by Proposition 2.3, the tropicalizations of these polynomials equal some tropical Schur polynomial. Therefore, as with skew Schur polynomials, one obtains immediate tropical semiring complexity implications:

- The polynomials (1) and (2) are dominated by  $s_{(n)}$ ; see [MoToYo17, Theorem 2.28] and [MoToYo17, Theorem 2.30]. Hence Proposition 2.3 shows their tropicalizations have  $O(n^3)$  complexity. In fact, since  $\text{Trop}(s_{(n)}) = \text{Trop}((x_1 + \cdots + x_n)^n)$  as tropical polynomials, they have  $O(n)$  complexity (see [FoGrNoSc16, Theorem 1.4] for a nontropical version of this statement).
- For (3), if  $\lambda = (\lambda_1, \dots, \lambda_\ell, 1^r)$  where each  $\lambda_i \geq 2$ , then  $q_\lambda$  is dominated by  $s_{a,b}$  where  $a = |\lambda| - \ell$  and  $b = \ell$  (see [MoToYo17, Theorem 2.3.2] and specifically its proof). Hence Proposition 2.3 asserts  $\text{Trop}(q_\lambda)$  has  $O(n^2 \cdot (|\lambda| - \ell))$  complexity.
- For a generic choice of  $q, t \in \mathbb{C}^2$ , it follows from [MoToYo17, Section 3.1] that if  $P_\lambda(X; q, t) \in \text{Sym}_n$  is the Macdonald polynomial, then  $\text{Trop}(P_\lambda(X; q, t)) = \text{Trop}(s_\lambda)$ . Hence  $\text{Trop}(P_\lambda(X; q, t))$  has  $O(n^2 \cdot \lambda_1)$  complexity.
- (5) and (6) are also indexed by partitions  $\lambda$  and dominated by  $s_\lambda$ . Thus, Proposition 2.3 implies their tropicalizations have  $O(n^2 \cdot \lambda_1)$  complexity.

#### ACKNOWLEDGEMENTS

AY was supported by an NSF grant. We thank Marc Snir for helpful communications.

#### REFERENCES

- [BiJoSt93] S. Billey, W. Jockusch and R. P. Stanley, *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. 2(1993), no. 4, 345–374.
- [EdGr87] P. Edelman and C. Greene, *Balanced tableaux*, Adv. in Math. 63 (1987), no. 1, 42–99.
- [FoGrNoSc16] S. Fomin, D. Grigoriev, D. Nogneng and E. Schost, *On semiring complexity of Schur polynomials*, preprint, 2016. arXiv:1608.05043
- [Fu97] W. Fulton, *Young tableaux. With applications to representation theory and geometry*. London Mathematical Society Student Texts, 35. Cambridge University Press, Cambridge, 1997.
- [GrKo16] D. Grigoriev and G. Koshevoy, *Complexity of tropical Schur polynomials*, Journal of Symbolic Computation, Volume 74, May–June 2016, Pages 46–54.
- [ItMiSh09] I. Itenberg, G. Mikhalkin, and E. Shustin, *Tropical algebraic geometry*. Second edition. Oberwolfach Seminars, 35. Birkhuser Verlag, Basel, 2009. x+104 pp.
- [JeSn82] M. Jerrum and M. Snir, *Some exact complexity results for straight-line computations over semirings*, J. Association for Computing Machinery, Vol 29, No. 3, July 1982, 874–897.
- [LaSc82] A. Lascoux and M. -P. Schützenberger, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. 295 (1982), 629–633.
- [MaSt15] D. Maclagan and B. Sturmfels, *Introduction to tropical geometry*. Graduate Studies in Mathematics, 161. American Mathematical Society, Providence, RI, 2015. xii+363 pp.
- [MoToYo17] C. Monical, N. Tokcan and A. Yong. *Newton polytopes in algebraic combinatorics*. Preprint, arXiv:1703.02583
- [Na06] H. Narayanan, *On the complexity of computing Kostka numbers and Littlewood-Richardson coefficients*, J. Alg. Comb., Vol. 24, N. 3, 2006, 347–354.
- [Ra52] R. Rado. *An inequality*. J. London Math. Soc., 27:1–6, 1952.
- [Re95] C. Reutenauer, *On symmetric functions, related Witt vectors, and the free Lie algebra*, Adv. Math. 110(1995), 234–246.
- [St84] R. P. Stanley, *On the number of reduced decompositions of elements of Coxeter groups*, European J. Combin. 5 (1984), no. 4, 359–372.

- [St99] ———, Enumerative combinatorics. Vol. 2. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. Cambridge Studies in Advanced Mathematics, 62. Cambridge University Press, Cambridge, 1999.
- [St91] J. R. Stembridge, *Immanants of totally positive matrices are nonnegative*, Bulletin of the London Mathematical Society, 23 (1991), no. 5, 422–428.
- [Va79] L. G. Valiant, *The complexity of computing the permanent*, Theoret. Comput. Sci., 8(2):189–201, 1979.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IDAHO, MOSCOW, ID 83844

*E-mail address:* awoo@uidaho.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801

*E-mail address:* ayong@uiuc.edu