

RAMANUJAN REACHES HIS HAND FROM HIS GRAVE TO SNATCH YOUR THEOREMS FROM YOU

BRUCE C. BERNDT

1. INTRODUCTION

Born on December 22, 1887, India's greatest mathematician, Srinivasa Ramanujan, began to record his discoveries in notebooks in about 1904 when he entered the Government College of Kumbakonam for what was to be only one year of study. For the next five years, Ramanujan did mathematics, mostly in isolation, while logging his findings without proofs in notebooks. Ramanujan moved to Madras in 1910 and, while working as a clerk in the Madras Port Trust Office, was encouraged by the Manager and Chief Accountant, S. Narayana Aiyar, and the Chairman, Sir Francis Spring, to write English mathematicians about his work. After communicating about 120 of his theorems to G. H. Hardy in early 1913, Ramanujan accepted Hardy's invitation to come to Cambridge, and so on March 17, 1914, Ramanujan departed India for England. At about that time, from letters that he wrote to friends back home in Madras [12, pp. 112–113; 123–125], Ramanujan ceased recording his theorems in notebooks to concentrate on publishing research that he was conducting in England. After returning to India in March, 1919, Ramanujan began to log entries in what was later to be called *Ramanujan's Lost Notebook*, which was found by George Andrews in the library at Trinity College, Cambridge in March, 1976. After a lengthy illness that confined him to nursing homes during his last two years in England, Ramanujan died on April 26, 1920 at the age of 32.

Ramanujan's earlier notebooks were not published until 1957 [25], and his later lost notebook was not published until 1988 [26]. Thus, not surprisingly, it transpired that some of Ramanujan's discoveries were hidden for many years and were in the meantime proved by others.

On June 1–5, 1987, a meeting was held at the University of Illinois to commemorate the centenary of the birth of Ramanujan and to survey the many areas of mathematics (and of physics) that have been profoundly influenced by his work. One of the speakers, R. William Gosper, remarked in his lecture, "How can you love this man? He continually reaches his hand from his grave to snatch your theorems from you." The purpose of this paper is to provide just a few of the many examples for which others have received credit for theorems, but unknown to them, their discoveries were not original with them; they were first unearthed by Ramanujan.

2. RAMANUJAN'S EARLY WORK

A perusal of issues of the *Journal of the Indian Mathematical Society* from its inception in 1907 to, say, the late 1920's shows that some of the earlier papers and problems that Ramanujan submitted to the *Journal of the Indian Mathematical Society* were much in the spirit

of mathematics popular in India at that time. Thus, it is not unexpected that some articles published in the *Journal* in the decade after Ramanujan's death might contain entries found many years later to be in Ramanujan's notebooks. In particular, two papers by M. B. Rao and M. V. Ayyar [27] and one by S. L. Malurkar [22] contain several entries from Chapter 14 in Ramanujan's second notebook [25], [7]. We offer one example from a paper by Malurkar, who studied physics at Cambridge University and who was Director of the observatories in Pune and Mumbai for much of his career.

Theorem 2.1. *Let $\zeta(s) := \sum_{n=1}^{\infty} n^{-s}$, $\operatorname{Re} s > 1$, denote the Riemann zeta function, and let B_n , $n \geq 0$, denote the n th Bernoulli number. If α and β are positive numbers such that $\alpha\beta = \pi^2$, and if r is a positive integer, then*

$$\begin{aligned} & (4\alpha)^{-r} \left(\frac{1}{2}\zeta(2r+1) + \sum_{m=1}^{\infty} \frac{1}{m^{2r+1}(e^{2m\alpha} - 1)} \right) \\ & - (-4\beta)^{-r} \left(\frac{1}{2}\zeta(2r+1) + \sum_{m=1}^{\infty} \frac{1}{m^{2r+1}(e^{2m\beta} - 1)} \right) \\ & = - \sum_{k=0}^{r+1} \frac{(-1)^k B_{2k} B_{2r+2-2k} \alpha^{r+1-k} \beta^k}{(2k)!(2r+2-2k)!}. \end{aligned} \tag{2.1}$$

Theorem 2.1 can be found as Entry 21(i) in Chapter 14 in Ramanujan's second notebook [7, pp. 275–276]. It also appears in a manuscript of Ramanujan that was published for the first time in its original handwritten form with his lost notebook [26, pp. 319–320, formula (28)]. This manuscript, along with commentaries, can also be found in [10] and [1]. The special case $\alpha = \beta = \pi$ was first proved by M. Lerch in 1901 [21]. There are now several proofs of Theorem 2.1, and references to these many proofs can be found in [7, p. 276] and [1]. An extensive generalization of Entry 2.1 can be found in Entry 20 of Chapter 16 in Ramanujan's first notebook [25], [8, pp. 429–432], and another one in [9]. Moreover, there are further generalizations, including analogues for Dirichlet L -functions.

Ramanujan's formula (2.1) is fascinating for several reasons. It is well known that, for each positive integer n , $\zeta(2n)$ can be explicitly evaluated as a rational multiple of π^{2n} containing the factor B_{2n} . However, except for the fact that $\zeta(3)$ is irrational, which was first proved by R. Apéry [2], we currently know no further information about the arithmetical nature of $\zeta(2n+1)$. If we set $r = 2n+1$, $n \geq 0$, and $\alpha = \beta = \pi$ in (2.1), we find that $\zeta(4n+3)$ can be represented as a rational multiple of π^{4n+3} involving Bernoulli numbers plus a rapidly convergent series. Thus, $\zeta(4n+3)$ is “almost” a rational multiple of π^{4n+3} .

3. BELL NUMBERS AND POLYNOMIALS

Most readers will be familiar with the name of E. T. Bell because of his popular book [5]. However, those with a combinatorial interest will identify him with Bell numbers and Bell polynomials. However, these numbers and polynomials should have Ramanujan's name attached to them, and not Bell's, because Ramanujan extensively studied these numbers and polynomials in Chapter 3 of his second notebook [25], [6], probably over 30 years before Bell wrote his papers [3], [4] on these numbers and polynomials.

The Bell polynomials $\varphi_n(x)$, $n \geq 0$, are defined by

$$\varphi_0(x) \equiv 1, \quad e^x \varphi_{n+1}(x) := \sum_{k=1}^{\infty} \frac{k^n x^k}{(k-1)!}, \quad n \geq 0.$$

Readers should verify that $\varphi_n(x)$ is a polynomial of degree n , and, in particular,

$$\begin{aligned} \varphi_1(x) &= x, \\ \varphi_2(x) &= x + x^2, \\ \varphi_3(x) &= x + 3x^2 + x^3, \\ \varphi_4(x) &= x + 7x^2 + 6x^3 + x^4. \end{aligned}$$

They can be generated by the exponential generating function

$$e^{x(e^a-1)} = \sum_{n=0}^{\infty} \frac{a^n}{n!} \varphi_n(x).$$

The n th Bell number $B(n)$, $n \geq 1$, is defined by

$$B(n) = \varphi_n(1).$$

For example,

$$B(1) = 1, \quad B(2) = 2, \quad B(3) = 5, \quad B(4) = 15, \quad B(5) = 52, \quad B(6) = 203.$$

Combinatorially, $B(n)$ is equal to the number of ways of partitioning a set of n objects into subsets. In early editions of the Japanese classic, *The Tale of Genji*, written by Lady Shikibu Murasaki early in the 11th century, an arrangement of five incense sticks can be found at the beginning of each of the 54 chapters, except for the first and last. Thus, the author knew that the 5th Bell number equals 52. Readers enjoying poetry might like to know that the number of ways of rhyming a sonnet is $B(14) = 190,899,322$.

4. KOSHLIAKOV'S FORMULA AND GUINAND'S FORMULA

The Russian mathematician, N. S. Koshliakov (1891–1958), is chiefly remembered by a formula that now bears his name [20]. However, most of his work has been unfortunately neglected, and consequently his contributions to mathematics under appreciated. In 1942, he was arrested on fabricated charges and sent to a labor camp in the Ural mountains. Classified as an invalid after suffering from complete exhaustion, he found time to do mathematics, often under extremely difficult conditions, before being released in 1949.

Theorem 4.1 (Koshliakov's Formula). *Let $K_0(z)$ denote the modified Bessel function of order 0, let $d(n)$ denote the number of positive divisors of the positive integer n , and let γ denote Euler's constant. If α and β are positive numbers such that $\alpha\beta = 1$, then*

$$\begin{aligned} & \sqrt{\alpha} \left(\frac{\gamma - \log(4\pi\alpha)}{\alpha} - 4 \sum_{n=1}^{\infty} d(n) K_0(2\pi n\alpha) \right) \\ &= \sqrt{\beta} \left(\frac{\gamma - \log(4\pi\beta)}{\beta} - 4 \sum_{n=1}^{\infty} d(n) K_0(2\pi n\beta) \right). \end{aligned} \quad (4.1)$$

Why is Koshliakov's formula interesting? Recall that the theta transformation formula

$$\sum_{n=-\infty}^{\infty} e^{-\pi n^2/\tau} = \sqrt{\tau} \sum_{n=-\infty}^{\infty} e^{-\pi n^2 \tau}, \quad \operatorname{Re} \tau > 0, \quad (4.2)$$

is the most common route to the functional equation of the Riemann zeta function

$$\pi^{-s/2} \Gamma\left(\frac{1}{2}s\right) \zeta(s) = \pi^{-(1-s)/2} \Gamma\left(\frac{1}{2}(1-s)\right) \zeta(1-s). \quad (4.3)$$

In fact, (4.2) and (4.3) are equivalent. Koshliakov's formula (4.1) is an analogue of (4.2), and, as shown by W. L. Ferrar [16] and F. Oberhettinger and K. L. Soni [24], is equivalent to the functional equation for $\zeta^2(s) = \sum_{n=1}^{\infty} d(n)n^{-s}$, which, of course, is obtained by squaring both sides of the functional equation (4.3).

Koshliakov published his proof of Theorem 4.1 in 1929 [20]. However, his formula can be found on page 253 of the publication containing the lost notebook [26], which was written in the last year of Ramanujan's life 1919–1920. The volume [26] contains further partial manuscripts and fragments written by Ramanujan; page 253 is such a page, and so it is possible that Ramanujan proved the formula earlier than 1919. Ramanujan clearly derived Koshliakov's formula from "Guinand's formula," which can be found on the same page, and which was first proved in print by A. P. Guinand [19] in 1955. Thus, on a single page, Ramanujan reached up his hand to snatch formulas that had brought fame to two mathematicians.

Theorem 4.2 (Guinand's Formula). *Let $\sigma_k(n) = \sum_{d|n} d^k$, let $\zeta(s)$ denote the Riemann zeta function, and let $K_\nu(z)$ denote the modified Bessel function of order ν . If α and β are positive numbers such that $\alpha\beta = \pi^2$, and if s is any complex number, then*

$$\begin{aligned} & \sqrt{\alpha} \sum_{n=1}^{\infty} \sigma_{-s}(n) n^{s/2} K_{s/2}(2n\alpha) - \sqrt{\beta} \sum_{n=1}^{\infty} \sigma_{-s}(n) n^{s/2} K_{s/2}(2n\beta) \\ &= \frac{1}{4} \Gamma\left(\frac{s}{2}\right) \zeta(s) \{\beta^{(1-s)/2} - \alpha^{(1-s)/2}\} + \frac{1}{4} \Gamma\left(-\frac{s}{2}\right) \zeta(-s) \{\beta^{(1+s)/2} - \alpha^{(1+s)/2}\}. \end{aligned} \quad (4.4)$$

Since $K_s(z) = K_{-s}(z)$, we see that (4.4) is invariant under the transformation $s \rightarrow -s$.

5. SNATCHING FROM GUINAND AGAIN

Ramanujan's lost notebook contains at least three further results that were first proved in print by Guinand. In this section we examine another beautiful transformation formula, thought first to have been established by Guinand [17], but found ensconced in a two-page partial manuscript published with the lost notebook [26, p. 220]. For further examples and details, see the forthcoming volume [1] by Andrews and the author.

To state this aforementioned formula, we require some definitions. Set

$$\psi(z) := \frac{\Gamma'(z)}{\Gamma(z)} \sim \log z - \frac{1}{2z} - \frac{1}{12z^2} + \frac{1}{120z^4} - \frac{1}{252z^6} + \cdots, \quad (5.1)$$

as $z \rightarrow \infty$, $|\arg z| < \pi$. Next, Riemann's ξ -function is defined by

$$\xi(s) := (s-1)\pi^{-s/2}\Gamma\left(1+\frac{1}{2}s\right)\zeta(s)$$

while the Riemann Ξ -function is given by

$$\Xi(t) := \xi\left(\frac{1}{2} + it\right).$$

Theorem 5.1. *Define*

$$\phi(x) := \psi(x) + \frac{1}{2x} - \log x.$$

If α and β are positive numbers such that $\alpha\beta = 1$, then

$$\begin{aligned} \sqrt{\alpha} \left\{ \frac{\gamma - \log(2\pi\alpha)}{2\alpha} + \sum_{n=1}^{\infty} \phi(n\alpha) \right\} &= \sqrt{\beta} \left\{ \frac{\gamma - \log(2\pi\beta)}{2\beta} + \sum_{n=1}^{\infty} \phi(n\beta) \right\} \\ &= -\frac{1}{\pi^{3/2}} \int_0^{\infty} \left| \Xi\left(\frac{1}{2}t\right) \Gamma\left(\frac{-1+it}{4}\right) \right|^2 \frac{\cos\left(\frac{1}{2}t \log \alpha\right)}{1+t^2} dt. \end{aligned} \quad (5.2)$$

Note that the asymptotic formula (5.1) ensures the convergence of the infinite series in (5.2). The appearance of the Riemann zeta function on the far right side of (5.2) is unexpected. Ramanujan's recording of (5.2) is given as formula (7) on page 220 of [26] amidst several examples of Fourier sine and cosine transforms. In particular, formula (3) on that page is given as

$$\int_0^{\infty} (\psi(1+x) - \log x) \cos(2\pi nx) dx = \frac{1}{2} (\psi(1+n) - \log n), \quad (5.3)$$

i.e., $\psi(1+x) - \log x$ is a self-reciprocal Fourier cosine transform. Guinand evidently discovered the first equality in (5.2) in the course of reading page proofs for [17], for he gives the identity in a footnote at the end of his paper. Furthermore, he writes that it “can be deduced from” (5.3). It is clear that both Ramanujan and Guinand employed (5.3) in their proofs. Referring to (5.2), Guinand asserts that “This formula also seems to have been overlooked.” In [18], Guinand provides another proof of the first part of (5.2) that also depends on (5.3), about which he writes, “Professor T. A. Brown tells me that he proved the self-reciprocal property of $\psi(1+x) - \log x$ some years ago, and that he communicated the result to the late Professor G. H. Hardy. Professor Hardy said that the result was also given in a progress report to the University of Madras by S. Ramanujan, but was not published elsewhere.” In March, 1913, Ramanujan received a scholarship from the University of Madras, with the only requirement being that he had to write quarterly reports about his research. Three such reports were written before he departed for England in March, 1914. Hardy's memory was perhaps flawed, because (5.3) cannot be found in Ramanujan's quarterly reports, which are detailed and discussed in [6], although it can be found in a partial manuscript [26, pp. 219–220], which evidently was also in Hardy's possession. The second equality in (5.2) was first proved by the author and A. Dixit [11], who gave two proofs of the first equality. Moreover, Dixit [14], [15] has established generalizations and analogues of (5.2).

6. DICKMAN'S FUNCTION

Dickman's function $\rho(u)$, first introduced by K. Dickman in 1930 [13], plays a central role in prime number theory. For $0 \leq u \leq 1$, let $\rho(u) \equiv 1$. For each integer $k \geq 1$, $\rho(u)$ is

defined inductively for $k \leq u \leq k + 1$ by

$$\rho(u) = \rho(k) - \int_k^u \rho(v-1) \frac{dv}{v}.$$

Dickman's function is continuous at $u = 1$ and differentiable for $u > 1$. Equivalently, $\rho(u)$ can be defined by the differential-difference equation

$$u\rho'(u) + \rho(u-1) = 0, \quad u > 1.$$

Let $P^+(n)$ denote the largest prime factor of the positive integer n , and set

$$\Psi(x, y) := |\{n \leq x : P^+(n) \leq y\}|. \quad (6.1)$$

On page 337 in his lost notebook [26], Ramanujan offers several asymptotic formulas for $\Psi(x, x^\epsilon)$, although in a different language. We quote just one instance. "Let $\phi(x)$ denote the number of numbers of the form

$$2^{a_2} 3^{a_3} 5^{a_5} \cdots p^{a_p}, \quad p \leq x^\epsilon,$$

not exceeding x . Then, for $\frac{1}{2} \leq \epsilon \leq 1$,

$$\phi(x) \sim x \left\{ 1 - \int_\epsilon^1 \frac{du_0}{u_0} \right\}." \quad (6.2)$$

His next formula is for $\frac{1}{3} \leq \epsilon \leq \frac{1}{2}$, and so on. In the notation (6.1), Ramanujan's function $\phi(x) = \Psi(x, x^\epsilon)$. Ramanujan hence proved Dickman's [13] famous asymptotic formula

$$\Psi(x, x^{1/u}) \sim x\rho(u), \quad x \rightarrow \infty,$$

while, in fact, giving a representation for $\rho(u)$ in terms of integrals, as in the first instance (6.2). According to the author's colleague, A. J. Hildebrand, Ramanujan's theorem is equivalent to a folklore theorem, which we cannot find located in the literature.

Theorem 6.1. *Define, for $u \geq 0$,*

$$I_0(u) := 1, \quad I_k(u) := \int \cdots \int_{\substack{t_1, \dots, t_k \geq 1 \\ t_1 + \cdots + t_k \leq u}} \frac{dt_1 \cdots dt_k}{t_1 \cdots t_k}, \quad k \geq 1.$$

Then, for $u \geq 0$,

$$\rho(u) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} I_k(u). \quad (6.3)$$

The series on the right-hand side of (6.3) is finite, since if $k > u$, then $I_k(u) = 0$, for the conditions $t_1, \dots, t_k \geq 1$ and $t_1 + \cdots + t_k \leq u$ are vacuous in this case. If we make the changes of variable $\epsilon = 1/u$ and $u_j = \epsilon t_j = t_j/u$, $1 \leq j \leq k$, then we obtain Ramanujan's theorem, the first instance of which is given in (6.2).

Excellent sources for information on the Dickman function and its prominence in prime number theory are G. Tenenbaum's treatise [28, Chapter III.5] and P. Moree's dissertation [23].

REFERENCES

- [1] G. E. Andrews and B. C. Berndt, *Ramanujan's Lost Notebook*, Part IV, Springer, New York, to appear.
- [2] R. Apéry, *Interpolation de fractions continues et irrationalité de certaines constantes*, Bull. Sect. Sci. III, Bibliothèque Nationale, Paris, 1981, 37–63.
- [3] E. T. Bell, *Exponential polynomials*, Ann. Math. **35** (1934), 258–277.
- [4] E. T. Bell, *The iterated exponential integrals*, Ann. Math. **39** (1938), 539–557.
- [5] E. T. Bell, *Men of Mathematics*, Simon and Schuster, New York, 1937.
- [6] B. C. Berndt, *Ramanujan's Notebooks*, Part I, Springer-Verlag, New York, 1985.
- [7] B. C. Berndt, *Ramanujan's Notebooks*, Part II, Springer-Verlag, New York, 1989.
- [8] B. C. Berndt, *Ramanujan's Notebooks*, Part IV, Springer-Verlag, New York, 1994.
- [9] B. C. Berndt, *Dedekind sums and a paper of G. H. Hardy*, J. London Math. Soc. (2) **13** (1976), 129–137.
- [10] B. C. Berndt, *An unpublished manuscript of Ramanujan on infinite series identities*, J. Ramanujan Math. Soc. **19** (2004), 57–74.
- [11] B. C. Berndt and A. Dixit, *A transformation formula involving the gamma and Riemann zeta functions in Ramanujan's lost notebook*, in *The Legacy of Alladi Ramakrishnan in the Mathematical Sciences*, K. Alladi, J. Klauder, and C. R. Rao, eds., Springer, New York, 2010, pp. 199–210.
- [12] B. C. Berndt and R. A. Rankin, *Ramanujan: Letters and Commentary*, American Mathematical Society, Providence, RI, 1995; London Mathematical Society, London, 1995.
- [13] K. Dickman, *On the frequency of numbers containing prime factors of a certain relative magnitude*, Arkiv Mat. Astron. Fys. **22** (1930), 1–14.
- [14] A. Dixit, *Analogues of a transformation formula of Ramanujan*, Internat. J. Number Thy., to appear.
- [15] A. Dixit, *Transformation formulas associated with integrals involving the Riemann Xi-function*, Monatsh. für Math., to appear.
- [16] W. L. Ferrar, *Some solutions of the equation $F(t) = F(t^{-1})$* , J. London Math. Soc. **11** (1936), 99–103.
- [17] A. P. Guinand, *Some formulae for the Riemann zeta-function*, J. London Math. Soc. **22** (1947), 14–18.
- [18] A. P. Guinand, *A note on the logarithmic derivative of the Gamma function*, Edinburgh Math. Notes **38** (1952), 1–4.
- [19] A. P. Guinand, *Some rapidly convergent series for the Riemann ξ -function*, Quart. J. Math. (Oxford) **6** (1955), 156–160.
- [20] N. S. Koshliakov, *On Voronoi's sum-formula*, Mess. Math. **58** (1929), 30–32.
- [21] M. Lerch, *Sur la fonction $\zeta(s)$ pour valeurs impaires de l'argument*, J. Sci. Math. Astron. pub. pelo Dr. F. Gomes Teixeira, Coimbra **14** (1901), 65–69.
- [22] S. L. Malurkar, *On the application of Herr Mellin's integrals to some series*, J. Indian Math. Soc. **16** (1925/26), 130–138.
- [23] P. Moree, *Psixyology and Diophantine Equations*, Ph. D. Thesis, Leiden, 1993.
- [24] F. Oberhettinger and K. L. Soni, *On some relations which are equivalent to functional equations involving the Riemann zeta function*, Math. Z. **127** (1972), 17–34.
- [25] S. Ramanujan, *Notebooks* (2 volumes), Tata Institute of Fundamental Research, Bombay, 1957.
- [26] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Narosa, New Delhi, 1988.
- [27] M. B. Rao and M. V. Ayyar, *On some infinite series and products*, Parts I, II, J. Indian Math. Soc. **15** (1923/24), 150–162, 233–247.
- [28] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, Cambridge University Press, Cambridge, 1995.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN STREET, URBANA, IL 61801, USA

E-mail address: berndt@illinois.edu