

---

# Contents

Preface	vii
Chapter 1. Elementary Approaches to Partitions and Their Generating Functions	1
§1.1. Texts in the Theory of Partitions and Basic Notation	1
§1.2. Definitions and Generating Functions	4
§1.3. A Proof Due to Sun Kim [73]	16
§1.4. Exercises	19
Chapter 2. MacMahon's Partition Analysis; Gaussian Binomial Coefficients	21
§2.1. MacMahon's Partition Analysis	21
§2.2. Elementary Partition Identities Involving Gaussian Binomial Coefficients	27
§2.3. Exercises	30
Chapter 3. Some Primary Theorems in the Theory of $q$ -Series	31
§3.1. Introduction	31
§3.2. Some Standard Theorems about ${}_2\phi_1(a, b; c; q; z)$	40
§3.3. The More General ${}_{r+1}\phi_r$	44
§3.4. Exercises	52
Chapter 4. Partition Identities Arising from $q$ -Series Identities	53
§4.1. Theorems Arising from Basic $q$ -Series Theorems	53
§4.2. Weak Form of Bailey's Lemma	64
§4.3. Strong Form of Bailey's Lemma	68
§4.4. Applications of Bailey's Lemma	70
§4.5. Exercises	72
Chapter 5. Analogues of Euler's Theorem; Partitions with Gaps	73

---

§5.1. Motivation	73
§5.2. The Rogers–Ramanujan Continued Fraction	93
§5.3. Exercises	102
Chapter 6. The Frobenius and Generalized Frobenius Symbols	103
§6.1. First Generalization of Frobenius Partitions	103
§6.2. Second Generalization of Frobenius Partitions	106
Chapter 7. Congruences for $p(n)$	109
§7.1. Introduction	109
§7.2. Ramanujan’s Congruence $p(5n + 4) \equiv 0 \pmod{5}$	110
§7.3. Ramanujan’s Congruence $p(7n + 5) \equiv 0 \pmod{7}$	118
§7.4. Ramanujan’s Congruence $p(11n + 6) \equiv 0 \pmod{11}$	124
§7.5. A More General Partition Function	129
§7.6. Exercises	133
Bibliography	135

---

# Preface

These lecture notes are based on a course in the theory of partitions given by Bruce Berndt at the University of Illinois at Urbana-Champaign in the spring of 2014. The notes are not in polished form and need to be reorganized. The lecturer had intended to give more applications to partitions from Ramanujan's lost notebook, in particular, from mock theta functions. However, time did not permit the inclusion of these topics.

Because the notes were prepared somewhat hastily while the course was being offered, misprints are likely abundant. Please send any comments and corrections to Bruce Berndt, [berndt@illinois.edu](mailto:berndt@illinois.edu).



---

## Chapter 1

# Elementary Approaches to Partitions and Their Generating Functions

### 1.1. Texts in the Theory of Partitions and Basic Notation

The texts that have been devoted to the theory of partitions are generally well written. The contents of each of these books moderately overlaps with these notes. However, these notes contain material that is not found in any of these texts. Two of them are difficult to obtain, namely, *Classical Partition Identities and Basic Hypergeometric Series* by W. Chu and L. Di Claudio [48], and *Partition Theory* by A.K. Agarwal, Padmavathamma, and M. V. Subbarao [1]. The most famous and broadest of books devoted to partition theory is *The Theory of Partitions* by G. E. Andrews [11], and the most elementary text is *Integer Partitions* by Andrews and K. Eriksson [20]. The latter is suitable for bright high school students and undergraduates, because it requires little background. The author's book, *Number Theory in the Spirit of Ramanujan* [29], as the title indicates, is a broad introduction to some of contributions of Ramanujan to number theory, and in particular, to some of his discoveries about partitions. The emphasis in this book, not unlike the book by Chu and Di Claudio, is partition theory from the viewpoint of  $q$ -series. Not surprisingly, there is some overlap with the lecturer's book [29] and these notes. As in [29], contributions of Ramanujan are emphasized in these notes. Also, like in [29],  $q$ -series are at the heart of our approach. Entries from Ramanujan's lost notebook [92] will also be discussed.

Little background in  $q$ -series is needed to read these notes. Most of our needs in  $q$ -series are developed *ab initio*, although in a few instances we may quote results from [11], [29], or G. Gasper and M. Rahman's 'bible' of the subject [59]. You may not know

what a  $q$ -series is, and indeed a general definition of a  $q$ -series may be difficult to give, but nonetheless we shall give an admittedly vague definition below.

For any nonnegative integer  $n$  and complex number  $a$ , set

$$(a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k). \quad (1.1.1)$$

If  $n = 0$ , the empty product above is interpreted to be equal to 1. For any complex numbers  $a$  and  $q$ ,  $|q| < 1$ , define

$$(a; q)_\infty := \lim_{n \rightarrow \infty} (a; q)_n = \prod_{k=0}^{\infty} (1 - aq^k).$$

If the base  $q$  is understood, then we often abbreviate the notation by writing  $(a)_n := (a; q)_n$  and  $(a)_\infty := (a; q)_\infty$ . In later chapters, it will be convenient to use another abbreviated notation

$$(a_1, a_2, \dots, a_m; q)_n := (a_1; q)_n (a_2; q)_n \cdots (a_m; q)_n. \quad (1.1.2)$$

Similarly,

$$(a_1, a_2, \dots, a_m; q)_\infty := (a_1; q)_\infty (a_2; q)_\infty \cdots (a_m; q)_\infty. \quad (1.1.3)$$

We also occasionally use the notation

$$[a]_\infty := [a; q]_\infty := (a; q)_\infty (q/a; q)_\infty \quad (1.1.4)$$

and

$$[a_1, a_2, \dots, a_m]_\infty := [a_1, a_2, \dots, a_m; q]_\infty := [a_1; q]_\infty [a_2; q]_\infty \cdots [a_m; q]_\infty. \quad (1.1.5)$$

A  $q$ -series generally will have summands containing various products of the sort  $(a; q)_n$ . For example,

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n^2} \quad \text{and} \quad \sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q; q)_n^2}$$

As we shall see later in this chapter, the first series is a generating function for the partition function, while the latter is one of Ramanujan's mock theta functions. Theta functions frequently appear in the theory of  $q$ -series. They do not contain  $q$ -products in their summands, but because of their ubiquitous appearances in the theory of  $q$ -series, we consider theta functions as  $q$ -series as well. We use Ramanujan's general definition of a theta function, namely,

$$f(a, b) := \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \quad |ab| < 1. \quad (1.1.6)$$

In particular, use is often made of the Jacobi triple product identity [27, p. 35, Entry 19], [29, Theorem 1.3.3, p. 10]

$$f(a, b) = (-a; ab)_\infty (-b; ab)_\infty (ab; ab)_\infty, \quad (1.1.7)$$

for which several proofs will be given in the sequel.

Further useful properties are given in the following two lemmas from Ramanujan's second notebook [91], [27, p. 34, Entry 18(iv); p. 48, Entry 31].

**Lemma 1.1.1.** *We have*

$$f(a, b) = f(b, a), \quad (1.1.8)$$

$$f(1, a) = 2f(a, a^3), \quad (1.1.9)$$

$$f(-1, a) = 0, \quad (1.1.10)$$

and, if  $n$  is any integer,

$$f(a, b) = a^{n(n+1)/2} b^{n(n-1)/2} f(a(ab)^n, b(ab)^{-n}). \quad (1.1.11)$$

A proof of Lemma 1.1.1 is left for the exercises.

**Lemma 1.1.2.** *Let  $U_n = a^{n(n+1)/2} b^{n(n-1)/2}$  and  $V_n = a^{n(n-1)/2} b^{n(n+1)/2}$ . Then, for any positive integer  $n$ ,*

$$f(a, b) = \sum_{r=0}^{n-1} U_r f\left(\frac{U_{n+r}}{U_r}, \frac{V_{n-r}}{U_r}\right). \quad (1.1.12)$$

Ramanujan actually wrote Lemma 1.1.2 in the form

$$\begin{aligned} f(U_1, V_1) &= f(U_n, V_n) + U_1 f\left(\frac{V_{n-1}}{U_1}, \frac{U_{n+1}}{U_1}\right) + V_1 f\left(\frac{U_{n-1}}{V_1}, \frac{V_{n+1}}{V_1}\right) \\ &\quad + U_2 f\left(\frac{V_{n-2}}{U_2}, \frac{U_{n+2}}{U_2}\right) + V_2 f\left(\frac{U_{n-2}}{V_2}, \frac{V_{n+2}}{V_2}\right) + \cdots, \end{aligned} \quad (1.1.13)$$

where the sum on the right-hand side contains  $n$  terms. However, by (1.1.11), for  $r \geq 1$ ,

$$V_r f\left(\frac{U_{n-r}}{V_r}, \frac{V_{n+r}}{V_r}\right) = U_{n-r} f\left(\frac{U_{n-r}^2 V_{n+r}}{V_r^3}, \frac{V_r}{U_{n-r}}\right) = U_{n-r} f\left(\frac{U_{2n-r}}{U_{n-r}}, \frac{V_r}{U_{n-r}}\right).$$

Thus, we see that the sums on the right sides of (1.1.12) and (1.1.13) agree.

**Proof of Lemma 1.1.2.** Using the definitions of  $U_n$  and  $V_n$ , we see that

$$\begin{aligned} f(a, b) &= \sum_{k=-\infty}^{\infty} a^{k(k+1)/2} b^{k(k-1)/2} \\ &= \sum_{k=-\infty}^{\infty} \sum_{r=0}^{n-1} a^{(nk+r)(nk+r+1)/2} b^{(nk+r)(nk+r-1)/2} \\ &= \sum_{k=-\infty}^{\infty} \sum_{r=0}^{n-1} U_r^{1-k^2} U_{n+r}^{k(k+1)/2} V_{n-r}^{k(k-1)/2} \\ &= \sum_{k=-\infty}^{\infty} \sum_{r=0}^{n-1} U_r \left(\frac{U_{n+r}}{U_r}\right)^{k(k+1)/2} \left(\frac{V_{n-r}}{U_r}\right)^{k(k-1)/2} \\ &= \sum_{r=0}^{n-1} U_r f\left(\frac{U_{n+r}}{U_r}, \frac{V_{n-r}}{U_r}\right). \end{aligned}$$

□

## 1.2. Definitions and Generating Functions

**Definition 1.2.1.** *The ordinary or unrestricted partition function  $p(n)$  is the number of representations of the positive integer  $n$  as a sum of positive integers, where different orders of the summands are not considered to be distinct.*

Thus,  $p(4) = 5$ , because there are five ways to write 4 as a sum of positive integers, namely, 4, 3+1, 2+2, 2+1+1, 1+1+1+1. A generating function for  $p(n)$  can be given by, for  $|q| < 1$ ,

$$\sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q; q)_{\infty}} = \sum_{n_1=0}^{\infty} q^{1 \cdot n_1} \sum_{n_2=0}^{\infty} q^{2n_2} \cdots \sum_{n_k=0}^{\infty} q^{kn_k} \cdots, \quad (1.2.1)$$

where we make the convention that  $p(0) = 1$ , and where we have expanded each factor  $1/(1 - q^k)$ ,  $k \geq 1$ , in the denominator of  $1/(q; q)_{\infty}$  into a geometric series. Thus, the first series on the far right side generates the number of 1's in a particular partition of  $n$ , say; the second series generates the number of 2's in a partition of  $n$ , etc. It is clear that every partition of  $n$  can be achieved, and achieved uniquely, in the product of geometric series on the right side of (1.2.1). This formal argument can easily be made rigorous.

**Definition 1.2.2.** *Let  $p_m(n)$  denote the number of partitions of  $n$  into parts that are not larger than  $m$ .*

It is clear that

$$\frac{1}{(q; q)_m} = \sum_{n=0}^{\infty} p_m(n)q^n. \quad (1.2.2)$$

We now put these observations in a theorem, originally due to Euler.

**Theorem 1.2.3.** *For  $|q| < 1$ ,*

$$\sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q; q)_{\infty}}. \quad (1.2.3)$$

**Proof.** Observe that

$$\frac{1}{(q; q)_{\infty}} - \frac{1}{(q; q)_m} = \sum_{n=m+1}^{\infty} p_m^*(n)q^n \leq \sum_{n=m+1}^{\infty} p(n)q^n, \quad (1.2.4)$$


where  $p_m^*(n)$  denotes the number of partitions of  $n$  into parts, such that at least one of the parts is greater than or equal to  $m+1$ . Now let  $m \rightarrow \infty$  in (1.2.4). Because the series on the far right-hand side of (1.2.4) is a convergent series, by the dominated convergence theorem, the first series on the right-hand side of (1.2.4) tends to 0 as  $m \rightarrow \infty$ . We also remark that the infinite series on the left-hand side of (1.2.1) or (1.2.3) converges for  $|q| < 1$ , because  $1/(q; q)_{\infty}$  is analytic for  $|q| < 1$ . □



In the sequel, we derive generating functions for several types of partitions. Rigorous proofs can generally be developed along the same lines as our proof above for Theorem 1.2.3. To avoid repetitions of the same kind of argument, we shall forego such proofs in the future.

**Definition 1.2.4.** *Let  $S$  be any subset of the natural numbers. Then  $p(S, m, n)$  denotes the number of partitions of  $n$  into exactly  $m$  parts taken from  $S$ . If  $S = \mathbb{N}$ , then we delete  $\mathbb{N}$  from our notation. In particular,  $p(m, n)$  is the number of partitions of  $n$  into precisely  $m$  parts.*

We now introduce the concept of a Ferrers graph, which originated with N. M. Ferrers in the 1850's. The idea was exploited by J. J. Sylvester, who was always grateful to Ferrers for his graphical interpretation of a partition. Arrange the parts in a partition of  $n$  in decreasing order, and graphically represent a part  $j$  by  $j$  dots, with the first dot in each part flush at the left margin. We provide an example.



**Figure 1.** Ferrers graph of  $29 = 7 + 7 + 6 + 5 + 2 + 2$

**Definition 1.2.5.** *The Durfee square for a partition of  $n$  is the largest square of nodes in the upper left-hand corner of the Ferrers diagram of the partition; if the number of such nodes on a side is  $s$ , then we say that we have a Durfee square of side  $s$ .*

Thus, for the partition  $7 + 7 + 6 + 5 + 2 + 2$ , the Durfee square has size 4. The idea of a Durfee square is due to W. P. Durfee, who was a student of J. J. Sylvester.

**Theorem 1.2.6.** *Let  $p_m^*(n)$  denote the number of partitions of  $n$  into parts with largest part exactly equal to  $m$ , and let  $p(m, n)$  be defined as in Definition 1.2.4. Then*

$$p(m, n) = p_m^*(n), \quad (1.2.5)$$


*which implies that the partitions of  $n$  in which the largest is precisely  $m$  are equinumerous with the partitions of  $n$  into exactly  $m$  parts.*

**Proof.** Consider the Ferrers graph of  $n$ , such as the one in Figure 1, giving a partition of  $p(6, 29)$ . Instead of reading the partition from top to bottom, read the Ferrers graph from left to right. Thus, in Figure 1, this partition, namely,  $29 = 6 + 6 + 4 + 4 + 4 + 3 + 2$ , is counted by  $p_6^*(29)$ . Every such partition counted by  $p(m, n)$  is thus uniquely associated with a partition counted by  $p_m^*(n)$ . Clearly, this establishes a bijection for partitions counted by  $p(m, n)$  with those counted by  $p_m^*(n)$ . Hence, (1.2.5) immediately follows.  $\square$

As a corollary of Theorem 1.2.6, we see that the number of partitions of  $n$  in which the number of parts is less than or equal to  $m$  is equal to the number of partitions of  $n$  in which the largest part is no more than  $m$ . If we let  $p(r, m, n)$  equal the number of partitions of  $n$  into exactly  $m$  parts with the largest being  $r$ , then we have shown that  $p(r, m, n) = p(m, r, n)$ .

**Definition 1.2.7.** *The conjugate of a partition of  $n$  represented by a Ferrers graph is the partition that one obtains by reading the graph from left to right.*

**Definition 1.2.8.** *A partition of  $n$  is self-conjugate if its conjugate is identical with that same partition.*



**Figure 2.** Ferrers graph of the self-conjugate partition  $7 + 6 + 4 + 4 + 2 + 2 + 1$

If we enumerate the nodes in each right angle beginning with the outside and proceeding inward, we easily see that each right angle has an odd number of nodes, and, moreover these odd numbers are distinct. Thus, we obtain a partition into distinct odd numbers. In Figure 2, the partition  $13 + 9 + 3 + 1$  is generated. Conversely, if we have a partition into distinct odd parts, then we can generate a unique self-conjugate partition. We easily see that we have a bijection between self-conjugate partitions and the set of partitions into odd, distinct integers.

**Theorem 1.2.9.** *The set of self-conjugate partitions of a positive integer  $n$  is equinumerous with the set of partitions into distinct odd parts.*

We now use the Ferrers graph in Table 2 to help us determine the generating function for self-conjugate partitions, or a generating function for partitions into odd, distinct parts, where we keep track of the number of odd, distinct parts.

**Theorem 1.2.10** (Euler). *We have*

$$(-xq; q^2)_\infty = \sum_{j=0}^{\infty} \frac{x^j q^{j^2}}{(q^2; q^2)_j}. \quad (1.2.6)$$

**Proof.** Observe that the left side of (1.2.6) is the generating function of a positive integer  $n$  into, say  $s$ , odd, distinct parts. This leads us to examine the Ferrers graph of a self-conjugate partition, as in Figure 2, because of the one-to-one correspondence with such partitions and those partitions into distinct odd parts. As in Figure 2, we identify the largest Durfee square of side  $s$ . But now note that  $s$  is identical to the number of odd, distinct parts. We know that this partition is also a self-conjugate partition, and so the partition  $\pi_1$ , reading from top to bottom, below the Durfee square is identical to the

partition  $\pi_2$ , reading from left to right, to the right of the Durfee square. Each part in each partition is  $\leq s$ . Now consider the union of  $\pi_1$  and  $\pi_2$  formed by doubling either  $\pi_1$  or  $\pi_2$ . In our picture, both of these partitions are given by  $2 + 2 + 1$ , and so the union is  $4 + 4 + 2$ . We obtain a partition into even parts, with each part  $\leq 2s$ . The generating function for such partitions is  $1/(q^2; q^2)_s$ . Thus, the generating function for all such partitions into odd distinct parts with Durfee square of side  $s$  is  $q^{s^2}x^s/(q^2; q^2)_s$ . Summing over all  $s$  completes the proof.  $\square$

**Definition 1.2.11.** Let  $Q(n)$  denote the number of partitions of  $n$  into distinct parts. More generally,  $Q(S, m, n)$  denotes the number of partitions of  $n$  into  $m$  distinct parts of  $S$ . If  $S = \mathbb{N}$ , we write  $Q(\mathbb{N}, m, n) = Q(m, n)$ , so that  $Q(m, n)$  is the number of partitions of  $n$  into exactly  $m$  distinct parts.

**Theorem 1.2.12.** The generating function for  $Q(n)$  is given by

$$\sum_{n=0}^{\infty} Q(n)q^n = (-q; q)_{\infty}. \quad (1.2.7)$$

**Proof.** Observe that  $(-q; q)_{\infty}$  yields only the first two terms of each geometric series on the right-hand side of (1.2.1). Thus, in each partition of  $n$ , each integer not larger than  $n$  can appear in a particular partition of  $n$  at most one time. Theorem 1.2.12 is thus immediate.  $\square$

**Definition 1.2.13.** Let  $p_o(n)$  denote the number of partitions of  $n$  into odd parts, and let  $p_e(n)$  denote the number of partitions of  $n$  into even parts.

Observe that

$$\frac{1}{(q; q^2)_{\infty}} = \sum_{n=0}^{\infty} p_o(n)q^n \quad \text{and} \quad \frac{1}{(q^2; q^2)_{\infty}} = \sum_{n=0}^{\infty} p_e(n)q^n. \quad (1.2.8)$$

**Theorem 1.2.14.** For each  $n \geq 1$ ,

$$Q(n) = p_o(n). \quad (1.2.9)$$

**Proof.** By Theorem 1.2.12 and (1.2.8),

$$\begin{aligned} \sum_{n=0}^{\infty} Q(n)q^n &= (-q; q)_{\infty} = \frac{(-q; q)_{\infty}(q; q)_{\infty}}{(q; q)_{\infty}} \\ &= \frac{(q^2; q^2)_{\infty}}{(q; q)_{\infty}} = \frac{1}{(q; q^2)_{\infty}} = \sum_{n=0}^{\infty} p_o(n)q^n. \end{aligned}$$

We now see that (1.2.9) follows.  $\square$

Theorem 1.2.14 is due to Euler and illustrates one of the fascinating aspects of the theory of partitions, namely, that the number of partitions of  $n$  of one particular type often equals the number of partitions of  $n$  of an entirely different type. We mention another famous example in illustration. The first of the two Rogers–Ramanujan identities

that we will prove in Chapter 5 has the following combinatorial interpretation. The number of partitions of a positive integer  $n$  into parts differing by at least 2 is equal to the number of partitions of  $n$  into parts that are congruent to either 1 or 4 modulo 5. For example, the two respective sets of partitions for 8 are

$$8 = 7 + 1 = 6 + 2 = 5 + 3;$$

$$6 + 1 + 1 = 4 + 4 = 4 + 1 + 1 + 1 + 1 = 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1.$$

Theorems 1.2.3 and 1.2.12 can easily be generalized.

**Theorem 1.2.15.** For  $|q| < 1$  and  $|z| < 1/|q|$ ,

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} p(S, m, n) z^m q^n = \prod_{n \in S} \frac{1}{1 - zq^n}, \quad (1.2.10)$$

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} Q(S, m, n) z^m q^n = \prod_{n \in S} (1 + zq^n). \quad (1.2.11)$$

**Proof.** Let  $S = \{n_1, n_2, \dots\}$ . We prove (1.2.10); the proof of (1.2.11) is similar. Observe that

$$\prod_{n \in S} \frac{1}{1 - zq^n} = \prod_{j=1}^{\infty} \sum_{m_j=0}^{\infty} z^{m_j} q^{n_j m_j} \quad (1.2.12)$$

$$= \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \dots z^{m_1+m_2+\dots} q^{n_1 m_1 + n_2 m_2 + \dots}. \quad (1.2.13)$$

Let  $\pi$  be a partition of  $n$  into  $m_1$   $n_1$ 's,  $m_2$   $n_2$ 's, etc., i.e.,  $n = n_1 m_1 + n_2 m_2 + \dots$ . The number of parts in this partition is thus  $m := m_1 + m_2 + \dots$ . Thus, from (1.2.12) and our discussion, we see that (1.2.10) holds.  $\square$

We emphasize that the power of  $z$  in (1.2.10) and (1.2.11) keeps track of the number of parts in each partition of  $n$ .

**Definition 1.2.16.** We define  $p^{(s)}(S, m, n)$  to be the number of partitions of  $n \in S$  into exactly  $m$  parts, with each part appearing no more than  $s$  times in any partition of  $n$ .

**Theorem 1.2.17.** If  $zq^n \neq 1$  for each nonnegative integer  $n$ , then

$$\begin{aligned} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} p^{(s)}(S, m, n) z^m q^n &= \prod_{n \in S} (1 + q^n z + q^{2n} z^2 + \dots + q^{sn} z^s) \\ &= \prod_{n \in S} \frac{1 - q^{(s+1)n} z^{s+1}}{1 - q^n z}. \end{aligned}$$

**Proof.** Using the definition of  $p^{(s)}(S, m, n)$ , we find that the first equality of Theorem 1.2.17 holds, while the second equality arises from summing the geometric series.  $\square$

Observe that  $p^{(1)}(S, m, n) = Q(S, m, n)$ . If there are no restrictions on the number of parts  $m$ , we drop  $m$  from the notation for partition functions. For example, if  $S = \mathbb{O}$ , the set of all positive odd integers, then  $p(\mathbb{O}, n)$  denotes the number of partitions of  $n$  into odd summands, or odd parts. As before, if  $S = \mathbb{N}$ , we delete  $\mathbb{N}$  from our notation. In alternative notations, we reformulate Theorem 1.2.14.

**Theorem 1.2.18.** *If  $S = \mathbb{O}$ , then*

$$p(\mathbb{O}, n) = Q(n) = p^{(1)}(n). \quad (1.2.14)$$

**Second Proof of Theorem 1.2.18.** Suppose that we have a partition into odd parts. Merge the odd parts by successively doubling. Repeat this process until all the parts are distinct. Note that this process will terminate, because each merging operation decreases the number of parts by 1, and by construction, the final parts will be distinct. For example, consider


$$3 + 3 + 3 + 1 + 1 + 1 + 1 \rightarrow 6 + 3 + 2 + 2 \rightarrow 6 + 3 + 4.$$

Suppose that we have a partition into distinct parts. We begin a ‘halving’ operation by successively halving each even part. Clearly, this process must terminate. For example, consider

$$6 + 4 + 3 \rightarrow 3 + 3 + 2 + 2 + 3 \rightarrow 3 + 3 + 3 + 1 + 1 + 1 + 1.$$

It is easy to see that these operations are inverses of each other, i.e., we have a bijection, and so we have reproved Theorem 1.2.18.  $\square$

Recall that in Definition 1.2.5 we defined a Durfee square of side  $s$ . In Figure 3 below, we indicate the Durfee square of size 4.



**Figure 3.** A Ferrers graph with Durfee square of size 4

Using Durfee squares, we can derive another generating function for  $p(n)$ .

**Theorem 1.2.19.** *We have*

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n^2} = \sum_{n=0}^{\infty} p(n)q^n. \quad (1.2.15)$$

**Proof.** Consider the nodes lying below the Durfee square of size  $s$  for an arbitrary partition of  $n$ . We note that we have the graphical representation of a partition  $\pi_1$  of  $m_1$ , where  $m_1$  equals the number of nodes of the partition to below the Durfee square. Moreover, each part is no larger than  $s$ . Now examine the nodes to the right of the Durfee square. Reading from left to right, we have a partition  $\pi_2$  of a number

$m_2$  with each part  $\leq s$ . For example, if  $\pi$  is the partition represented in Figure 3, then  $s = 4$ ,  $\pi_1$  is the partition  $2+2$ , and  $\pi_2$  is the partition  $4+3+2$ . The number of choices for  $\pi_1$  is  $p(\{1, 2, \dots, s\}, m_1) =: p_s(m_1)$ , and the number of choices for  $\pi_2$  equals  $p(\{1, 2, \dots, s\}, m_2) =: p_s(m_2)$ . Note that  $n = s^2 + m_1 + m_2$ . The generating function for all partitions of  $n$ , with fixed  $s$  is thus

$$\sum_{n=0}^{\infty} q^n \sum_{n=s^2+m_1+m_2} p_s(m_1)p_s(m_2) = \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} q^{s^2+m_1+m_2} = \frac{q^{s^2}}{(q; q)_s^2}. \quad (1.2.16)$$

We now sum (1.2.16) over  $s$ ,  $0 \leq s < \infty$ , to generate all partitions. Thus,

$$\sum_{n=0}^{\infty} p(n)q^n = \sum_{s=0}^{\infty} \frac{q^{s^2}}{(q; q)_s^2},$$

which is the same as (1.2.15), and so the proof of Theorem 1.2.19 is complete.  $\square$

In view of Theorem 1.2.15, it is natural to ask if there is a generating function for  $p(\mathbb{N}, m, n)$  generalizing Theorem 1.2.19. We derive such a generating function in the next theorem.

**Theorem 1.2.20.** *We have*

$$\sum_{m, n=0}^{\infty} p(\mathbb{N}, m, n)z^m q^n = \frac{1}{(zq; q)_{\infty}} = \sum_{n=0}^{\infty} \frac{z^n q^{n^2}}{(zq; q)_n (q; q)_n}. \quad (1.2.17)$$

**Proof.** The argument is similar to that in the proof of Theorem 1.2.19. Let  $m_1$  denote the number of nodes in a partition below the Durfee square of size  $s$ , and let  $m_2$  denote the number of nodes in the partition  $\pi_2$ , with its nodes lying to the right of the Durfee square of size  $s$ . Thus,  $n = s^2 + m_1 + m_2$ . However, we now want to keep track of the number of parts  $r$  in the partition  $\pi_1$ . The number of such partitions is  $p(\{1, 2, \dots, s\}, r, m_1)$ , while the number of partitions  $\pi_2$  is  $p(\{1, 2, \dots, s\}, m_2)$ . Note that the number of parts of  $\pi$  is equal to  $s + r$ . Hence, for fixed  $s$ , the generating function for the number of partitions of  $n$  is equal to

$$\begin{aligned} & \sum_{n=0}^{\infty} q^n \sum_{r=0}^{\infty} \sum_{s^2+m_1+m_2=n} z^{r+s} p(\{1, 2, \dots, s\}, r, m_1) p(\{1, 2, \dots, s\}, m_2) \\ &= \sum_{r=0}^{\infty} \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} q^{s^2+m_1+m_2} z^{r+s} p_s(r, m_1) p_s(m_2) \\ &= \frac{q^{s^2} z^s}{(zq; q)_s (q; q)_s}. \end{aligned} \quad (1.2.18)$$

We now sum over  $s$ ,  $0 \leq s < \infty$ , and use (1.2.10) to deduce that

$$\sum_{s=0}^{\infty} \frac{q^{s^2} z^s}{(zq; q)_s (q; q)_s} = \frac{1}{(zq; q)_{\infty}}.$$

This completes the proof.  $\square$

**Definition 1.2.21.** Let  $Q_m(n)$  stand for the number of partitions of  $n$  into exactly  $m$  distinct parts.

**Definition 1.2.22.** The Ferrers triangle for a partition of  $n$  is the largest right, isosceles triangle in the upper left-hand corner of the Ferrers graph of  $n$ .

**Theorem 1.2.23.** The generating function for  $Q_m(n)$  is given by

$$\sum_{n=0}^{\infty} Q_m(n)q^n = \frac{q^{m(m+1)/2}}{(q; q)_m}. \quad (1.2.19)$$

**Proof.** Take a partition of  $n$  into exactly  $m$  distinct parts. We see that the largest Durfee triangle has  $\frac{1}{2}m(m+1)$  nodes. We will have a partition of  $n - \frac{1}{2}m(m+1)$  into no more than  $m$  parts to the right of this Durfee triangle. There are no further restrictions, and so the generating function for all those partitions lying to the right of the Durfee triangle is  $1/(q; q)_m$ . Thus, for a fixed  $m$ , the generating function for such partitions of  $n$  with Durfee triangle of side  $m$  is equal to

$$\frac{q^{m(m+1)/2}}{(q; q)_m}.$$

□

Our next theorem is an analogue of Theorem 1.2.20 and a refinement of Theorem 1.2.23. Recall that  $Q_m(n)$  denotes the number of partitions of  $n$  into precisely  $m$  distinct parts.

**Definition 1.2.24.** The length of a partition  $\ell(\pi)$  is the number of parts of  $\pi$ .

**Theorem 1.2.25.** The generating function for  $Q_m(n)$  is given by

$$\begin{aligned} \sum_{m,n=0}^{\infty} Q_m(n)z^m q^n &= \prod_{n=1}^{\infty} (1 + zq^n) \\ &= \sum_{s=0}^{\infty} z^s q^{s(3s+1)/2} (zq^{2s+1} + 1) \frac{(-zq; q)_s}{(q; q)_s}. \end{aligned} \quad (1.2.20)$$

**Proof.** Suppose that  $\pi$  is a partition of  $n$  with distinct parts and Durfee square of side  $s$ . We consider two cases: 1) The lower edge of the Durfee square constitutes a complete part of  $n$ . 2) The lower edge of the Durfee square does not constitute a complete part of  $n$ . We regard  $s$  as fixed.

Case 1. Let  $\pi_1$  denote the partition below the Durfee square, where we read the parts from top to bottom. Note that  $\pi_1$  is a partition of  $n$ , say, with  $w$ , say, distinct parts, each of which is  $\leq s - 1$ . Let  $\pi_2$  denote the partition to the right of the Durfee square, where we read from top to bottom. Let us say that  $\pi_2$  is a partition of  $m$ , say, with exactly  $s - 1$  distinct parts. We see that  $\ell(\pi) = s + \ell(\pi_1)$ . The generating function

for these partitions is, for  $s$  fixed,

$$\begin{aligned}
& \sum_{N=0}^{\infty} \sum_{M=0}^{\infty} z^M q^N \sum_{s^2+m+n=N} Q_{s-1}(m) \sum_{s+w=M} Q(\{1, 2, \dots, s-1\}, w, n) \\
&= z^s q^{s^2} \sum_{n=0}^{\infty} \sum_{w=0}^{\infty} Q(\{1, 2, \dots, s-1\}, w, n) z^w q^n \sum_{m=0}^{\infty} Q_{s-1}(m) q^m \\
&= z^s q^{s^2} (-zq; q)_{s-1} \frac{q^{s(s-1)/2}}{(q; q)_{s-1}}, \tag{1.2.21}
\end{aligned}$$

by Theorems 1.2.15 and 1.2.23.

Case 2. As in Case 1,  $\pi_1$  is the partition represented below the Durfee square of side  $s$ , where we read from top to bottom. The partition  $\pi_2$  is located to the right of the Durfee square, and we again read the partition from top to bottom. The parts are distinct, and there are exactly  $s$  of them. The generating function for these partitions is

$$\begin{aligned}
& \sum_{N=0}^{\infty} \sum_{M=0}^{\infty} z^M q^N \sum_{s^2+m+n=N} Q_s(m) \sum_{s+w=M} Q(\{1, 2, \dots, s\}, w, n) \\
&= z^s q^{s^2} (-zq; q)_s \frac{q^{s(s+1)/2}}{(q; q)_s}, \tag{1.2.22}
\end{aligned}$$

by Theorems 1.2.15 and 1.2.23 once again. We now sum over  $s$ ,  $1 \leq s < \infty$ , in (1.2.21), and sum over  $s$ ,  $0 \leq s < \infty$ , in (1.2.22). Hence,

$$\begin{aligned}
(-zq; q)_{\infty} &= \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} Q_m(n) z^m q^n \\
&= \sum_{s=1}^{\infty} z^s q^{s^2} (-zq; q)_{s-1} \frac{q^{s(s-1)/2}}{(q; q)_{s-1}} \\
&\quad + \sum_{s=0}^{\infty} z^s q^{s^2} (-zq; q)_s \frac{q^{s(s+1)/2}}{(q; q)_s} \\
&= \sum_{s=0}^{\infty} z^{s+1} q^{s^2+2s+1} (-zq; q)_s \frac{q^{s(s+1)/2}}{(q; q)_s} \\
&\quad + \sum_{s=0}^{\infty} z^s q^{s^2+s(s+1)/2} \frac{(-zq; q)_s}{(q; q)_s} \\
&= \sum_{s=0}^{\infty} z^s q^{s(3s+1)/2} (zq^{2s+1} + 1) \frac{(-zq; q)_s}{(q; q)_s},
\end{aligned}$$

which is (1.2.20). □

**Corollary 1.2.26** (Euler's Pentagonal Number Theorem). *For  $|q| < 1$ ,*

$$(q; q)_{\infty} = \sum_{s=-\infty}^{\infty} (-1)^s q^{s(3s-1)/2}. \tag{1.2.23}$$



**Proof.** Set  $z = -1$  in (1.2.20) and replace  $s$  by  $-s - 1$  in the second series in the second step below. Therefore,

$$\begin{aligned}
(q; q)_\infty &= \sum_{s=0}^{\infty} (-1)^s q^{s(3s+1)/2} (1 - q^{2s+1}) \\
&= \sum_{s=0}^{\infty} (-1)^s q^{s(3s+1)/2} - \sum_{s=0}^{\infty} (-1)^s q^{s(3s+1)/2 + 2s+1} \\
&= \sum_{s=0}^{\infty} (-1)^s q^{s(3s+1)/2} + \sum_{s=-1}^{-\infty} (-1)^s q^{s(3s+1)/2} \\
&= \sum_{s=-\infty}^{\infty} (-1)^s q^{s(3s+1)/2} \\
&= \sum_{s=-\infty}^{\infty} (-1)^s q^{s(3s-1)/2}, \tag{1.2.24}
\end{aligned}$$

where we replaced  $s$  by  $-s$ . □

**Definition 1.2.27.** *The numbers  $s(3s - 1)/2$ ,  $1 \leq s < \infty$ , are called the pentagonal numbers, while the numbers  $s(3s - 1)/2$ ,  $-\infty < s < \infty$ , are called the generalized pentagonal numbers.*

Observe that (1.2.23) tells us that the cardinality of the set of partitions of  $n$  with an even number of distinct parts is usually equal to the cardinality of the set of partitions of  $n$  with an odd number of distinct parts. We make this observation more precise.

**Corollary 1.2.28.** *Let  $D_e(n)$  denote the number of partitions of  $n$  into an even number of distinct parts, and let  $D_o(n)$  denote the number of partitions of  $n$  into an odd number of distinct parts. Then*

$$D_e(n) - D_o(n) = \begin{cases} 0, & \text{if } n \text{ is not a generalized pentagonal number,} \\ (-1)^s & \text{if } n \text{ is a generalized pentagonal number } \frac{1}{2}s(3s - 1). \end{cases} \tag{1.2.25}$$

Corollary 1.2.26 leads to a method for numerically calculating values of  $p(n)$ , as we shall see in the next theorem. This recurrence relation was used by P. A. MacMahon to calculate  $p(n)$ ,  $1 \leq n \leq 200$ , and by H. Gupta to determine  $p(n)$  for  $201 \leq n \leq 300$ . If one wants to calculate a certain value of  $p(n)$  on a computer, then the computer will likely employ Theorem 1.2.29 to do so.

**Theorem 1.2.29.** *For each positive integer  $N$ ,*

$$p(N) = \sum_{\substack{k(3k \pm 1)/2 + n = N \\ 0 \leq n < N}} (-1)^{k-1} p(n). \tag{1.2.26}$$

**Proof.** From the generating function (1.2.1) for  $p(n)$  and the pentagonal number theorem, Corollary 1.2.23, we see that

$$1 = \frac{(q; q)_\infty}{(q; q)_\infty} = \sum_{k=-\infty}^{\infty} (-1)^k q^{k(3k-1)/2} \sum_{n=0}^{\infty} p(n) q^n. \quad (1.2.27)$$

Equating coefficients of  $q^N$ ,  $N > 0$ , in (1.2.27), we find that

$$0 = \sum_{k(3k\pm 1)/2+n=N} (-1)^k p(n). \quad (1.2.28)$$

Solving (1.2.28) for  $p(N)$ , we complete the proof of (1.2.26).  $\square$

Instead of using  $(q; q)_\infty$ , we can take its logarithmic derivative to derive a recurrence formula for  $\sigma(n) := \sum_{d|n} d$ , as we now demonstrate.

**Theorem 1.2.30.** *If  $n \in \mathbb{N}$ , then*

$$\sum_{\substack{k(3k\pm 1)/2+j=n \\ 1 \leq j \leq n}} (-1)^k \sigma \left( n - \frac{k(3k-1)}{2} \right) = \begin{cases} (-1)^k n, & \text{if } n = \frac{1}{2}k(3k-1), \\ 0, & \text{otherwise.} \end{cases}$$

**Proof.** Let  $F(q) = (q; q)_\infty$ . We first observe that

$$q \frac{F'(q)}{F(q)} = - \sum_{j=1}^{\infty} \frac{j q^j}{1 - q^j} = - \sum_{j=1}^{\infty} j \sum_{k=1}^{\infty} q^{jk} = - \sum_{n=1}^{\infty} \sigma(n) q^n. \quad (1.2.29)$$

By (1.2.23) and (1.2.29),

$$q F'(q) = F(q) \frac{q F'(q)}{F(q)} = - \sum_{k=-\infty}^{\infty} (-1)^k q^{k(3k-1)/2} \sum_{j=1}^{\infty} \sigma(j) q^j. \quad (1.2.30)$$

On the other hand, by (1.2.23),

$$q F'(q) = \sum_{n=-\infty}^{\infty} (-1)^n \frac{n(3n-1)}{2} q^{n(3n-1)/2}. \quad (1.2.31)$$

If we equate coefficients of  $q^n$ ,  $n \geq 1$ , in (1.2.30) and (1.2.31), we conclude that

$$\sum_{k=-\infty}^{\infty} (-1)^k \sigma \left( n - \frac{k(3k-1)}{2} \right) = \begin{cases} (-1)^k n, & \text{if } n = \frac{1}{2}k(3k-1), \\ 0, & \text{otherwise,} \end{cases}$$

and this completes the proof.  $\square$

The next result falls under the same purview as the two preceding theorems.

**Theorem 1.2.31.** *For  $n \geq 1$ ,*

$$\sigma(n) = \sum_{k=-\infty}^{\infty} (-1)^{k+1} \frac{k(3k+1)}{2} p \left( n - \frac{k(3k+1)}{2} \right). \quad (1.2.32)$$

**Proof.** Employing (1.2.29) and (1.2.30), we find that

$$\begin{aligned} -\sum_{n=1}^{\infty} \sigma(n)q^n &= \frac{qF'(q)}{F(q)} = \frac{1}{F(q)} \cdot qF'(q) \\ &= \frac{1}{(q; q)_{\infty}} \sum_{k=-\infty}^{\infty} (-1)^k \frac{k(3k+1)}{2} q^{k(3k+1)/2}. \end{aligned}$$

Equating coefficients of  $q^n$  on both sides above, we complete the proof.  $\square$

We now offer a bijective proof of Corollary 1.2.26, or Theorem 1.2.28, which is due to F. Franklin in 1881. *Franklin's Bijection* has been employed numerous times in bijective proofs in the theory of partitions.

**Second Proof of Corollaries 1.2.26 and 1.2.28.** Consider the Ferrers graph of a partition of  $n$  into distinct parts. Identify the Durfee square of side  $s$ . To the right of this square, there will be a Durfee triangle with side less than or equal to  $s$ . Reading from top to bottom, we have a partition  $\pi_1$  with distinct parts below the square, with the largest part being not greater than  $s$ . Reading from left to right, we see that we have a partition  $\pi_2$  in which each part is no larger than  $s$ . We consider two cases: 1) Each part of  $\pi_2$  is not larger than  $s-1$ . 2) One part of  $\pi_2$  is equal to  $s$ .

Case 1. If the smallest part in the upper region is smaller than any part in the lower region, transfer that smallest part to the lower region. Otherwise, the smallest part in the lower region is transferred to the upper region. In the case of a tie, we transfer the smallest part in the lower region to the upper region. Note that such a transfer changes the parity of the number of parts. Of course, this procedure breaks down if there are not any parts in both regions. As an example, consider the partition  $\pi = 8 + 4 + 3 + 2 + 1$ . We observe that the Durfee square has side 3, and the Durfee triangle has side 2. In the lower region,  $\pi_1 = 2 + 1$ ; in the upper region,  $\pi_2 = 1 + 1 + 1$ . The smallest part of these two partitions is equal to 1 in each instance. Thus, we transfer the part 1 in the lower region to a part 1 in the upper region. The parities of the partitions in each case changes with the transfer. This establishes a one-to-one correspondence between partitions in those cases when at least one of the partitions  $\pi_1, \pi_2$  is non-empty. Hence, for such partitions,  $D_e(n) - D_o(n) = 0$ . We now need to consider those cases when  $\pi_1$  and  $\pi_2$  are both empty. In Case 1, the lower side of the Durfee square is equal to  $s$ , and this implies that the Durfee triangle has side equal to  $s-1$ . Thus, the number of nodes in a partition giving such a Ferrers graph is  $s^2 + s(s-1)/2 = s(3s-1)/2$ . If  $s$  is odd, then there is one more odd part than even part. In such a case,  $D_e(n) - D_o(n) = -1 = (-1)^s$ . If  $s$  is even, then the number of parts with even parity equals the number with odd parity, and so there is an 'extra' partition with an even number of distinct parts.

Case 2. The argument is identical to that in Case 1 when at least one of  $\pi_1, \pi_2$  is non-empty. Thus, we obtain equal numbers of partitions with an odd number of distinct parts, and with an even number of distinct parts. There are cases, however, when both  $\pi_1$  and  $\pi_2$  are empty. In such cases, the side of the Durfee triangle is equal to  $s$ . Thus, if  $s$  is even, we conclude that  $D_e(n) - D_o(n) = 1 = (-1)^s$ . If  $s$  is odd, then we obtain one

additional partition with an odd number of parts, and  $D_e(n) - D_o(n) = -1 = (-1)^s$ . To illustrate this case, consider the partition  $\pi = 9 + 8 + 6 + 3 + 1$ . Here,  $s = 3$ , which is also equal to the side of the Durfee triangle. We thus obtain one additional partition with an odd number of distinct parts.

Thus, Corollaries 1.2.26 and 1.2.28 have been established.  $\square$

### 1.3. A Proof Due to Sun Kim [73]

We conclude our chapter on elementary methods with a generalization of Theorem 1.2.31. We first give a rather straightforward proof of this generalization giving a formula for a certain divisor function as a sum of certain partition functions. We then give an elegant combinatorial proof of this theorem due to Sun Kim [73].

**Theorem 1.3.1.** *Let  $p_{n,m}(N)$  equal the number of partitions of  $N$  into parts  $\equiv n, m$ , or  $0 \pmod{L}$ , where  $m + n = L$ . Let*

$$\sigma_{m,n}(k) = \sum_{\substack{d|k \\ d \equiv m, n, 0 \pmod{L}}} d.$$

Then

$$-\sigma_{n,m}(N) = \sum_{k=-\infty}^{\infty} (-1)^k \frac{k^2(m+n) + k(m-n)}{2} p_{m,n} \left( N - \frac{k^2(m+n) + k(m-n)}{2} \right). \quad (1.3.1)$$

**First Proof of Theorem 1.3.1.** If we logarithmically differentiate the Jacobi triple product representation for  $f(-q^m, q^n)$ , given in (1.1.7), where  $m$  and  $n$  are arbitrary nonnegative integers, we find that

$$\begin{aligned} q \frac{\frac{d}{dq} f(-q^m, -q^n)}{f(-q^m, -q^n)} &= - \sum_{k=1}^{\infty} \frac{(km+n)q^{km+n}}{1-q^{km+n}} \\ &\quad - \sum_{k=1}^{\infty} \frac{(m+kn)q^{m+kn}}{1-q^{m+kn}} - \sum_{k=1}^{\infty} \frac{k(m+n)q^{k(m+n)}}{1-q^{k(m+n)}} \\ &= - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} (km+n)q^{(km+n)(j+1)} \\ &\quad - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} (m+kn)q^{(m+kn)(j+1)} - \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} k(m+n)q^{k(m+n)(j+1)} \\ &= - \sum_{k=1}^{\infty} \sum_{\substack{d|k \\ d \equiv n, m, 0 \pmod{L}}} dq^k \\ &= - \sum_{k=1}^{\infty} \sigma_{n,m}(k)q^k. \end{aligned}$$

On the other hand,

$$\begin{aligned} q \frac{\frac{d}{dq} f(-q^m, -q^n)}{f(-q^m, -q^n)} &= q \frac{q}{dq} \sum_{k=-\infty}^{\infty} (-1)^k q^{k^2(m+n)/2+k(m-n)/2} \sum_{j=0}^{\infty} p_{m,n}(j) q^j \\ &= \sum_{k=-\infty}^{\infty} (-1)^k \frac{k^2(m+n) + k(m-n)}{2} q^{k^2(m+n)/2+k(m-n)/2} \sum_{j=0}^{\infty} p_{m,n}(j) q^j \\ &= \sum_{N=0}^{\infty} \sum_{k=-\infty}^{\infty} (-1)^k \frac{k^2(m+n) + k(m-n)}{2} p_{m,n} \left( N - \frac{k^2(m+n) + k(m-n)}{2} \right) q^N. \end{aligned}$$

If we equate the coefficients of  $q^N$  on the far right-hand sides of the two strings of equalities above, we immediately deduce (1.3.1)  $\square$

We now give a beautiful combinatorial proof of (1.3.1) that is due to Sun Kim [73].

**Second Proof of Theorem 1.3.1.** We begin with some definitions.

**Definition 1.3.2.** Let  $D_{m,n}(N)$  denote the set of partitions  $d_{m,n}(N)$  of  $N$  into distinct parts that are  $\equiv m, n, 0 \pmod{L}$ . Let  $P_{m,n}(N)$  denote the set of unrestricted partitions  $p_{m,n}(N)$  of  $N$  into parts that are  $\equiv m, n, 0 \pmod{L}$ . Let

$$A_{m,n}(N) = \{(\pi, \lambda) : |\pi| + |\lambda| = N, \pi \in D_{m,n}, \lambda \in P_{m,n}\}.$$

Lastly, let  $\ell(\pi)$  denote the number of parts of the partition  $\pi$ ; we similarly define  $\ell(\lambda)$ .

Let

$$B_{m,n}(N) := \sum_{(\pi, \lambda) \in A_{m,n}(N)} (-1)^{\ell(\pi)} |\pi|.$$

Our task is to show that  $B_{m,n}(N)$  is equal to both the left and right sides of (1.3.1).

We first attempt to reach the right-hand side of (1.3.1). For brevity, set  $b = |\lambda|$ , so that  $|\pi| = N - b$ . Thus, we can write

$$B_{m,n}(N) = \sum_{b=0}^N (N-b) p_{m,n}(b) \sum_{\pi \in D_{m,n}(N-b)} (-1)^{\ell(\pi)}. \quad (1.3.2)$$

First, by the Jacobi triple product identity (1.1.7) and the definition of  $p_{m,n}(N)$ ,

$$\frac{1}{f(-q^m, -q^n)} = \frac{1}{(q^m; q^L)_\infty (q^n; q^L)_\infty (q^L; q^L)_\infty} = \sum_{N=0}^{\infty} p_{m,n}(N) q^N.$$

Second, by the definition of  $f(-q^m, -q^n)$ , the Jacobi triple product identity (1.1.7), and the definition of  $d_{m,n}(N)$ ,

$$\begin{aligned} \sum_{k=-\infty}^{\infty} (-1)^k q^{(m+n)k^2/2+(m-n)k/2} &= \sum_{k=-\infty}^{\infty} (-1)^k q^{k(k+1)m/2+k(k-1)n/2} \\ &= f(-q^m, -q^n) = (q^m; q^L)_\infty (q^n; q^L)_\infty (q^L; q^L)_\infty = \sum_{N=0}^{\infty} (-1)^{\ell(\pi)} d_{m,n}(N) q^N. \end{aligned}$$

Thus, if

$$\pi \in D_{m,n} \left( \frac{(m+n)k^2 + (m-n)k}{2} \right),$$

then

$$b = N - \frac{(m+n)k^2 + (m-n)k}{2}.$$

Now remember that  $\pi$  is a partition into parts congruent to  $m$ ,  $n$ , or  $0$  modulo  $L = m+n$ , and so the number being partitioned is composed of a linear combination of  $m$ 's,  $n$ 's, and  $L$ 's. Now one can think of  $(m+n)k^2/2 + (m-n)k/2$  as  $k(k+1)/2$   $m$ 's and  $k(k-1)/2$   $n$ 's for a total of  $k^2$   $m$  and  $n$ 's. But, generally, the partition also contains some  $L$ 's, each of which contains one  $m$  and one  $n$ . Hence, we see that

$$(-1)^k = (-1)^{k^2} = (-1)^{\ell(\pi)}.$$

Hence, substituting into (1.3.2) the values we have just determined for  $b$ ,  $N - b$ , and  $(-1)^{\ell(\pi)}$ , we find that

$$B_{m,n}(N) = \sum_{k=-\infty}^{\infty} (-1)^k \frac{(m+n)k^2 + (m-n)k}{2} p_{m,n} \left( N - \frac{(m+n)k^2 + (m-n)k}{2} \right). \quad (1.3.3)$$

Demonstrating that  $B_{m,n}(N)$  equals the left side of (1.3.1) is more difficult. First, we show that

$$\#\{(\pi, \lambda) \in A_{m,n}(\lambda) : \ell(\pi) \text{ is even}\} = \#\{(\pi, \lambda) \in A_{m,n}(\lambda) : \ell(\pi) \text{ is odd}\} \quad (1.3.4)$$

For any partition  $\pi^*$ , define

$$s(\pi^*) = \text{smallest part of } \pi^*; \quad s(\pi^*) = \infty, \text{ if } \pi^* = \emptyset.$$

If  $s(\pi) \leq s(\lambda)$ , then move  $s(\pi)$  to  $\lambda$ . If  $\pi'$  and  $\lambda'$  are the new partitions thus formed, we observe that  $s(\lambda') < s(\pi')$  and that  $(-1)^{\ell(\pi)} = -(-1)^{\ell(\pi')}$ . If  $s(\lambda) < s(\pi)$ , then move  $s(\lambda)$  to  $\pi$ . If  $\pi'$  and  $\lambda'$  are the newly formed partitions, we see that  $s(\pi') \leq s(\lambda')$  and that  $(-1)^{\ell(\pi)} = -(-1)^{\ell(\pi')}$ . We thus have established an involution between the two sets of partitions with  $s(\pi) \leq s(\lambda)$  and  $s(\lambda) < s(\pi)$ , with  $\ell(\pi)$  changing by 1 with each movement of a smallest part from one set to the other. This then establishes the equality in (1.3.4).

We now subdivide our partitions  $\pi$  according to their smallest parts  $a$ . In the case that  $s(\pi) > s(\lambda)$ , we can think of this situation as arising from moving the smallest part  $a$  from a partition  $\pi'$  into one of size  $|\pi| - a$  with sign  $(-1)^{\ell(\pi)-1}$ . Thus,

$$\begin{aligned} B_{m,n}(N) &:= \sum_{(\pi, \lambda) \in A_{m,n}(N)} (-1)^{\ell(\pi)} |\pi| \\ &= \sum_{s(\pi) \leq s(\lambda)} (-1)^{\ell(\pi)} |\pi| + \sum_{s(\pi) > s(\lambda)} (-1)^{\ell(\pi)} |\pi| \\ &= \sum_{a=1}^N \sum_{\substack{s(\pi)=a \\ s(\pi) \leq s(\lambda)}} (-1)^{\ell(\pi)} (|\pi| - (|\pi| - a)) \end{aligned}$$

$$= \sum_{a=1}^N a \sum_{\substack{s(\pi)=a \\ s(\pi) \leq s(\lambda)}} (-1)^{\ell(\pi)}. \quad (1.3.5)$$

In view of (1.3.1) and (1.3.5), it now suffices to show that

$$\sum_{\substack{s(\pi)=a \\ s(\pi) \leq s(\lambda)}} (-1)^{\ell(\pi)} = \begin{cases} -1, & \text{if } a|N, \\ 0, & \text{otherwise.} \end{cases} \quad (1.3.6)$$

Let  $L(\pi^*)$  denote the largest part of a partition  $\pi^*$ , with the convention that  $L(\pi^*) = 0$ , if  $\pi^* = \emptyset$ . Consider  $(\pi, \lambda) \in A_{m,n}(N)$ , with  $s(\pi) = a \leq s(\lambda)$ . Let  $\pi = a + \mu$ . If  $L(\mu) \geq L(\lambda)$  and  $\mu \neq \emptyset$ , then move  $L(\mu)$  to  $\lambda$ . If  $L(\mu) < L(\lambda)$ , except when  $L(\mu) = 0 < L(\lambda) = a$ , then move  $L(\lambda)$  to  $\mu$ . We thus obtain a new partition pair  $(\pi', \lambda')$  with

$$s(\pi') = a \leq s(\lambda'), \quad (-1)^{\ell(\pi)} = -(-1)^{\ell(\pi')}.$$

However, this map fails in two cases. First, suppose that  $\pi = a$ , i.e.,  $\mu = \emptyset$ , and  $\lambda = \emptyset$ . Then there is nothing to move. Second, suppose that  $\pi = a$  and  $\lambda = a + a + \cdots + a$ . Again, in this case, there is nothing to move, for if we did move  $a$  from  $\lambda$ , we would not obtain a new partition  $\pi'$  with distinct parts, i.e., we would have two parts equaling  $a$ . Thus, in these exceptional cases,  $a|N$ . Thus,

$$\sum_{\substack{s(\pi)=a \\ s(\pi) \leq s(\lambda)}} (-1)^{\ell(\pi)} = \begin{cases} -1, & \text{if } a|N, \\ 0, & \text{otherwise,} \end{cases}$$

i.e., (1.3.6) has been demonstrated. Hence,

$$B_{m,n}(N) := \sum_{(\pi, \lambda) \in A_{m,n}(N)} (-1)^{\ell(\pi)} |\pi| = - \sum_{\substack{a|N \\ a \equiv m, n, L \pmod{L}}} = -\sigma_{m,n}(N). \quad (1.3.7)$$

Hence, (1.3.3) and (1.3.7) taken together yield (1.3.1) to complete the proof.  $\square$

## 1.4. Exercises

1. Prove that the number of partitions of  $n$  into parts  $\equiv 1, 2 \pmod{3}$  is equal to  $p^{(2)}(n)$ .
2. Prove that

$$\begin{aligned} p(n|\text{even (odd) number of odd parts}) & \\ &= p(n|\text{distinct parts, with an even (odd) number of odd parts}) \end{aligned} \quad (1.4.1)$$

**Example 1.4.1.** Let  $n = 4$ . Those partitions with an even number of odd parts are:  $3 + 1, 1 + 1 + 1 + 1$ . The partitions of 4 with an even number of distinct odd parts are:  $4, 3 + 1$ .

3. Let  $k \geq 2$  and  $n \geq 1$ . Prove that

$$\begin{aligned} p(n|\text{no part is divisible by } k) \\ = p(n|\text{there are less than } k \text{ copies of each part}). \end{aligned}$$

Observe that if  $k = 2$  in Exercise 3, then its conclusion gives  $p(\mathbb{O}, n) = Q(n)$ , i.e., we obtain Theorem 1.2.14.

4. Prove Lemma 1.1.1.



---

## Chapter 2

# MacMahon's Partition Analysis; Gaussian Binomial Coefficients

### 2.1. MacMahon's Partition Analysis

In this chapter, we examine an elementary, but powerful, method due to P. A. MacMahon in the study of partitions. (The initials P. A. stand for Percy Alexander.) In his time, he was known as Major MacMahon, because he served as a Major in the British army for 20 years before he became a mathematician.

Recall that in Chapter 1,  $p(m, n)$  designates the number of partitions with exactly  $m$  parts, while  $p_m^*(n)$  denoted the number of partitions of  $n$  with largest part  $m$ . However, in Theorem 1.2.6, we showed that  $p(m, n) = p_m^*(n)$ . As a simple corollary, we observed that the number of partitions of an integer  $n$  into parts  $\leq m$  is equal to the number of partitions of  $n$  into  $\leq m$  parts. Since these two numbers are the same, in this chapter we shall think of  $p_m(n)$  in the latter interpretation.

**Definition 2.1.1.** *The number of partitions of  $n$  into no more than  $m$  parts is to be denoted by  $p_m(n)$  in the sequel.*

We put the  $m$  parts of a partition  $\pi$  in decreasing order, say  $n_1 \geq n_2 \geq \cdots \geq n_m \geq 0$ . Note that we allow some of the parts to be equal to 0. Thus, we can write the generating function for  $p_m(n)$  in the form

$$\sum_{n=0}^{\infty} p_m(n)q^n = \sum_{n_1 \geq n_2 \geq \cdots \geq n_m \geq 0} q^{n_1+n_2+\cdots+n_m}.$$

Consider the sum

$$\sum_{n_1, n_2, \dots, n_m \geq 0} q^{n_1+n_2+\cdots+n_m} \lambda_1^{n_1-n_2} \lambda_2^{n_2-n_3} \cdots \lambda_{m-1}^{n_{m-1}-n_m}.$$

To consider only those terms with nonnegative powers of  $\lambda_j$ ,  $1 \leq j \leq m-1$ , we introduce the restrictions  $n_1 \geq n_2, n_2 \geq n_3, \dots, n_{m-1} \geq n_m$ . We now define an operator that performs exactly this task.

**Definition 2.1.2.** Define an operator  $\Omega_{\geq}$  on multiple Laurent series so that it annihilates terms with negative exponents and sets any remaining  $\lambda_j$ 's equal to 1.

Note that

$$\begin{aligned} \sum_{n=0}^{\infty} p_m(n) q^n &= \Omega_{\geq} \sum_{n_1, n_2, \dots, n_m \geq 0} q^{n_1+n_2+\dots+n_m} \lambda_1^{n_1-n_2} \lambda_2^{n_2-n_3} \dots \lambda_{m-1}^{n_{m-1}-n_m} \\ &= \Omega_{\geq} \sum_{n_1=0}^{\infty} (q\lambda_1)^{n_1} \sum_{n_2=0}^{\infty} (q\lambda_2/\lambda_1)^{n_2} \dots \sum_{n_{m-1}=0}^{\infty} (q\lambda_{m-1}/\lambda_{m-2})^{n_{m-1}} \sum_{n_m=0}^{\infty} (q/\lambda_{m-1})^{n_m} \\ &= \Omega_{\geq} \frac{1}{(1-q\lambda_1)(1-q\lambda_2/\lambda_1) \dots (1-q\lambda_{m-1}/\lambda_{m-2})(1-q/\lambda_{m-1})}. \end{aligned} \quad (2.1.1)$$

We now explicitly calculate the effect of the operator  $\Omega_{\geq}$  on special cases of the identity above.

**Lemma 2.1.3.**

$$\Omega_{\geq} \frac{1}{(1-\lambda x)(1-y/\lambda)} = \frac{1}{(1-x)(1-xy)}. \quad (2.1.2)$$

**Proof.** Setting  $k = n - m$  below, we find that

$$\begin{aligned} \Omega_{\geq} \frac{1}{(1-\lambda x)(1-y/\lambda)} &= \Omega_{\geq} \sum_{n=0}^{\infty} (\lambda x)^n \sum_{m=0}^{\infty} (y/\lambda)^m = \sum_{n \geq m \geq 0} x^n y^m \\ &= \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} x^{m+k} y^m = \sum_{k=0}^{\infty} x^k \sum_{m=0}^{\infty} (xy)^m \\ &= \frac{1}{(1-x)(1-xy)}. \end{aligned}$$

□

The following lemma is (1.2.2), for which we wrote, "It is clear that ...". It is also a special case of Theorem 1.2.15, equation (1.2.10). To illustrate MacMahon's partition analysis, we will prove it again here.

**Lemma 2.1.4.** Recall that  $p_m(n)$  is defined in Definition 2.1.1. Then

$$\sum_{n=0}^{\infty} p_m(n) q^n = \frac{1}{(q; q)_m}.$$

**Proof.** We apply Lemma 2.1.3 several times. In the first instance, we replace  $x$  by  $q$ ,  $\lambda$  by  $\lambda_1$ , and  $y$  by  $\lambda_2 q$ ; in the second application, we replace  $x$  by  $q^2$ ,  $\lambda$  by  $\lambda_2$ , and  $y$  by

$q\lambda_3$ , etc. Hence, by (2.1.1) and Lemma 2.1.3,

$$\begin{aligned} \sum_{n=0}^{\infty} p_m(n)q^n &= \Omega \frac{1}{(1-q\lambda_1)(1-q\lambda_2/\lambda_1)\cdots(1-q\lambda_{m-1}/\lambda_{m-2})(1-q/\lambda_{m-1})} \\ &= \Omega \frac{1}{(1-q)(1-\lambda_2q^2)(1-q\lambda_3/\lambda_2)\cdots(1-q\lambda_{m-1}/\lambda_{m-2})(1-q/\lambda_{m-1})} \\ &= \Omega \frac{1}{(1-q)(1-q^2)(1-q^3\lambda_3)\cdots(1-q\lambda_{m-1}/\lambda_{m-2})(1-q/\lambda_{m-1})} \\ &= \cdots \frac{1}{(1-q)(1-q^2)\cdots(1-q^m)}, \end{aligned}$$

and so the proof of Lemma 2.1.4 has been completed.  $\square$

**Corollary 2.1.5.** *The partitions of  $n$  into no more than  $m$  parts are equinumerous with the partitions of  $n$  into parts  $\leq m$ .*

**Lemma 2.1.6.** *If  $\alpha$  is a nonnegative integer, then*

$$\Omega \frac{\lambda^{-\alpha}}{(1-\lambda x)(1-y/\lambda)} = \frac{x^\alpha}{(1-x)(1-xy)}.$$

**Proof.** Putting  $k = n - m - \alpha$  below, we find that

$$\begin{aligned} \Omega \frac{\lambda^{-\alpha}}{(1-\lambda x)(1-y/\lambda)} &= \Omega \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} x^n y^m \lambda^{n-m-\alpha} \\ &= \sum_{m=0}^{\infty} \sum_{n \geq m+\alpha} x^n y^m \\ &= \sum_{k=0}^{\infty} x^{m+k+\alpha} \sum_{m=0}^{\infty} y^m. \end{aligned}$$

Summing the geometric series on the right-hand side above, we complete the proof.  $\square$

**Definition 2.1.7.** *Let  $\Delta(n)$  denote the number of incongruent triangles with positive integral sides and perimeter  $n$ .*

**Theorem 2.1.8.** *A generating function for  $\Delta(n)$  is given by*

$$\sum_{n=0}^{\infty} \Delta(n)q^n = \frac{q^3}{(1-q^2)(1-q^3)(1-q^4)}. \quad (2.1.3)$$

**Proof.** Let  $n_1 \geq n_2 \geq n_3$  denote the lengths of the sides of the triangle. Note that  $n_2 + n_3 \geq n_1 + 1$ . We make three applications of Lemma 2.1.6. In the first instance, replace  $\lambda$  by  $\lambda_1$ ,  $x$  by  $q/\lambda_3$ , and  $y$  by  $q\lambda_2\lambda_3$ . In the second, replace  $\lambda$  by  $\lambda_2$ ,  $x$  by  $q^2$ , and

$y$  by  $q\lambda_3$ . In the third application, replace  $\lambda$  by  $\lambda_3$ ,  $x$  by  $q^3$ ,  $y$  by  $q$ , and  $\alpha$  by 1. Hence,

$$\begin{aligned}
\sum_{n=0}^{\infty} \Delta(n)q^n &= \Omega_{\geq} \sum_{n_1, n_2, n_3=0}^{\infty} q^{n_1+n_2+n_3} \lambda_1^{n_1-n_2} \lambda_2^{n_2-n_3} \lambda_3^{n_2+n_3-n_1-1} \\
&= \Omega_{\geq} \frac{\lambda_3^{-1}}{(1-q\lambda_1/\lambda_3)(1-q\lambda_2\lambda_3/\lambda_1)(1-q\lambda_3/\lambda_2)} \\
&= \Omega_{\geq} \frac{\lambda_3^{-1}}{(1-q/\lambda_3)(1-q^2\lambda_2)(1-q\lambda_3/\lambda_2)} \\
&= \Omega_{\geq} \frac{\lambda_3^{-1}}{(1-q/\lambda_3)(1-q^2)(1-q^3\lambda_3)} \\
&= \frac{q^3}{(1-q^3)(1-q^4)(1-q^2)}.
\end{aligned}$$

□

**Corollary 2.1.9.**  $\Delta(n)$  is equal to the number of partitions of  $n$  into 2's, 3's, and 4's, with at least one 3.

**Theorem 2.1.10.** We have

$$\Omega_{\geq} \frac{1}{(1-\lambda x)(1-y_1/\lambda)(1-y_2/\lambda) \cdots (1-y_j/\lambda)} = \frac{1}{(1-x)(1-xy_1) \cdots (1-xy_j)}, \quad (2.1.4)$$

$$\Omega_{\geq} \frac{1}{(1-\lambda x)(1-\lambda y)(1-z/\lambda)} = \frac{1-xyz}{(1-x)(1-y)(1-xz)(1-yz)}, \quad (2.1.5)$$

$$\Omega_{\geq} \frac{1}{(1-\lambda x)(1-\lambda y)(1-z/\lambda^2)} = \frac{1+xyz-x^2yz-xy^2z}{(1-x)(1-y)(1-x^2z)(1-y^2z)}. \quad (2.1.6)$$

**Proof of (2.1.4).** We first observe that the case  $j = 1$  of (2.1.4) is identical with Lemma 2.1.3. Thus, we shall proceed by induction on  $j$ . Suppose that (2.1.4) is true up to and including  $j - 1$ . Next, we check that

$$\frac{1}{(1-y_{j-1}/\lambda)(1-y_j/\lambda)} = \frac{1}{y_{j-1}-y_j} \left( \frac{y_{j-1}}{1-y_{j-1}/\lambda} - \frac{y_j}{1-y_j/\lambda} \right). \quad (2.1.7)$$

Hence, using the foregoing identity and induction, we find that

$$\begin{aligned}
&\Omega_{\geq} \frac{1}{(1-\lambda x)(1-y_1/\lambda)(1-y_2/\lambda) \cdots (1-y_j/\lambda)} \\
&= \frac{1}{y_{j-1}-y_j} \Omega_{\geq} \left( \frac{y_{j-1}}{(1-\lambda x)(1-y_1/\lambda) \cdots (1-y_{j-2}/\lambda)(1-y_{j-1}/\lambda)} \right. \\
&\quad \left. - \frac{y_j}{(1-\lambda x)(1-y_1/\lambda) \cdots (1-y_{j-2}/\lambda)(1-y_j/\lambda)} \right) \\
&= \frac{1}{y_{j-1}-y_j} \Omega_{\geq} \left( \frac{y_{j-1}}{(1-x)(1-xy_1) \cdots (1-xy_{j-2})(1-xy_{j-1})} \right)
\end{aligned}$$

$$\begin{aligned}
& - \frac{y_j}{(1-x)(1-xy_1)\cdots(1-xy_{j-2})(1-xy_j)} \\
& = \frac{1}{(1-x)(1-xy_1)\cdots(1-xy_j)},
\end{aligned}$$

and so the proof of (2.1.4) is complete.  $\square$

**Proof.** (2.1.5). Using (2.1.7) and Lemma 2.1.3, we find that

$$\begin{aligned}
& \Omega \frac{1}{(1-\lambda x)(1-\lambda y)(1-z/\lambda)} \\
& = \Omega \frac{1}{\geq x-y} \left( \frac{x}{1-\lambda x} - \frac{y}{1-\lambda y} \right) \frac{1}{1-z/\lambda} \\
& = \frac{x}{(x-y)(1-x)(1-xz)} - \frac{y}{(x-y)(1-y)(1-yz)} \\
& = \frac{x(1-y)(1-yz) - y(1-x)(1-xz)}{(x-y)(1-x)(1-y)(1-xz)(1-yz)} \\
& = \frac{x + xy^2z - y - x^2yz}{(x-y)(1-x)(1-y)(1-xz)(1-yz)} \\
& = \frac{(x-y)(1-xyz)}{(x-y)(1-x)(1-y)(1-xz)(1-yz)}.
\end{aligned}$$

$\square$

We leave the proof of (2.1.6) as an exercise.

Recall that Theorem 1.2.23 provides a generating function for  $Q_m(n)$ . We shall use MacMahon's partition analysis to give an alternative proof of Theorem 1.2.23.

**Second Proof of Theorem 1.2.23** Let  $n = n_1 + n_2 + \cdots + n_m$ , and suppose that  $n_j \geq n_{j+1} + 1$ ,  $1 \leq j \leq m-1$ . Also assume that  $n_m \geq 1$ . Then, with several applications of Lemma 2.1.6 with  $\alpha = 1$ ,

$$\begin{aligned}
\sum_{n=0}^{\infty} Q_m(n)q^n & = \Omega \sum_{n_1, n_2, \dots, n_m=0}^{\infty} q^{n_1+n_2+\cdots+n_m} \lambda_1^{n_1-n_2-1} \lambda_2^{n_2-n_3-1} \cdots \lambda_{m-1}^{n_{m-1}-n_m-1} \lambda_m^{n_m-1} \\
& = \Omega \frac{\lambda_1^{-1} \lambda_2^{-1} \cdots \lambda_m^{-1}}{\geq (1-\lambda_1 q)(1-\lambda_2 q/\lambda_1) \cdots (1-\lambda_m q/\lambda_{m-1})} \\
& = \Omega \frac{q \lambda_2^{-1} \cdots \lambda_m^{-1}}{\geq (1-q)(1-\lambda_2 q^2)(1-\lambda_3 q/\lambda_2) \cdots (1-\lambda_m q/\lambda_{m-1})} \\
& = \Omega \frac{q \cdot q^2 \lambda_3^{-1} \cdots \lambda_m^{-1}}{\geq (1-q)(1-q^2)(1-q^3 \lambda_3)(1-\lambda_4 q/\lambda_3) \cdots (1-\lambda_m q/\lambda_{m-1})} \\
& = \cdots \Omega \frac{q \cdot q^2 \cdot q^3 \cdots q^{m-1} \lambda_m^{-1}}{\geq (1-q)(1-q^2) \cdots (1-q^{m-1})(1-q^m \lambda_m)} \\
& = \frac{q \cdot q^2 \cdot q^3 \cdots q^{m-1} \cdot q^m}{(1-q)(1-q^2) \cdots (1-q^{m-1})(1-q^m)},
\end{aligned}$$

where in our last application of Lemma 2.1.6,  $y = 0$ . Combining the powers of  $q$  in the last line above, we complete our second proof of Theorem 1.2.23.  $\square$

**Definition 2.1.11.** Let  $Q_m^{(k,\ell)}(n)$  denote the number of partitions of  $n$  into exactly  $m$  distinct parts, where each part (in descending order) differs from the next by at least  $k$ , and where the smallest part is  $\geq \ell$ .

**Theorem 2.1.12.** For  $Q_m^{(k,\ell)}(n)$  as defined above,

$$\sum_{n=0}^{\infty} Q_m^{(k,\ell)}(n)q^n = \frac{q^{\ell m + km(m-1)/2}}{(q; q)_m}.$$

Note that when  $k = \ell = 1$ , Theorem 2.1.12 reduces to Theorem 1.2.23.

**Proof.** We observe that

$$\begin{aligned} \sum_{n=0}^{\infty} Q_m^{(k,\ell)}(n)q^n &= \Omega_{\geq} \sum_{n_1, n_2, \dots, n_m=0}^{\infty} q^{n_1 + n_2 + \dots + n_m} \lambda_1^{n_1 - n_2 - k} \lambda_2^{n_2 - n_3 - k} \dots \lambda_{m-1}^{n_{m-1} - n_m - k} \lambda_m^{n_m - \ell} \\ &= \Omega_{\geq} \frac{\lambda_1^{-k} \lambda_2^{-k} \dots \lambda_m^{-\ell}}{(1 - \lambda_1 q)(1 - \lambda_2 q / \lambda_1) \dots (1 - \lambda_m q / \lambda_{m-1})}. \end{aligned}$$

The remainder of the proof follows exactly along the same lines as the proof for Theorem 1.2.23 that we gave above, and so we leave it as an exercise for readers.  $\square$

**Definition 2.1.13.** We let  $p_m(j, n)$  denote the number of partitions of  $n$  into at most  $m$  parts, with the largest part being  $j$ . Let  $Q_m(j, n)$  equal the number of partitions of  $n$  into exactly  $m$  distinct parts, with the largest part being  $j$ .

**Theorem 2.1.14.** We have

$$\sum_{j, n=0}^{\infty} p_m(j, n)z^j q^n = \frac{1}{(zq; q)_m}.$$

**Proof.** Note that

$$\begin{aligned} \sum_{j, n=0}^{\infty} p_m(j, n)z^j q^n &= \Omega_{\geq} \sum_{n_1, n_2, \dots, n_m=0}^{\infty} z^{n_1} q^{n_1 + n_2 + \dots + n_m} \lambda_1^{n_1 - n_2} \lambda_2^{n_2 - n_3} \dots \lambda_{m-1}^{n_{m-1} - n_m} \\ &= \Omega_{\geq} \frac{1}{(1 - zq\lambda_1)(1 - q\lambda_2/\lambda_1) \dots (1 - q\lambda_{m-1}/\lambda_{m-2})(1 - q/\lambda_{m-1})}. \end{aligned}$$

We now proceed as in the proof of Theorem 1.2.23, and because the details are similar, we shall forego giving them.  $\square$

**Theorem 2.1.15.** We have

$$\sum_{j, n=0}^{\infty} Q_m(j, n)z^j q^n = \frac{z^m q^{m(m+1)/2}}{(zq; q)_m}.$$

**Proof.** We first observe that we can write the generating function for  $Q_m(j, n)$  in the form

$$\begin{aligned} & \sum_{j,n=0}^{\infty} Q_m(j, n) z^j q^n \\ &= \Omega_{\substack{\geq \\ n_1, n_2, \dots, n_m=0}} \sum_{n_1, n_2, \dots, n_m=0}^{\infty} z^{n_1} q^{n_1+n_2+\dots+n_m} \lambda_1^{n_1-n_2-1} \lambda_2^{n_2-n_3-1} \dots \lambda_{m-1}^{n_{m-1}-n_m-1} \lambda_m^{n_m-1}. \end{aligned}$$

The remainder of the argument is similar to that given in our second proof of Theorem 1.2.23, earlier in this chapter, and so we omit it.  $\square$

## 2.2. Elementary Partition Identities Involving Gaussian Binomial Coefficients

**Definition 2.2.1.** Let  $p(N, M, n)$  denote the number of partitions of  $n$  into at most  $M$  parts, each  $\leq N$ . Let  $Q(N, M, n)$  denote the number of partitions of  $n$  into exactly  $M$  distinct parts, each  $\leq N$ .

**Definition 2.2.2.** The Gaussian coefficient, or the  $q$ -binomial coefficient, or the  $q$ -Gaussian polynomial, is defined by

$$\begin{bmatrix} n \\ m \end{bmatrix}_q = \begin{cases} \frac{(q; q)_n}{(q; q)_m (q; q)_{n-m}}, & 0 \leq m \leq n, \\ 0, & \text{otherwise.} \end{cases}$$

**Definition 2.2.3.** Define an operator  $[z^j]$  by

$$[z^j] \sum_{n=0}^{\infty} a_n z^n = a_j.$$

Note that

$$\begin{aligned} \sum_{j=0}^N a_j &= \sum_{j=0}^N [z^j] \sum_{n=0}^{\infty} a_n z^n = [z^N] \sum_{j=0}^{\infty} z^j \sum_{n=0}^{\infty} a_n z^n \\ &= [z^N] \frac{1}{1-z} \sum_{n=0}^{\infty} a_n z^n. \end{aligned} \tag{2.2.1}$$

**Theorem 2.2.4.** Recall that  $p(N, M, n)$  and  $Q(N, M, n)$  are defined in Definition 2.2.1. Then

$$\sum_{n=0}^{\infty} p(N, M, n) q^n = \begin{bmatrix} M+N \\ M \end{bmatrix}_q, \tag{2.2.2}$$

$$\sum_{n=0}^{\infty} Q(N, M, n) q^n = q^{M(M+1)/2} \begin{bmatrix} N \\ M \end{bmatrix}_q. \tag{2.2.3}$$

We are putting the cart before the horse by offering the following proof, because the primary ingredient in the proof is the  $q$ -binomial theorem, Theorem 3.1.2, the initial theorem in Chapter 3.

**Proof.** We prove (2.2.2). First note that

$$[z^h] \sum_{j,n=0}^{\infty} p_M(j, n) z^j q^n = \sum_{n=0}^{\infty} p_M(j, n) q^n \quad (2.2.4)$$

and

$$\sum_{h=0}^N p_M(h, n) = p(N, M, n). \quad (2.2.5)$$

Thus, using (2.2.5), (2.2.4), Theorem 2.1.14, (2.2.1), and finally the  $q$ -binomial theorem, Theorem 3.1.2, we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} p(N, M, n) q^n &= \sum_{n=0}^{\infty} \sum_{h=0}^N p_M(h, n) q^n \\ &= \sum_{h=0}^N [z^h] \sum_{j,n=0}^{\infty} p_M(j, n) z^j q^n \\ &= \sum_{h=0}^N [z^h] \frac{1}{(zq; q)_M} \\ &= [z^N] \frac{1}{(z; q)_{M+1}} \\ &= [z^N] \frac{(zq^{M+1}; q)_{\infty}}{(z; q)_{\infty}} \\ &= [z^N] \sum_{k=0}^{\infty} \frac{(q^{M+1}; q)_k}{(q; q)_k} z^k \\ &= [z^N] \sum_{k=0}^{\infty} \frac{(q^{M+1}; q)_k (q; q)_M}{(q; q)_k (q; q)_M} z^k \\ &= [z^N] \sum_{k=0}^{\infty} \frac{(q; q)_{M+k}}{(q; q)_k (q; q)_M} z^k \\ &= [z^N] \sum_{k=0}^{\infty} \begin{bmatrix} M+k \\ M \end{bmatrix}_q z^k \\ &= \begin{bmatrix} M+N \\ M \end{bmatrix}_q, \end{aligned}$$

and this completes the proof of (2.2.2).

We now turn to the proof of (2.2.3). Utilizing analogues of (2.2.4) and (2.2.5), Theorem 2.1.15, (2.2.1), and finally a corollary of the  $q$ -binomial theorem, Theorem



3.1.2, we arrive at

$$\begin{aligned}
\sum_{n=0}^{\infty} Q(N, M, n)q^n &= \sum_{h=0}^N [z^h] \sum_{j,n=0}^{\infty} Q_M(j, n)z^j q^n \\
&= \sum_{h=0}^N [z^h] \frac{z^M q^{M(M+1)/2}}{(zq; q)_M} \\
&= [z^N] \frac{z^M q^{M(M+1)/2}}{(z; q)_{M+1}} \\
&= [z^N] z^M q^{M(M+1)/2} \sum_{k=0}^{\infty} \begin{bmatrix} M+k \\ M \end{bmatrix}_q z^k \\
&= q^{M(M+1)/2} \begin{bmatrix} N \\ M \end{bmatrix}_q,
\end{aligned}$$

where we chose the  $k = (N - M)$ th coefficient of the series.  $\square$

**Theorem 2.2.5.** *We have*

$$\sum_{n,M=0}^{\infty} Q(N, M, n)z^M q^n = (-zq; q)_N. \quad (2.2.6)$$

**Proof.** Applying (2.2.3), we arrive at

$$\begin{aligned}
\sum_{n,M=0}^{\infty} Q(N, M, n)z^M q^n &= \sum_{M=0}^{\infty} z^M \sum_{n=0}^{\infty} Q(N, M, n)q^n \\
&= \sum_{M=0}^{\infty} z^M q^{M(M+1)/2} \begin{bmatrix} N \\ M \end{bmatrix}_q.
\end{aligned} \quad (2.2.7)$$

We need to put the Gaussian binomial coefficients in a different form. To that end,

$$\begin{aligned}
\begin{bmatrix} N \\ M \end{bmatrix}_q &= \frac{(q)_N}{(q)_M (q)_{N-M}} = \frac{(q^{N-M+1})_M}{(q)_M} \\
&= \frac{(1 - q^{N-M+1})(1 - q^{N-M+2}) \cdots (1 - q^N)}{(q)_M} \\
&= (-1)^M q^{NM - M(M-1)/2} \frac{(1 - q^{-N+M-1})(1 - q^{-N+M-2}) \cdots (1 - q^{-N})}{(q)_M} \\
&= (-1)^M q^{NM - M(M-1)/2} \frac{(q^{-N})_M}{(q)_M}.
\end{aligned}$$

Employing the calculation above in (2.2.7), we deduce that

$$\begin{aligned}
\sum_{n,M=0}^{\infty} Q(N, M, n)z^M q^n &= \sum_{M=0}^{\infty} \frac{(q^{-N})_M}{(q)_M} (-zq^{N+1})^M \\
&= \frac{(-zq)_{\infty}}{(-zq^{N+1})_{\infty}} = (-zq)_N,
\end{aligned}$$

where we applied the  $q$ -binomial theorem, Theorem 3.1.2, in the penultimate equality.  $\square$

Generally, if the base  $q$  is clear, the subscript  $q$  is deleted from the Gaussian coefficient.

### 2.3. Exercises

1. Find a combinatorial proof of Corollary 2.1.9.

2. Prove (2.1.6).

3. Prove that

$$\sum_{m,n=0}^{\infty} Q_m^{(2,1)}(n) z^m q^n = \sum_{m=0}^{\infty} \frac{z^m q^{m^2}}{(q; q)_m}.$$

4. Prove that

$$\sum_{m,n=0}^{\infty} Q_m^{(2,2)}(n) z^m q^n = \sum_{m=0}^{\infty} \frac{z^m q^{m^2+m}}{(q; q)_m}.$$

When  $z = 1$ , the two functions on the right-hand sides in Exercises 3,4 are the Rogers–Ramanujan functions.

5. Prove the *two* analogues of Pascal's formula for the ordinary binomial coefficients:

$$\begin{bmatrix} n \\ m \end{bmatrix} = \begin{bmatrix} n-1 \\ m \end{bmatrix} + q^{n-m} \begin{bmatrix} n-1 \\ m-1 \end{bmatrix}, \quad (2.3.1)$$

$$\begin{bmatrix} n \\ m \end{bmatrix} = \begin{bmatrix} n-1 \\ m-1 \end{bmatrix} + q^m \begin{bmatrix} n-1 \\ m \end{bmatrix}. \quad (2.3.2)$$

6. Prove that

$$\lim_{q \rightarrow 1} \begin{bmatrix} n \\ m \end{bmatrix} = \binom{n}{m}.$$

---

## Chapter 3

# Some Primary Theorems in the Theory of $q$ -Series

### 3.1. Introduction

As we remarked in Chapter 1, a  $q$ -series, sometimes also dubbed an Eulerian series, generally, but not always, has at least one  $q$ -product in its summands. There is one class of  $q$ -series, called *basic hypergeometric series*, for which an enormous and beautiful theory has been developed. We now define a  ${}_r\phi_s$  basic hypergeometric series.

**Definition 3.1.1.** *If  $r$  and  $s$  are nonnegative integers and  $|q| < 1$ , then*

$$\begin{aligned} {}_r\phi_s(a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_s; q, z) &\equiv_r \phi_s \left[ \begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix}; q, z \right] \\ &:= \sum_{n=0}^{\infty} \frac{(a_1; q)_n (a_2; q)_n \cdots (a_r; q)_n}{(b_1; q)_n (b_2; q)_n \cdots (b_s; q)_n (q; q)_n} \left[ (-1)^n q^{n(n-1)/2} \right]^{1+s-r} z^n, \quad |z| < 1. \end{aligned} \quad (3.1.1)$$

Of course, conditions, depending on  $r$ ,  $s$ , and other parameters, need to be imposed for existence and convergence. Note that when  $r = s + 1$ , the expression involving square brackets is identically equal to 1. In these notes, all of the basic hypergeometric series that appear will be instances when  $r = s + 1$ , in which case the series converges for  $|z| < 1$ . We also have convergence on  $|z| = 1$  if  $\operatorname{Re}(b_1 + \cdots + b_s - (a_1 + \cdots + a_r)) > 0$ . For a fuller discussion of convergence, consult Gasper and Rahman's text [59, pp. 4–5].

The most useful theorem in  $q$ -series is the  $q$ -binomial theorem.

**Theorem 3.1.2** ( $q$ -analogue of the binomial theorem). *For  $|q|, |z| < 1$ ,*

$$\sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} z^n = \frac{(az)_{\infty}}{(z)_{\infty}}. \quad (3.1.2)$$

**First Proof of Theorem 3.1.2.** Note that the product on the right side of (3.1.2) converges uniformly on compact subsets of  $|z| < 1$  and so represents an analytic function on  $|z| < 1$ . Thus, we may write

$$F(z) := \frac{(az)_\infty}{(z)_\infty} = \sum_{n=0}^{\infty} A_n z^n, \quad |z| < 1. \quad (3.1.3)$$

From the product representation in (3.1.3), we can readily verify that

$$(1-z)F(z) = (1-az)F(qz). \quad (3.1.4)$$

Equating coefficients of  $z^n$ ,  $n \geq 1$ , on both sides of (3.1.4), we find that

$$A_n - A_{n-1} = q^n A_n - aq^{n-1} A_{n-1},$$

or

$$A_n = \frac{1 - aq^{n-1}}{1 - q^n} A_{n-1}, \quad n \geq 1. \quad (3.1.5)$$

Iterating (3.1.5) and using the value  $A_0 = 1$ , which is readily apparent from (3.1.3), we deduce that

$$A_n = \frac{(a)_n}{(q)_n}, \quad n \geq 0. \quad (3.1.6)$$

Using (3.1.6) in (3.1.3), we complete the proof of (3.1.2).  $\square$

**Second Proof of Theorem 3.1.2.** Define

$$f_a(z) := \sum_{k=0}^{\infty} \frac{(a)_k}{(q)_k} z^k. \quad (3.1.7)$$

Then

$$\begin{aligned} \frac{f_a(z) - f_a(qz)}{z} &= \sum_{k=0}^{\infty} \left( \frac{(a)_k}{(q)_k} z^{k-1} - \frac{(a)_k}{(q)_k} q^k z^{k-1} \right) \\ &= \sum_{k=0}^{\infty} \frac{(a)_k}{(q)_k} (1 - q^k) z^{k-1} \\ &= (1-a) \sum_{k=1}^{\infty} \frac{(aq)_{k-1}}{(q)_{k-1}} z^{k-1} \\ &= (1-a) f_{aq}(z). \end{aligned} \quad (3.1.8)$$

Rearrange (3.1.8) to achieve the form

$$f_a(z) - f_a(qz) = (1-a)z f_{aq}(z). \quad (3.1.9)$$

Next, consider

$$\begin{aligned} f_a(z) - f_{aq}(z) &= \sum_{k=1}^{\infty} \frac{(aq)_{k-1}}{(q)_k} \{(1-a) - (1-aq^k)\} z^k \\ &= -a \sum_{k=1}^{\infty} \frac{(aq)_{k-1}}{(q)_{k-1}} z^k = -az f_{aq}(z). \end{aligned}$$

In other words,

$$f_a(z) = (1 - az)f_{aq}(z). \quad (3.1.10)$$

From (3.1.9) and (3.1.10),

$$f_a(z) - f_a(qz) = \frac{(1-a)z}{1-az} f_a(z),$$

which we can further write in the form

$$f_a(z) = \frac{1-az}{1-z} f_a(qz). \quad (3.1.11)$$

If we iterate (3.1.11) a total of  $n$  times, we find that

$$f_a(z) = \frac{(az)_n}{(z)_n} f_a(q^n z).$$

Letting  $n \rightarrow \infty$ , we conclude that

$$f_a(z) = \frac{(az)_\infty}{(z)_\infty} f_a(0) = \frac{(az)_\infty}{(z)_\infty}, \quad (3.1.12)$$

because, from (3.1.7),  $f_a(0) = 1$ . Hence, (3.1.2) has been proved.  $\square$

We now state two useful corollaries of Theorem 3.1.2.

**Corollary 3.1.3.** For  $|q|, |z| < 1$ ,

$$\sum_{n=0}^{\infty} \frac{z^n}{(q)_n} = \frac{1}{(z)_\infty}. \quad (3.1.13)$$

**Proof.** Equality (3.1.13) is an immediate consequence of (3.1.2) by setting  $a = 0$ .  $\square$

Suppose that we write (3.1.13) in the form

$$(1-z) \sum_{n=0}^{\infty} \frac{z^n}{(q)_n} = \sum_{m=0}^{\infty} z^m \left\{ \frac{1}{(q)_m} - \frac{1}{(q)_{m-1}} \right\} = \frac{1}{(zq)_\infty}, \quad (3.1.14)$$

where we define  $1/(q)_{-1} = 0$ . Recall that the generating function for  $p(n)$  is  $1/(q)_\infty$ . Thus, the function on the far right side of (3.1.14) generates  $p(n)$  with the power of  $z$  keeping track of the number of parts in each partition. However, the number of partitions of  $n$  with  $m$  parts equals the number of partitions of  $n$  with largest part  $m$ . Note that, indeed, the middle expression in (3.1.14) is generating the partitions of  $n$  with largest part  $m$ .

**Corollary 3.1.4.** For  $z \in \mathbb{Z}$ ,

$$\sum_{n=0}^{\infty} \frac{(-z)^n q^{n(n-1)/2}}{(q)_n} = (z)_\infty. \quad (3.1.15)$$

**Proof.** Replace  $a$  by  $a/b$  and  $z$  by  $bz$  in (3.1.2) to find that, for  $|bz| < 1$ ,

$$\sum_{n=0}^{\infty} \frac{(a/b)_n}{(q)_n} (bz)^n = \frac{(az)_{\infty}}{(bz)_{\infty}}. \quad (3.1.16)$$

Now let  $b \rightarrow 0$ . We note that

$$\begin{aligned} \lim_{b \rightarrow 0} (a/b)_n b^n &= \lim_{b \rightarrow 0} \left(1 - \frac{a}{b}\right) \left(1 - \frac{aq}{b}\right) \cdots \left(1 - \frac{aq^{n-1}}{b}\right) b^n \\ &= (-a)^n q^{n(n-1)/2}. \end{aligned} \quad (3.1.17)$$

With the use of (3.1.17), equality (3.1.15) now follows upon setting  $a = 1$ .  $\square$

In letting  $b \rightarrow 0$  in the proof above, we have swept the details “under the rug.” We give a complete justification, as in [29, pp. 9–10]. There will be many instances in the sequel where we will encounter further interchanges of limiting processes. In each case, a rigorous argument can be constructed along the lines as we will do below.

Let  $|q| \leq M < 1$ . Fix  $\epsilon > 0$  such that  $0 < 2\epsilon < 1 - M$ . Suppose that  $|b| < \epsilon$ . Let  $N_0$  be the unique non-negative integer such that

$$\begin{aligned} |b| + |a|M^k &\geq 2\epsilon, \quad 0 \leq k \leq N_0, \\ |b| + |a|M^{N_0} &< 2\epsilon. \end{aligned}$$

Then, for  $|b| \leq \epsilon$  and  $n \geq N_0$ ,

$$\begin{aligned} \left| \frac{(a/b)_n}{q^n} b^n \right| &\leq \frac{1}{(1-M)^n} \prod_{k=0}^{n-1} (|b| + |a|M^k) \\ &\leq \frac{(\epsilon + |a|)^{N_0} (2\epsilon)^{n-N_0}}{(1-M)^n} \\ &= \left( \frac{\epsilon + |a|}{2\epsilon} \right)^{N_0} \left( \frac{2\epsilon}{1-M} \right)^n. \end{aligned}$$

Since  $2\epsilon < 1 - M$ ,

$$\sum_{n=N_0}^{\infty} \left( \frac{2\epsilon}{1-M} \right)^n < \infty.$$

Thus,

$$\sum_{n=0}^{\infty} \frac{(a/b)_n}{(q)_n} (bz)^n$$

converges uniformly on  $|b| \leq \epsilon$  and for fixed  $z$  by the Weierstrass  $M$ -test. Thus, letting  $b \rightarrow 0$  under the summation sign is justified.

In Chapter 1, we defined Ramanujan’s general theta function  $f(a, b)$  and stated the Jacobi triple product identity. For convenience, we restate them again here. Let

$$f(a, b) := \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \quad |ab| < 1. \quad (3.1.18)$$

The most useful theorem in the theory of theta functions is indeed the Jacobi triple product identity [27, p. 35, Entry 19], [29, Theorem 1.3.3, p. 10].

**Theorem 3.1.5.** *For each theta function  $f(a, b)$ ,*

$$f(a, b) = (-a; ab)_\infty (-b; ab)_\infty (ab; ab)_\infty. \quad (3.1.19)$$

The first proof that we give is based on the  $q$ -binomial theorem and a theorem of Rothe, which in fact is just another version of Theorem 2.2.5.

**Theorem 3.1.6** (Rothe). *For any positive integer  $N$ ,*

$$\sum_{j=0}^N \begin{bmatrix} N \\ j \end{bmatrix} (-1)^j x^j q^{j(j-1)/2} = (x; q)_N. \quad (3.1.20)$$

**Proof.** In the  $q$ -binomial theorem, Theorem 3.1.2, replace  $z$  by  $xq^N$  and let  $a = q^{-N}$  to find that

$$\sum_{j=0}^{\infty} \frac{(q^{-N})_j}{(q)_j} x^j q^{Nj} = \frac{(x)_\infty}{(xq^N)_\infty} = (x; q)_N. \quad (3.1.21)$$

We leave it to readers to verify that

$$(q^{-N})_j = (-1)^j q^{-Nj+j(j-1)/2} \frac{(q)_N}{(q)_{N-j}}. \quad (3.1.22)$$

Putting (3.1.22) in (3.1.21), we deduce that

$$(x; q)_N = \sum_{j=0}^{\infty} \frac{(q)_N}{(q)_j (q)_{N-j}} (-1)^j x^j q^{j(j-1)/2} = \sum_{j=0}^N \begin{bmatrix} N \\ j \end{bmatrix} (-1)^j x^j q^{j(j-1)/2}. \quad (3.1.23)$$

Thus, the proof of (3.1.20) is complete.  $\square$

**Proof of Theorem 3.1.5.** In Theorem 3.1.6, let  $N = 2n$  and replace the index  $j$  by  $k + n$  to see that

$$\sum_{k=-n}^n \begin{bmatrix} 2n \\ k+n \end{bmatrix} (-1)^{k+n} q^{(k+n)(k+n-1)/2} x^{k+n} = (x; q)_{2n}. \quad (3.1.24)$$

Replace  $x$  by  $xq^{-n}$  in (3.1.24) and use a calculation almost identical to that in (3.1.23) to find that

$$(xq^{-n}; q)_{2n} = (xq^{-n}; q)_n (x; q)_n = (-1)^n q^{-n(n+1)/2} x^n (q/x; q)_n (x; q)_n. \quad (3.1.25)$$

Thus, from (3.1.24), with the use of (3.1.25),

$$\sum_{k=-n}^n \frac{(q; q)_{2n} (-1)^{k+n} q^{(k+n)(k+n-1)/2} x^{k+n} q^{-n(k+n)}}{(q; q)_{n+k} (q; q)_{n-k}} = (-1)^n q^{-n(n+1)/2} x^n (q/x; q)_n (x; q)_n,$$

which can be put in the more simplified form

$$\sum_{k=-n}^n \frac{(q; q)_{2n} (-1)^k q^{k(k-1)/2} x^k}{(q; q)_{n+k} (q; q)_{n-k}} = (q/x; q)_n (x; q)_n. \quad (3.1.26)$$

Letting  $n \rightarrow \infty$  in (3.1.26), and using Tannery's Theorem or the Dominated Convergence Theorem to justify taking the limit inside the summation sign, we conclude that

$$\sum_{k=-\infty}^{\infty} \frac{(-1)^k q^{k(k-1)/2} x^k}{(q; q)_{\infty}} = (q/x; q)_{\infty} (x; q)_{\infty}, \quad (3.1.27)$$

which is easily seen to be equivalent to (3.1.19).  $\square$

We offer three corollaries of Theorem 3.1.5. If we set  $x = -q$  in (3.1.27) and replace  $k$  by  $-k-1$  for the negative indexed terms on the left-hand side, and use Theorem 1.2.10, we deduce the following corollary.

**Corollary 3.1.7.** *We have*

$$\psi(q) := \sum_{k=0}^{\infty} q^{k(k+1)/2} = (-q; q)_{\infty}^2 (q; q)_{\infty} = \frac{(q^2; q^2)_{\infty}}{(q; q^2)_{\infty}}. \quad (3.1.28)$$

We have introduced Ramanujan's notation  $\psi(q)$  on the left side of (3.1.28). If we set  $a = b = q$  in (3.1.19), we obtain the following corollary, where we use Ramanujan's notation  $\varphi(q)$ .

**Corollary 3.1.8.** *We have*

$$\varphi(q) := \sum_{n=-\infty}^{\infty} q^{n^2} = (-q; q^2)_{\infty}^2 (q^2; q^2)_{\infty}. \quad (3.1.29)$$

If we replace  $x$  by  $-xq$  in (3.1.27), we see that

$$\sum_{k=0}^{\infty} q^{k(k+1)/2} (x^k + x^{-k-1}) = (-xq; q)_{\infty} (-1/x; q)_{\infty} (q; q)_{\infty}. \quad (3.1.30)$$

Divide both sides of (3.1.30) by  $x+1$ , and then let  $x \rightarrow -1$ , with the aid of L'Hospital's rule, we deduce the next corollary.

**Corollary 3.1.9** (Jacobi's Identity).

$$\sum_{k=0}^{\infty} (-1)^k (2k+1) q^{k(k+1)/2} = (q; q)_{\infty}^3. \quad (3.1.31)$$

The second proof that we give is due to J. J. Sylvester and Hathaway.

**Second Proof of Theorem 3.1.5.** We shall prove (3.1.19) in the form

$$(-zq; q)_{\infty} (-1/z; q)_{\infty} (q; q)_{\infty} = \sum_{n=-\infty}^{\infty} z^n q^{n(n+1)/2}. \quad (3.1.32)$$

Set

$$\phi(z) := (-zq; q)_{\infty} (-1/z; q)_{\infty}. \quad (3.1.33)$$

It is easily verified that  $\phi(z)$  satisfies the functional equation

$$\phi(zq) = \frac{1}{zq} \phi(z). \quad (3.1.34)$$



From its definition (3.1.33), we easily see that  $\phi(z)$  is analytic in a deleted neighborhood of the origin, and so we shall write

$$\phi(z) = \sum_{n=-\infty}^{\infty} c_n z^n, \quad 0 < |z| < \infty. \quad (3.1.35)$$

By (3.1.34),

$$\sum_{n=-\infty}^{\infty} c_n z^n = zq \sum_{n=-\infty}^{\infty} c_n (zq)^n = \sum_{n=-\infty}^{\infty} c_{n-1} (zq)^n, \quad (3.1.36)$$

Equating coefficients of  $z^n$  on both sides of (3.1.36), we deduce the recurrence relation

$$c_n = q^n c_{n-1}. \quad (3.1.37)$$

Assume now that  $n \geq 1$ . By successively iterating (3.1.37), we find that

$$c_n = q^{n(n+1)/2} c_0, \quad n \geq 1. \quad (3.1.38)$$

If  $n \leq 0$ , successive iterations of (3.1.37) yield

$$c_{n-1} = q^{-n(-n+1)/2} c_0,$$

or, with the replacement of  $n$  by  $n+1$ ,

$$c_n = q^{n(n+1)/2} c_0, \quad n \leq -1. \quad (3.1.39)$$

Hence, (3.1.38) and (3.1.39) show that for all integers  $n$

$$c_n = q^{n(n+1)/2} c_0, \quad -\infty < n < \infty. \quad (3.1.40)$$

Hence, we have shown that

$$\phi(z) = c_0 \sum_{n=-\infty}^{\infty} q^{n(n+1)/2} z^n. \quad (3.1.41)$$

There remains the task of computing  $c_0$ .

Now return to the definition of  $\phi(z)$  given in (3.1.33), and note that the constant term  $c_0$  is of course equal to the constant term in the product of the two infinite products on the left-hand side. The constant term arises from multiplying expressions of the type  $zq^{a_j}$  by expressions of the type  $z^{-1}q^{b_k}$ . A typical term will be of the form

$$(zq^{a_1})(zq^{a_2}) \cdots (zq^{a_r})(z^{-1}q^{b_1})(z^{-1}q^{b_2}) \cdots (z^{-1}q^{b_r}), \quad (3.1.42)$$

where  $a_1 > a_2 > \cdots > a_r \geq 1$ ,  $b_1 > b_2 > \cdots > b_r \geq 0$ . To more clearly understand how many of these expressions can arise, we define the Frobenius Symbol.

**Definition 3.1.10.** Consider the Ferrers graph of a partition of a positive integer  $n$ . Let  $r$  be the size of a Durfee square. Form the diagonal of the Durfee square, which will have  $r$  nodes. To the right of the diagonal is a graphical representation of a partition of no more than  $r$  distinct parts, reading from top to bottom, say  $a_1, a_2, \dots, a_r$ . To the left of the diagonal is a graphical representation of another partition of no more than  $r$  distinct parts, reading from left to right, namely  $b_1, b_2, \dots, b_r$ , say. Thus,  $n = r + \sum_{j=1}^r (a_j + b_j)$ . A

matrix representation corresponding to these two partitions can be given by the Frobenius symbol

$$\begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ b_1 & b_2 & \cdots & b_r \end{pmatrix}.$$

Note that, possibly,  $a_r = 0$  and/or  $b_r = 0$ .

As an example, consider the partition  $15 = 5 + 4 + 4 + 2$ . Here,  $r = 3$ . To the right of the diagonal is the partition  $4+2+1$  of 7; below the diagonal is the partition  $3+2$  of 5. Hence, the Frobenius symbol for our original partition is given by

$$\begin{pmatrix} 4 & 2 & 1 \\ 3 & 2 & 0 \end{pmatrix}.$$

We resume our proof of (3.1.19). We see that with each partition of  $n$ , there corresponds a unique Frobenius symbol of the form

$$\begin{pmatrix} a_1 - 1 & a_2 - 1 & \cdots & a_r - 1 \\ b_1 & b_2 & \cdots & b_r \end{pmatrix}.$$

Conversely, each Frobenius symbol is associated with a unique partition. Moreover, by (3.1.19), (3.1.33), and (3.1.42), each Frobenius symbol corresponds to a unique multiplication of series arising from  $r$  terms of the sort  $(1 + zq^n)$ ,  $n \geq 1$ , and  $r$  expressions of the form  $(1 + q^{n-1}/z)$ ,  $n \geq 1$ . Thus, the constant term  $c_0$  arises from all of the partitions of all nonnegative integers  $n$ , i.e.,

$$c_0 = \sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q; q)_{\infty}}. \quad (3.1.43)$$

Returning to our example above, we write  $15 = 3 + \{(5-1) + (3-1) + (2-1)\} + \{3+2+0\}$ , with the corresponding Frobenius symbol being written in the form

$$\begin{pmatrix} 5-1 & 3-1 & 2-1 \\ 3 & 2 & 0 \end{pmatrix}.$$

Putting (3.1.43) into (3.1.41) and noting that our goal was to prove (3.1.32), we have completed our proof of Theorem 3.1.5.  $\square$

**Third Proof of Theorem 3.1.5.** We provide a beautiful bijective proof that is due to E. M. Wright [104].

Recall Ramanujan's definition of  $f(a, b)$  from (1.1.6), in which we set  $a = x, b = y$  and put it in the form

$$(-x; xy)_{\infty}(-y; xy)_{\infty} = \frac{1}{(xy; xy)_{\infty}} \sum_{n=-\infty}^{\infty} x^{n(n+1)/2} y^{n(n-1)/2}. \quad (3.1.44)$$

The left-hand side of (3.1.44) is the generating function for  $\alpha(n, m)$ , the number of bipartite partitions of the forms

$$(a, a-1), (b-1, b), \quad (3.1.45)$$

where  $a$  and  $b$  are positive integers. Before equating coefficients of  $x^n y^m$ ,  $m, n \geq 0$ , on both sides of (3.1.44), we slightly rewrite (3.1.44) in the form

$$\sum_{n,m=0}^{\infty} \alpha(n,m) x^n y^m = \sum_{j=0}^{\infty} p(j) x^j y^j \sum_{r=-\infty}^{\infty} x^{r(r+1)/2} y^{r(r-1)/2}. \quad (3.1.46)$$

Note that

$$j + \frac{1}{2}r(r+1) = n \quad \text{and} \quad j + \frac{1}{2}r(r-1) = m. \quad (3.1.47)$$

Hence,

$$n - \frac{1}{2}r(r+1) = m - \frac{1}{2}r(r-1)$$

, i.e.,

$$r = n - m \quad (3.1.48)$$

Putting (3.1.48) into the first equation of (3.1.47), we find that

$$j = n - \frac{1}{2}(n-m)(n-m+1) \quad (3.1.49)$$

and conclude from (3.1.46) that

$$\alpha(n,m) = p(n - \frac{1}{2}(n-m)(n-m+1)). \quad (3.1.50)$$

Without loss of generality, assume that  $n \geq m$ . Recall from (3.1.48) that  $m = n - r$ . We shall demonstrate that each partition of  $(n, n - r)$  into parts of the form given in (3.1.45) corresponds bijectively with the partition

$$n = \sum_{j=1}^{v+r} a_j + \sum_{j=1}^v (b_j - 1), \quad 1 \leq a_1 < a_2 < \dots, \quad 1 \leq b_1 < b_2 < \dots. \quad (3.1.51)$$

Write  $k = n - \frac{1}{2}r(r+1)$ . Thus, the right-hand side of (3.1.50) is  $p(k)$ . If  $k < 0$ , i.e.,  $n < \frac{1}{2}r(r+1)$ , then  $p(k) = 0$ , as we show forthwith. Now,

$$n \geq \sum_{j=1}^r a_j \geq \sum_{j=1}^r j = \frac{1}{2}r(r+1) \quad (3.1.52)$$

which contradicts the fact that  $n < \frac{1}{2}r(r+1)$ . Therefore, there are no solutions to (3.1.51) if  $k < 0$ . If  $k = 0$ , then  $n = \frac{1}{2}r(r+1)$ . Thus,  $p(k) = p(0) = 1$ , and there exists 1 solution to (3.1.51), namely,  $v = 0$ ,  $a_j = j$ .

Suppose lastly that  $k > 0$ . We will accompany our proof with an example:  $31 = 9 + 9 + 6 + 4 + 2 + 1$ . Above the Ferrers graph of the partition, place a right-angled triangle of  $r$  rows, the lowest of which has  $r$  nodes. The triangle should be situated such that one side of  $r$  nodes is adjacent to the top row of the original Ferrers graph and that the other side of length  $r$  is along the left vertical side of the Ferrers graph. In our example, we take  $r = 2$ . We now have  $n = k + \frac{1}{2}r(r+1)$  nodes. Next, draw a diagonal line lying just above the hypotenuse, extending through the Ferrers graph, and dividing the Ferrers graph into two parts. The set of nodes below the diagonal has  $r + v$  columns, for some particular  $v$ . In our example,  $v = 3$ . The columns have different numbers of nodes, and so we obtain a partition into distinct parts. In our example, we have  $a_1 = 1$ ,  $a_2 = 3$ ,  $a_3 = 4$ ,  $a_4 = 6$ , and  $a_5 = 8$ . To the right of the diagonal, we have a Ferrers graph

of  $v$  rows. Note that the last row might be empty, but the last two rows cannot both be empty. The partition to the right of the diagonal will have  $v$  rows of distinct lengths. We denote the parts of this partition by  $b_1 - 1, b_2 - 1, \dots, b_v - 1$ , where  $1 \leq b_1 < b_2 < \dots$ . In our example,  $b_1 = 2, b_2 = 6$ , and  $b_3 = 7$ . Thus, we compose the partition

$$n = \sum_{j=1}^{v+r} a_j + \sum_{j=1}^v (b_j - 1).$$

This process can be reversed. We start with a solution of (3.1.51) and construct a graph, as in our second graph above. We delete the  $r$  rows or  $\frac{1}{2}r(r+1)$  nodes at the top of the graph, and so obtain a partition of  $k$ . The correspondence is one-to-one, and so we have established (3.1.50), which is the arithmetical interpretation of the Jacobi triple product identity.  $\square$

### 3.2. Some Standard Theorems about

$${}_2\phi_1(a, b; c; q; z)$$

Recall that at the beginning of this chapter, we gave the definition of a general basic hypergeometric series (3.1.1). The theory for the basic hypergeometric function

$${}_2\phi_1(a, b; c; z) := {}_2\phi_1(a, b; c; q, z) := \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(c; q)_n (q; q)_n} z^n, \quad |z| < 1,$$

is more extensive than those for other pairs of indices, but we emphasize that it cannot be separated from the theories for other values of  $r$  and  $s$ . E. Heine [68] was the first to systematically study  ${}_2\phi_1(a, b; c; z)$ , although special cases were studied earlier by Cauchy, Euler, Gauss, and others. We now prove a few basic properties that we shall utilize in the sequel. Several theorems in the  ${}_2\phi_1$  theory are analogues of fundamental theorems in the theory of the ordinary or Gaussian hypergeometric series  ${}_2F_1$ .

**Theorem 3.2.1** (Heine's Transformation). *For  $|q|, |z|, |b| < 1$ ,*

$${}_2\phi_1(a, b; c; z) = \frac{(b)_\infty (az)_\infty}{(c)_\infty (z)_\infty} {}_2\phi_1(c/b, z; az; b). \quad (3.2.1)$$

**Proof.** Using the  $q$ -binomial theorem, Theorem (3.1.2), twice, we find that

$$\begin{aligned} {}_2\phi_1(a, b; c; z) &= \frac{(b)_\infty}{(c)_\infty} \sum_{n=0}^{\infty} \frac{(a)_n z^n (cq^n)_\infty}{(q)_n (bq^n)_\infty} \\ &= \frac{(b)_\infty}{(c)_\infty} \sum_{n=0}^{\infty} \frac{(a)_n z^n}{(q)_n} \sum_{m=0}^{\infty} \frac{(c/b)_m (bq^n)^m}{(q)_m} \\ &= \frac{(b)_\infty}{(c)_\infty} \sum_{m=0}^{\infty} \frac{(c/b)_m b^m}{(q)_m} \sum_{n=0}^{\infty} \frac{(a)_n (zq^m)^n}{(q)_n} \\ &= \frac{(b)_\infty}{(c)_\infty} \sum_{m=0}^{\infty} \frac{(c/b)_m b^m (azq^m)_\infty}{(q)_m (zq^m)_\infty} \end{aligned}$$

$$= \frac{(b)_\infty (az)_\infty}{(c)_\infty (z)_\infty} \sum_{m=0}^{\infty} \frac{(c/b)_m (z)_m}{(q)_m (az)_m} b^m,$$

which is what we wanted to prove.  $\square$

**Theorem 3.2.2** (*q*-analogue of Gauss's Theorem). *If  $|c| < |ab|$  and  $|q| < 1$ , then*

$${}_2\phi_1\left(a, b; c; \frac{c}{ab}\right) = \frac{(c/a)_\infty (c/b)_\infty}{(c/(ab))_\infty (c)_\infty}. \quad (3.2.2)$$

**Proof.** Applying Heine's transformation followed by the *q*-analogue of the binomial theorem, we see that

$$\begin{aligned} {}_2\phi_1\left(a, b; c; \frac{c}{ab}\right) &= \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n (q)_n} \left(\frac{c}{ab}\right)^n \\ &= \frac{(b)_\infty (c/b)_\infty}{(c)_\infty (c/(ab))_\infty} \sum_{n=0}^{\infty} \frac{(c/b)_n (c/(ab))_n}{(c/b)_n (q)_n} b^n \\ &= \frac{(b)_\infty (c/b)_\infty}{(c)_\infty (c/(ab))_\infty} \frac{(c/a)_\infty}{(b)_\infty}. \end{aligned}$$

After cancellation, we obtain the desired result.  $\square$

If we set  $a = q^{-n}$  in Theorem 3.2.2, we obtain the following famous corollary.

**Theorem 3.2.3** (*q*-analogue of the Chu–Vandermonde Theorem). *For each nonnegative integer  $n$ ,*

$${}_2\phi_1\left(q^{-n}, b; c; \frac{cq^n}{b}\right) = \frac{(c/b; q)_n}{(c; q)_n}. \quad (3.2.3)$$

**Theorem 3.2.4** (Bailey's Theorem). *For  $|q| < \min(1, |b|)$ ,*

$${}_2\phi_1(a, b; qa/b; -q/b) = \frac{(aq; q^2)_\infty (-q; q)_\infty (q^2 a/b^2; q^2)_\infty}{(qa/b; q)_\infty (-q/b; q)_\infty}. \quad (3.2.4)$$

**Proof.** Applying Heine's transformation with  $a$  and  $b$  switched and Theorem 3.1.2, we find that

$$\begin{aligned} {}_2\phi_1(a, b; qa/b; -q/b) &= \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(qa/b; q)_n (q)_n} \left(-\frac{q}{b}\right)^n \\ &= \frac{(a; q)_\infty (-q; q)_\infty}{(qa/b; q)_\infty (-q/b; q)_\infty} \sum_{n=0}^{\infty} \frac{(q/b; q)_n (-q/b; q)_n}{(-q; q)_n (q; q)_n} a^n \\ &= \frac{(a; q)_\infty (-q; q)_\infty}{(qa/b; q)_\infty (-q/b; q)_\infty} \sum_{n=0}^{\infty} \frac{(q^2/b^2; q^2)_n}{(q^2; q^2)_n} a^n \\ &= \frac{(a; q)_\infty (-q; q)_\infty}{(qa/b; q)_\infty (-q/b; q)_\infty} \frac{(q^2 a/b^2; q^2)_\infty}{(a; q^2)_\infty} \\ &= \frac{(aq; q^2)_\infty (-q; q)_\infty (q^2 a/b^2; q^2)_\infty}{(qa/b; q)_\infty (-q/b; q)_\infty}. \end{aligned}$$

□

**Corollary 3.2.5.** *We have*

$$\sum_{n=0}^{\infty} \frac{(a)_n}{(q)_n} q^{n(n+1)/2} = (aq; q^2)_{\infty} (-q; q)_{\infty}. \quad (3.2.5)$$

**Proof.** Set  $b = 1/\beta$  in Theorem 3.2.4 and then let  $\beta \rightarrow 0$ . To that end,

$$(aq/b)_n \rightarrow 1, \quad \frac{(b)_n}{b^n} = (1/\beta)_n \beta^n \rightarrow (-1)^n q^{n(n-1)/2}.$$

Then the desired result follows immediately from (3.2.4). □

Note that if we set  $a = q$  in (3.2.5), we obtain the familiar product representation for Ramanujan's function  $\psi(q)$ , defined in (3.1.28), i.e.,

$$\sum_{n=0}^{\infty} q^{n(n+1)/2} = (q^2; q^2)_{\infty} (-q; q)_{\infty} = \frac{(q^2; q^2)_{\infty}}{q; q^2}_{\infty},$$

by Euler's theorem.

**Theorem 3.2.6** (*q-analogue of Euler's Transformation*). *For  $|z|, |abz/c| < 1$ ,*

$${}_2\phi_1(a, b; c; z) = \frac{(abz/c)_{\infty}}{(z)_{\infty}} {}_2\phi_1(c/a, c/b; c; abz/c). \quad (3.2.6)$$

**Proof.** By Heine's transformation (3.2.1),

$${}_2\phi_1(a, b; c; z) = \frac{(b)_{\infty}(az)_{\infty}}{(c)_{\infty}(z)_{\infty}} {}_2\phi_1(c/b, z; az; b).$$

With the roles of  $c/b$  and  $z$  reversed, we apply Heine's transformation a second time to deduce that

$${}_2\phi_1(a, b; c; z) = \frac{(b)_{\infty}(az)_{\infty}}{(c)_{\infty}(z)_{\infty}} \frac{(c/b)_{\infty}(bz)_{\infty}}{(az)_{\infty}(b)_{\infty}} {}_2\phi_1(abz/c, b; bz; c/b). \quad (3.2.7)$$

With the roles of  $abz/c$  and  $b$  reversed, we apply Heine's transformation a third time to obtain our final identity

$$\begin{aligned} {}_2\phi_1(a, b; c; z) &= \frac{(c/b)_{\infty}(bz)_{\infty}}{(c)_{\infty}(z)_{\infty}} \frac{(abz/c)_{\infty}(c)_{\infty}}{(bz)_{\infty}(c/b)_{\infty}} {}_2\phi_1(c/a, c/b; c; abz/c) \\ &= \frac{(abz/c)_{\infty}}{(z)_{\infty}} {}_2\phi_1(c/a, c/b; c; abz/c). \end{aligned} \quad (3.2.8)$$

□

Because (3.2.7) is used so frequently in applications, we are going to elevate it from a step in a proof of a  $q$ -analogue of Euler's transformation to a corollary.

**Corollary 3.2.7** (*Second Iterate of Heine's Transformation*). *We have*

$${}_2\phi_1(a, b; c; z) = \frac{(c/b)_{\infty}(bz)_{\infty}}{(c)_{\infty}(z)_{\infty}} {}_2\phi_1(abz/c, b; bz; c/b). \quad (3.2.9)$$

Our next result is a corollary of Heine's transformation, Theorem 3.2.1.

**Theorem 3.2.8.** For  $|z| < 1$ ,

$$\sum_{n=0}^{\infty} \frac{(z)_n}{(q)_n} a^n q^{n(n+1)/2} = (z)_{\infty} (-aq)_{\infty} \sum_{n=0}^{\infty} \frac{z^n}{(q)_n (-aq)_n}. \quad (3.2.10)$$

**Proof.** Setting  $c = 0$  in Heine's transformation, Theorem 3.2.1, we find that

$$\sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(q)_n} z^n = \frac{(b)_{\infty} (az)_{\infty}}{(z)_{\infty}} \sum_{n=0}^{\infty} \frac{(z)_n}{(az)_n (q)_n} b^n.$$

Let  $z = c$  and then put  $b = z$  above. Thus,

$$\sum_{n=0}^{\infty} \frac{(a)_n (z)_n}{(q)_n} c^n = \frac{(z)_{\infty} (ac)_{\infty}}{(c)_{\infty}} \sum_{n=0}^{\infty} \frac{(c)_n}{(ac)_n (q)_n} z^n. \quad (3.2.11)$$

Next, replace  $a$  by  $-qa/c$  and then let  $c$  tend to 0 in (3.2.11). Noting that

$$\lim_{c \rightarrow 0} (-qa/c)_n c^n = q^{n(n+1)/2} a^n,$$

we readily deduce that 3.2.10 holds.  $\square$

**Second Proof of Theorem 3.2.8.** We begin by writing (3.2.10) in the form

$$\sum_{m=0}^{\infty} \frac{a^m q^{m(m+1)/2}}{(q)_m} \frac{1}{(q^m z)_{\infty}} = \sum_{n=0}^{\infty} \frac{z^n}{(q)_n} (-aq^{n+1})_{\infty}. \quad (3.2.12)$$

Let  $L(n, m, N)$  and  $R(n, m, N)$  denote the coefficients of  $z^n a^m q^N$  on the left- and right-hand sides, respectively, of (3.2.12). A bipartition is an ordered pair of partitions. The two types of partitions need not be of the same type. We show that bipartitions are enumerated on each side of (3.2.12), and that these numbers are equal.

Let  $P_r^k$  be the set of partitions consisting of exactly  $r$  parts, with each part  $\geq k$ . Let  $D_r^k$  be the set of partitions of  $r$  distinct parts, with each part  $\geq k$ . We see that

$$\frac{a^m q^{m(m+1)/2}}{(q)_m} \quad (3.2.13)$$

enumerates partitions with exactly  $m$  distinct parts, i.e., parts in  $D_m^1$ . To see this, we first consider a partition of  $1/(q)_m$  of less than or equal to  $m$  parts and write the parts in decreasing order. Now  $1 + 2 + \cdots + m = m(m+1)/2$ , and so we take our partition from  $1/(q)_m$  and add the parts  $m, m-1, \dots, 1$  to it in decreasing order. We thus have a partition into *exactly*  $m$  distinct parts. Note that the power of  $a$  is the number of parts in the partition. Thus, (3.2.13) generates the partitions in  $D_m^1$ . The terms

$$\frac{1}{(q^m z)_{\infty}} \quad (3.2.14)$$

generates partitions with each part at least equal to  $m$ . The power of  $z$ , say  $n$ , tells us how many parts we have in a particular partition. Thus, (3.2.14) generates partitions in  $P_n^m$ . In summary,  $L(n, m, N)$  is equal to the number of bipartite partitions

$$D_m^1 \oplus P_n^m. \quad (3.2.15)$$

On the right-hand side of (3.2.12), consider first

$$(-aq^{n+1})_\infty, \quad (3.2.16)$$

which generates partitions into distinct parts, with  $n+1$  being the smallest possible part. The power of  $a$  indicates the number of parts. Furthermore,

$$\frac{1}{(q)_n} \quad (3.2.17)$$

generates partitions into less than or equal to  $n$  parts. Take each such partition and borrow, in the language of Ferrers' graphs,  $nm$  nodes from each partition from the partitions generated from (3.2.16), so that instead of having  $m$  parts at least  $n+1$  in size, we now have  $m$  parts at least 1 in size. So now, when we previously had  $\leq n$  parts, by adding  $m$  nodes to each of these  $\leq n$  parts, we now have exactly  $n$  parts, with at least  $m$  in each part. Thus, we are then counting partitions generated by

$$D_m^1 \oplus P_n^m. \quad (3.2.18)$$

Note that this transfer of  $nm$  nodes does not affect the number of parts, i.e., the power of  $a$  indicating the number of parts is the same. The power  $n$  of  $z$  is also exactly the number of parts in  $P_n^m$ . From (3.2.15) and (3.2.18), we complete the proof.  $\square$

### 3.3. The More General ${}_{r+1}\phi_r$

Let us return to Definition 3.1.1 with  $r$  replaced by  $r+1$  and  $s=r$ . Thus, we are going to study

$$\begin{aligned} {}_{r+1}\phi_r(a_1, a_2, \dots, a_{r+1}; b_1, b_2, \dots, b_r; q, z) &\equiv {}_{r+1}\phi_r \left[ \begin{matrix} a_1, a_2, \dots, a_{r+1} \\ b_1, b_2, \dots, b_r \end{matrix}; q, z \right] \\ &:= \sum_{n=0}^{\infty} \frac{(a_1; q)_n (a_2; q)_n \cdots (a_{r+1}; q)_n}{(b_1; q)_n (b_2; q)_n \cdots (b_r; q)_n (q; q)_n} z^n, \quad |z| < 1. \end{aligned} \quad (3.3.1)$$

We begin with one of the classic theorems in basic hypergeometric series, due to Pfaff and Saalschütz. Note that the series on the left-hand side of (3.3.2) below terminates.

**Theorem 3.3.1.** *For each positive integer  $n$ ,*

$${}_3\phi_2 \left( \begin{matrix} a, b, q^{-n} \\ c, abq^{1-n}/c \end{matrix}; q; q \right) = \frac{(c/a)_n (c/b)_n}{(c)_n (c/(ab))_n}. \quad (3.3.2)$$



**Proof.** By the  $q$ -binomial theorem, Theorem 3.1.2,

$$\frac{(abz/c)_\infty}{(z)_\infty} = \sum_{k=0}^{\infty} \frac{(ab/c)_k}{(q)_k} z^k, \quad |z| < 1. \quad (3.3.3)$$

Next, by the  $q$ -analogue of Euler's transformation, Theorem 3.2.6, and (3.3.3),

$$\begin{aligned} {}_2\phi_1(a, b; c; q; z) &= \frac{(abz/c)_\infty}{(z)_\infty} \sum_{m=0}^{\infty} \frac{(c/a)_m (c/b)_m}{(c)_m (q)_m} \left(\frac{abz}{c}\right)^m \\ &= \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \frac{(ab/c)_k}{(q)_k} \frac{(c/a)_m (c/b)_m}{(c)_m (q)_m} \left(\frac{ab}{c}\right)^m z^{m+k}. \end{aligned} \quad (3.3.4)$$

Equating coefficients of  $z^n$ ,  $n \geq 0$ , on both sides of (3.3.4), we find that

$$\frac{(a)_n (b)_n}{(c)_n (q)_n} = \sum_{m=0}^n \frac{(ab/c)_{n-m}}{(q)_{n-m}} \frac{(c/a)_m (c/b)_m}{(c)_m (q)_m} \left(\frac{ab}{c}\right)^m. \quad (3.3.5)$$

It can readily be shown that

$$(a)_{n-m} = \frac{(a)_n}{(q^{1-n}/a)_m} \left(-\frac{q}{a}\right)^m q^{m(m-1)/2 - nm}, \quad (3.3.6)$$

which we leave as an exercise. Applying (3.3.6) with  $a$  replaced by  $ab/c$  and  $q$ , respectively, in (3.3.5), and simplifying, we find that

$$\frac{(a)_n (b)_n}{(c)_n (ab/c)_n} = \sum_{m=0}^n \frac{(q^{-n})_m (c/a)_m (c/b)_m}{(q^{1-n}c/(ab))_m (c)_m (q)_m} q^m. \quad (3.3.7)$$

If we replace the pair  $a, b$  by the pair  $c/a, c/b$ , respectively, in (3.3.7), we deduce (3.3.2) to complete the proof.  $\square$

Our next goal is to utilize the Pfaff–Saalschütz Theorem to derive a general transformation formula that enables us to reduce certain  ${}_{r+2}\phi_{r+1}$  series to  ${}_r\phi_{r-1}$  series. This reduction transformation can be used repeatedly, with the aim of reaching a hypergeometric series that perhaps can be summed in closed form. Our final result can also be viewed as a form of Bailey's Theorem, which is an enormous tool for producing  $q$ -series identities.

Return to Theorem 3.3.1 and replace  $n, a, b$ , and  $c$  by  $k, aq^k, aq/(bc)$ , and  $aq/b$ , respectively. Hence,

$$\begin{aligned} {}_3\phi_2(q^{-k}, aq^k, aq/(bc); aq/b, aq/c; q) \\ = \frac{(q^{1-k}/b)_k (c)_k}{(aq/b)_k (cq^{-k}/a)_k} = \frac{(b)_k (c)_k}{(aq/b)_k (aq/c)_k} \left(\frac{aq}{bc}\right)^k, \end{aligned} \quad (3.3.8)$$

after some elementary manipulation. Multiply (3.3.8) by  $(q^{-n})_k A_k / (q)_k$ ,  $0 \leq k \leq n$ , where there are no restrictions on  $A_k$ ,  $0 \leq k \leq n$ . Also multiply both sides by  $(bc/(aq))^k$ .

Now sum both sides from  $k = 0$  to  $k = n$  to arrive at

$$\sum_{k=0}^n \frac{(b)_k (c)_k (q^{-n})_k A_k}{(aq/b)_k (aq/c)_k (q)_k} = \sum_{k=0}^n \sum_{j=0}^k \frac{(aq/(bc))_j (aq^k)_j (q^{-k})_j (q^{-n})_k}{(aq/b)_j (aq/c)_j (q)_j (q)_k} q^j \left(\frac{bc}{aq}\right)^k A_k. \quad (3.3.9)$$

Invert the order of summation in (3.3.9), so that in the inner sum  $j \leq k \leq n$ , and in the outer sum,  $0 \leq j \leq n$ . Next, let  $k = i + j$ . Thus, using the more compact notation (1.1.2), we can rewrite (3.3.9) in the form

$$\begin{aligned} & \sum_{k=0}^n \frac{(b)_k (c)_k (q^{-n})_k A_k}{(aq/b)_k (aq/c)_k (q)_k} \\ &= \sum_{j=0}^n \sum_{i=0}^{n-j} \frac{(aq/(bc), aq^{i+j}, q^{-i-j}; q)_j (q^{-n}; q)_{i+j}}{(aq/b, aq/c, q; q)_j (q; q)_{i+j}} q^j \left(\frac{bc}{aq}\right)^{i+j} A_{i+j}. \end{aligned} \quad (3.3.10)$$

To simplify (3.3.10), we employ the elementary identities

$$(q^{-i-j}; q)_j = (-1)^j (q^{i+1})_j q^{-ij-j(j+1)/2}, \quad (3.3.11)$$

$$\frac{(q^{-i-j}; q)_j}{(q)_{i+j}} = \frac{(-1)^j q^{-ij-j(j+1)/2}}{(q)_i}, \quad (3.3.12)$$

$$(aq^{i+j}; q)_j = \frac{(aq^j; q)_j (aq^{2j}; q)_i}{(aq^j; q)_i}, \quad (3.3.13)$$

$$(q^{-n}; q)_{i+j} = (q^{-n}; q)_j (q^{j-n}; q)_i. \quad (3.3.14)$$

Using (3.3.11)–(3.3.14) in (3.3.10), we deduce the following theorem, which can be regarded as a version of *Bailey's Lemma*.

**Theorem 3.3.2.** *We have*

$$\begin{aligned} \sum_{k=0}^n \frac{(b, c, q^{-n}; q)_k}{(aq/b, aq/c, q; q)_k} A_k &= \sum_{j=0}^n \frac{(aq/(bc), aq^j, q^{-n}; q)_j}{(aq/b, aq/c, q; q)_j} (-1)^j q^{-j(j-1)/2} \\ &\quad \times \sum_{i=0}^{n-j} \frac{(aq^{2j}, q^{j-n}; q)_i}{(aq^j, q; q)_i} q^{-ij} \left(\frac{bc}{aq}\right)^{i+j} A_{i+j}. \end{aligned} \quad (3.3.15)$$

If we set

$$A_k = \frac{(a, a_1, \dots, a_r; q)_k}{(b_1, b_2, \dots, b_{r+1}; q)_k} z^k$$

and note that

$$\frac{(aq^j)_j (a)_{i+j}}{(aq^j)_i} = (a)_{2j},$$

then Theorem 3.3.2 takes the following shape.

**Theorem 3.3.3.** *For any nonnegative integer  $r$ ,*

$$\begin{aligned} & {}_{r+4}\phi_{r+3} \left[ \begin{matrix} a, b, c, a_1, \dots, a_r, q^{-n} \\ aq/b, aq/c, b_1, \dots, b_{r+1} \end{matrix}; q; z \right] \\ &= \sum_{j=0}^n \frac{(aq/(bc), a_1, \dots, a_r, q^{-n}; q)_j}{(aq/b, aq/c, q, b_1, \dots, b_{r+1}; q)_j} (a; q)_{2j} \left( -\frac{bcz}{aq} \right)^j q^{-j(j-1)/2} \\ & \quad \times {}_{r+2}\phi_{r+1} \left[ \begin{matrix} aq^{2j}, a_1q^j, \dots, a_rq^j, q^{j-n} \\ b_1q^j, \dots, b_{r+1}q^j \end{matrix}; q; \frac{bcz}{aq^{j+1}} \right]. \end{aligned} \quad (3.3.16)$$

The thrust of Theorem 3.3.3 is that we can reduce the evaluation of a  ${}_{r+4}\phi_{r+3}$  to the evaluation of a  ${}_{r+2}\phi_{r+1}$ . If this is not possible, then we reiterate (3.3.16) until we find that, for a “small” value of  $r$ , we can evaluate the series using one of our standard theorems.

**Definition 3.3.4.** *In the notation (3.3.1), we say that  ${}_{r+1}\phi_r$  is well-poised if*

$$qa_1 = a_2b_1 = a_3b_2 = \dots = a_{r+1}b_r. \quad (3.3.17)$$

**Definition 3.3.5.** *In the notation (3.3.1), we say that  ${}_{r+1}\phi_r$  is very-well-poised if, in addition to (3.3.17),*

$$a_2 = q\sqrt{a_1}, \quad a_3 = -q\sqrt{a_1}. \quad (3.3.18)$$

In the next theorem, we evaluate in closed form certain very-well-poised  ${}_4\phi_3$ -functions.

**Theorem 3.3.6.** *If  $\delta_{n,0}$  denotes the “Kronecker delta” and  $n$  is a nonnegative integer, then*

$${}_4\phi_3 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq^{n+1} \end{matrix}; q; q^n \right] = \delta_{n,0}. \quad (3.3.19)$$

**Proof.** In Theorem 3.3.3, set

$$b = q\sqrt{a}, c = -q\sqrt{a}, a_k = b_k, 1 \leq k \leq r, b_{r+1} = aq^{n+1}.$$

Hence,

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq^{n+1} \end{matrix}; q; z \right] \\ &= \sum_{j=0}^n \frac{(-q^{-1}, q^{-n}; q)_j (a; q)_{2j}}{(q, \sqrt{a}, -\sqrt{a}, aq^{n+1}; q)_j} (qz)^j q^{-j(j-1)/2} {}_2\phi_1 \left[ \begin{matrix} aq^{2j}, q^{j-n} \\ aq^{j+n+1} \end{matrix}; q; -zq^{1-j} \right]. \end{aligned} \quad (3.3.20)$$

Next, let  $z = q^n$  and apply Bailey’s Theorem 3.2.4 to deduce that

$${}_2\phi_1 \left[ \begin{matrix} aq^{2j}, q^{j-n} \\ aq^{j+n+1} \end{matrix}; q; -q^{1+n-j} \right] = \frac{(-q; q)_\infty (aq^{2j+1}, aq^{2n+2}; q^2)_\infty}{(aq^{1+n+j}, -q^{1+n-j}; q)_\infty}. \quad (3.3.21)$$

Let  $R$  denote the right-hand side of (3.3.20). Putting (3.3.21) in  $R$ , we find that

$$R = (-q; q)_\infty \sum_{j=0}^n \frac{(-q^{-1}, q^{-n}; q)_j (a; q)_{2j} (aq^{2j+1}, aq^{2n+2}; q^2)_\infty}{(q, \sqrt{a}, -\sqrt{a}, aq^{n+1}; q)_j (aq^{1+n+j}, -q^{1+n-j}; q)_\infty} q^{(n+1)j} q^{-j(j-1)/2}. \quad (3.3.22)$$

We now record the simplifications, which we leave as exercises,

$$\frac{(a; q)_{2j}}{(\sqrt{a}, -\sqrt{a}; q)_j} = (aq; q^2)_j, \quad (3.3.23)$$

$$(aq; q^2)_j (aq^{2j+1}; q^2)_\infty = (aq; q^2)_\infty, \quad (3.3.24)$$

$$(aq^{n+1}; q)_j (aq^{n+j+1}; q)_\infty = (aq^{n+1}; q)_\infty, \quad (3.3.25)$$

and

$$(-q^{1+n-j}; q)_\infty = (-q^{-n}; q)_j (-q^{n+1}; q)_\infty q^{nj-j(j-1)/2}. \quad (3.3.26)$$

Utilizing (3.3.23)–(3.3.26) in (3.3.22), we find that

$$\begin{aligned} R &= \frac{(-q; q)_\infty (aq; q^2)_\infty (aq^{2n+2}; q^2)_\infty}{(aq^{n+1}; q)_\infty (-q^{n+1}; q)_\infty} \sum_{j=0}^n \frac{(-q^{-1}, q^{-n}; q)_j}{(q, -q^{-n}; q)_j} q^j \\ &= \frac{(-q; q)_n (aq; q)_n}{(\sqrt{aq}; q)_n (-\sqrt{aq}; q)_n} {}_2\phi_1(-q^{-1}, q^{-n}; -q^{-n}; q; q). \end{aligned} \quad (3.3.27)$$

If  $n = 0$ , we easily see that  $R = 1$ . If  $n > 0$ , we use the  $q$ -analogue of the Chu–Vandermonde Theorem, Theorem 3.2.3, with  $b = -q^{-1}$  and  $c = -q^{-n}$  to conclude that

$${}_2\phi_1(-q^{-1}, q^{-n}; -q^{-n}; q; q) = \frac{(q^{-n+1}; q)_n}{(-q^{-n}; q)_n} = 0. \quad (3.3.28)$$

Recalling that  $R$  denotes the right-hand side of (3.3.20), that  $R$  is given by (3.3.27), and that  $R$  is evaluated by (3.3.28) for  $n > 0$  and in the line above for  $n = 0$ , we complete the proof of Theorem 3.3.6.  $\square$

In the next theorem, we sum a very-well-poised  ${}_6\phi_5$ .

**Theorem 3.3.7.** *For each nonnegative integer  $n$ ,*

$${}_6\phi_5 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq^{n+1}; q; \frac{aq^{n+1}}{bc} \end{matrix} \right] = \frac{(aq, aq/(bc); q)_n}{(aq/b, aq/c; q)_n}. \quad (3.3.29)$$

**Proof.** Let  $r = 2$  and

$$a_1 = q\sqrt{a}, a_2 = -q\sqrt{a}, b_1 = \sqrt{a}, b_2 = -\sqrt{a}, b_3 = aq^{n+1}$$

in Theorem 3.3.3 to find that

$$\begin{aligned} &{}_6\phi_5 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq^{n+1}; q; z \end{matrix} \right] \\ &= \sum_{j=0}^n \frac{(aq/(bc), q\sqrt{a}, -q\sqrt{a}, q^{-n}; q)_j (a; q)_{2j}}{(q, \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq^{n+1}; q)_j} \left( -\frac{bcz}{aq} \right)^j q^{-j(j-1)/2} \\ &\quad \times {}_4\phi_3 \left[ \begin{matrix} aq^{2j}, q^{j+1}\sqrt{a}, -q^{j+1}\sqrt{a}, q^{j-n} \\ q^j\sqrt{a}, -q^j\sqrt{a}, aq^{j+n+1}; q; \frac{bcz}{aq^{j+1}} \end{matrix} \right]. \end{aligned} \quad (3.3.30)$$

Now let  $z = aq^{n+1}/(bc)$  in (3.3.30) and apply Theorem 3.3.6 with  $a$  replaced by  $aq^{2j}$  and  $n$  replaced by  $n-j$ . Thus, in the sum on  $j$  in (3.3.30), all of the terms equal 0 except the term with  $j = n$ . Hence,

$$\begin{aligned} & {}_6\phi_5 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq^{n+1}; q; z \end{matrix} \right] \\ &= \frac{(aq/(bc), q\sqrt{a}, -q\sqrt{a}, q^{-n}; q)_n (a; q)_{2n}}{(q, \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq^{n+1}; q)_n} (-1)^n q^{n^2} q^{-n(n-1)/2}. \end{aligned} \quad (3.3.31)$$

Since

$$\frac{(q\sqrt{a}, -q\sqrt{a}; q)_n}{(\sqrt{a}, -\sqrt{a}; q)_n} = \frac{1 - aq^{2n}}{1 - a}, \quad (3.3.32)$$

$$\frac{(a; q)_{2n}(1 - aq^{2n})}{(1 - a)(q, aq^{n+1}; q)_n} = \frac{(aq; q)_n}{(q; q)_n},$$

and

$$(q^{-n}; q)_n = (-1)^n (q; q)_n q^{-n(n+1)/2},$$

we find from (3.3.31) that

$${}_6\phi_5 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq^{n+1}; q; z \end{matrix} \right] = \frac{(aq, aq/(bc); q)_n}{(aq/b, aq/c; q)_n},$$

which completes the proof of Theorem 3.3.7.  $\square$

The next theorem is Watson's transformation formula for a very-well-poised terminating  ${}_8\phi_7$ . It is one of the most useful theorems in  $q$ -series, especially in the theory of partitions.

**Theorem 3.3.8** ( $q$ -analogue of Whipple's Theorem). *For each nonnegative integer  $n$ ,*

$$\begin{aligned} & {}_8\phi_7 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, d, e, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq/d, aq/e, aq^{n+1}; q; \frac{a^2q^{n+2}}{bcde} \end{matrix} \right] \\ &= \frac{(aq, aq/(de); q)_n}{(aq/d, aq/e; q)_n} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, d, e, aq/(bc) \\ aq/b, aq/c, deq^{-n}/a; q; q \end{matrix} \right]. \end{aligned} \quad (3.3.33)$$

**Proof.** With an appeal to Theorem 3.3.3, we set

$$\begin{aligned} a_1 &= q\sqrt{a}, a_2 = -q\sqrt{a}, a_3 = d, a_4 = e, \\ b_1 &= \sqrt{a}, b_2 = -\sqrt{a}, b_3 = aq/d, b_4 = aq/e, b_5 = aq^{n+1}. \end{aligned}$$

Also, set

$$z = \frac{a^2q^{2+n}}{bcde}.$$

Hence,

$$\begin{aligned}
& {}_8\phi_7 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, d, e, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq/d, aq/e, aq^{n+1}; q; \frac{a^2q^{n+2}}{bcde} \end{matrix} \right] \\
&= \sum_{j=0}^n \frac{(aq/(bc), q\sqrt{a}, -q\sqrt{a}, d, e, q^{-n}; q)_j (a; q)_{2j}}{(q, \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq/d, aq/e, aq^{n+1}; q)_j} \left( -\frac{aq^{n+1}}{de} \right)^j q^{-j(j-1)/2} \\
&\quad \times {}_6\phi_5 \left[ \begin{matrix} aq^{2j}, q^{j+1}\sqrt{a}, -q^{j+1}\sqrt{a}, dq^j, eq^j, q^{j-n} \\ q^j\sqrt{a}, -q^j\sqrt{a}, aq^{j+1}/d, aq^{j+1}/e, aq^{j+n+1}; q; \frac{aq^{n+1-j}}{de} \end{matrix} \right].
\end{aligned} \tag{3.3.34}$$

We now use Theorem 3.3.7 with  $a$  replaced by  $aq^{2j}$ ,  $b = dq^j$ ,  $c = eq^j$ , and  $n$  replaced by  $n - j$  to evaluate the  ${}_6\phi_5$  above. Consequently,

$$\begin{aligned}
& {}_6\phi_5 \left[ \begin{matrix} aq^{2j}, q^{j+1}\sqrt{a}, -q^{j+1}\sqrt{a}, dq^j, eq^j, q^{j-n} \\ q^j\sqrt{a}, -q^j\sqrt{a}, aq^{j+1}/d, aq^{j+1}/e, aq^{j+n+1}; q; \frac{aq^{n+1-j}}{de} \end{matrix} \right] \\
&= \frac{\left( aq^{2j+1}, \frac{aq^{2j+1}}{deq^{2j}} \right)_{n-j}}{\left( \frac{aq^{2j+1}}{dq^j}, \frac{aq^{2j+1}}{eq^j} \right)_{n-j}} = \frac{\left( aq^{2j+1}, \frac{aq}{de} \right)_{n-j}}{\left( \frac{aq^{j+1}}{d}, \frac{aq^{j+1}}{e} \right)_{n-j}} \\
&= \frac{(aq^{2j+1})_n (aq/(de))_n (dq^{-n-j}/a)_j (eq^{-n-j}/a)_j}{(q^{-n-2j}/a)_j (q^{-n}de/a)_j (aq^{j+1}/d)_n (aq^{j+1}/e)_n}, \tag{3.3.35}
\end{aligned}$$

where we made four applications of (3.3.6). First, using (3.3.32), we readily find that

$$\frac{(q\sqrt{a}, -q\sqrt{a}; q)_j}{(\sqrt{a}, -\sqrt{a}; q)_j} (a; q)_{2j} (aq^{2j+1}; q)_n = (aq; q)_{n+2j}. \tag{3.3.36}$$

Second, a brief calculation shows that

$$(q^{-n-2j}/a; q)_j = (aq^{n+j+1}; q)_j (-1)^j q^{-nj-j(3j+1)/2} a^{-j}. \tag{3.3.37}$$

Third, we readily see that

$$\frac{(aq; q)_{n+2j}}{(aq^{n+j+1}; q)_j (aq^{n+1}; q)_j} = (aq; q)_n. \tag{3.3.38}$$

Fourth,

$$\frac{(dq^{-n-j}/a; q)_j}{(aq/d; q)_j (aq^{j+1}/d; q)_n} = \frac{(-1)^j d^j a^{-j} q^{-nj-j(j+1)/2}}{(aq/d; q)_n}. \tag{3.3.39}$$

Similarly,

$$\frac{(eq^{-n-j}/a; q)_j}{(aq/e; q)_j (aq^{j+1}/e; q)_n} = \frac{(-1)^j e^j a^{-j} q^{-nj-j(j+1)/2}}{(aq/e; q)_n}. \tag{3.3.40}$$

Hence, using (3.3.35)–(3.3.40) in (3.3.34), after an enormous amount of simplification, we finally find that

$$\begin{aligned} {}_8\phi_7 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, d, e, q^{-n} \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, aq/d, aq/e, aq^{n+1}; q; \frac{a^2 q^{n+2}}{bcde} \end{matrix} \right] \\ = \frac{(aq; q)_n (aq/(de); q)_n}{(aq/d; q)_n (aq/e; q)_n} \sum_{j=0}^n \frac{(aq/bc, d, e, q^{-n}; q)_j}{(q, aq/b, aq/c, q^{-n} de/a; q)_j} q^j. \end{aligned}$$

Thus, the proof of Theorem 3.3.8 is complete.  $\square$

**Definition 3.3.9.** *If  $qa_1a_2 \cdots a_{r+1} = b_1b_2 \cdots b_r$ , then we say that  ${}_{r+1}\phi_r$  is balanced or Saalschützian.*

Note that the  ${}_4\phi_3$  in Watson's transformation is balanced.

Watson's  ${}_8\phi_7$  transformation is not found in any of Ramanujan's published papers, his earlier notebooks, or his lost notebook. However, the following corollary is Entry 7 in Chapter 16 of his second notebook [91], [27, p. 16].

**Corollary 3.3.10.** *We have*

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(a)_k (d/b)_k (d/c)_k (d/q)_k (1-dq^{2k-1})}{(b)_k (c)_k (d/a)_k (q)_k (1-d/q)} \left( \frac{bc}{a} \right)^k q^{k(k-1)} \\ = \frac{(a)_{\infty} (d)_{\infty}}{(b)_{\infty} (c)_{\infty}} \sum_{k=0}^{\infty} \frac{(b/a)_k (c/a)_k}{(d/a)_k (q)_k} a^k. \quad (3.3.41) \end{aligned}$$

**Proof.** We let  $n \rightarrow \infty$  in Theorem 3.3.8. On the left-hand side of (3.3.33), we observe that

$$\lim_{n \rightarrow \infty} \frac{(q^{-n})_k q^{nk}}{(aq^{n+1})_k} = (-1)^k q^{k(k-1)/2},$$

while on the right-hand side of (3.3.33), we see that

$$\lim_{n \rightarrow \infty} \frac{(q^{-n})_k}{(deq^{-n}/a)_k} = \left( \frac{a}{de} \right)^k.$$

Thus, so far, we have shown from (3.3.33) that

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(a)_k (b)_k (c)_k (d)_k (e)_k (1-aq^{2k})}{(aq/b)_k (aq/c)_k (aq/d)_k (aq/e)_k (1-a)(q)_k} \left( -\frac{a^2}{bcde} \right)^k q^{k(k+3)/2} \\ = \frac{(aq, aq/(de))_{\infty}}{(aq/d, aq/e)_{\infty}} {}_3\phi_2 \left[ \begin{matrix} aq/(bc), d, e, aq \\ aq/b, aq/c; de \end{matrix} \right]. \quad (3.3.42) \end{aligned}$$

Next, we let  $c \rightarrow \infty$ . Observing that

$$\lim_{c \rightarrow \infty} \frac{(c)_k c^{-k}}{(aq/c)_k} = (-1)^k q^{k(k-1)/2},$$

we see that (3.3.42) reduces to

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(a)_k (b)_k (d)_k (e)_k (1 - aq^{2k})}{(aq/b)_k (aq/d)_k (aq/e)_k (1 - a)(q)_k} \left( \frac{a^2}{bde} \right)^k q^{k(k+1)} \\ = \frac{(aq, aq/(de))_{\infty}}{(aq/d, aq/e)_{\infty}} \sum_{k=0}^{\infty} \frac{(d)_k (e)_k}{(aq/b)_k (q)_k} \left( \frac{aq}{de} \right)^k. \end{aligned} \quad (3.3.43)$$

Now replace  $a$ ,  $b$ ,  $d$ , and  $e$  by  $d/q$ ,  $a$ ,  $d/b$ , and  $d/c$ , respectively. Hence, (3.3.43) and the  $q$ -analogue of Euler's transformation, Theorem 3.2.6, yield

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{(a)_k (d/b)_k (d/c)_k (d/q)_k (1 - dq^{2k-1})}{(b)_k (c)_k (d/a)_k (q)_k (1 - d/q)} \left( \frac{bc}{a} \right)^k q^{k(k-1)} \\ = \frac{(d)_{\infty} (bc/d)_{\infty}}{(b)_{\infty} (c)_{\infty}} \sum_{k=0}^{\infty} \frac{(d/b)_k (d/c)_k}{(d/a)_k (q)_k} \left( \frac{bc}{d} \right)^k \\ = \frac{(d)_{\infty} (bc/d)_{\infty}}{(b)_{\infty} (c)_{\infty}} \frac{(a)_{\infty}}{(bc/d)_{\infty}} \sum_{k=0}^{\infty} \frac{(b/a)_k (c/a)_k}{(d/a)_k (q)_k} a^k. \end{aligned} \quad (3.3.44)$$

After minor simplification, we complete the proof.  $\square$

### 3.4. Exercises

1. Prove that  $q$ -binomial coefficients are polynomials in  $q$ .
2. For each positive integer  $N$ , establish a result to H.A. Rothe, namely,

$$\sum_{k=0}^N \begin{bmatrix} N \\ k \end{bmatrix}_q (-1)^k q^{k(k-1)/2} x^k = (x; q)_N.$$

3. For each positive integer  $N$  and  $|x| < 1$ , prove that

$$\sum_{k=0}^{\infty} \begin{bmatrix} N+k-1 \\ k \end{bmatrix}_q x^k = \frac{1}{(x; q)_N}.$$

4. Verify (3.3.6).
5. Verify (3.3.11)–(3.3.14).
6. Verify (3.3.23)–(3.3.24).
7. Reverse the order of summation in the  $q$ -analogue of the Chu–Vandermonde Theorem, Theorem 3.2.3, to show that

$${}_2\phi_1(q^{-n}, b; c; q; q) = \frac{(c/b; q)_n}{(c; q)_n} b^n. \quad (3.4.1)$$



---

## Chapter 4

# Partition Identities Arising from $q$ -Series Identities

### 4.1. Theorems Arising from Basic $q$ -Series Theorems

We begin with a famous identity due to Lebesgue.

**Theorem 4.1.1.** For  $b \in \mathbb{C}$ ,

$$\sum_{n=0}^{\infty} \frac{(-bq; q)_n}{(q; q)_n} q^{n(n+1)/2} = \frac{(-bq^2; q^2)_{\infty}}{(q; q^2)_{\infty}}. \quad (4.1.1)$$

**First Proof of Theorem 4.1.1.** Employing Theorem 3.2.8 with  $a = 1$  and  $z = -bq$ , Corollary 3.1.3, and Euler's identity,

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(-bq; q)_n}{(q; q)_n} q^{n(n+1)/2} &= (-bq; q)_{\infty} (-q; q)_{\infty} \sum_{n=0}^{\infty} \frac{(-bq)^n}{(q; q)_n (-q; q)_n} \\ &= (-bq; q)_{\infty} (-q; q)_{\infty} \sum_{n=0}^{\infty} \frac{(-bq)^n}{(q^2; q^2)_n} \\ &= (-bq; q)_{\infty} (-q; q)_{\infty} \frac{1}{(-bq; q^2)_{\infty}} \\ &= \frac{(-bq^2; q^2)_{\infty}}{(q; q^2)_{\infty}}. \end{aligned}$$

□

**Second Proof of Theorem 4.1.1.** Replacing  $b$  by  $1/b$  in Bailey's Theorem 3.2.4, we find that

$${}_2\phi(a, 1/b; qab; -qb) = \frac{(aq; q^2)_\infty (-q; q)_\infty (q^2 ab^2; q^2)_\infty}{(qab; q)_\infty (-qb; q)_\infty}. \quad (4.1.2)$$

Now,

$$\lim_{b \rightarrow 0} (1/b; q)_n (-qb)^n = q^{n(n+1)/2}. \quad (4.1.3)$$

Letting  $b$  tend to 0 on both sides of (4.1.2) and using (4.1.3) and Euler's theorem, we find that

$$\sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} q^{n(n+1)/2} = \frac{(aq; q^2)_\infty}{(q; q^2)_\infty}.$$

Letting  $a = -bq$ , we complete the proof.  $\square$

The following beautiful result is due to N. J. Fine [54]; its proof can also be found in Andrews' text [11, pp. 26–27]. Observe that Theorem 4.1.2 is a refinement of Euler's theorem, Theorem 1.2.3.

**Theorem 4.1.2.** *The number of partitions of a positive integer  $n$  into distinct parts with largest part  $k$  equals the number of partitions of  $n$  into odd parts, such that  $2k + 1$  equals the largest part plus twice the number of parts.*

We interpret Theorem 4.1.2 as an equivalent  $q$ -series identity.

**Theorem 4.1.3.** *We have*

$$\sum_{j=0}^{\infty} (-q; q)_j (tq)^{j+1} = \sum_{j=0}^{\infty} \frac{t^{j+1} q^{2j+1}}{(tq; q^2)_{j+1}}. \quad (4.1.4)$$

**Proof.** Write

$$\sum_{j=0}^{\infty} (-q; q)_j (tq)^{j+1} = \sum_{j=0}^{\infty} (1+q)(1+q^2) \cdots (1+q^j) q^{j+1} t^{j+1}.$$

Let  $k = j + 1$ . We readily see that the series above generates partitions of  $n$  into distinct parts, with largest part  $k$ .

On the other hand,

$$\sum_{j=0}^{\infty} \frac{t^{j+1} q^{2j+1}}{(tq; q^2)_{j+1}} = \sum_{j=0}^{\infty} t^{j+1} q^{2j+1} \sum_{n_1=0}^{\infty} (tq)^{n_1} \sum_{n_2=0}^{\infty} (tq^3)^{n_2} \cdots \sum_{n_{j+1}=0}^{\infty} (tq^{2j+1})^{n_{j+1}}. \quad (4.1.5)$$

For a partition  $\pi$ , let  $\nu(\pi)$  denote the number of parts of  $\pi$ . Note that  $1 + n_1 + n_2 + \cdots + n_{j+1} = \nu(\pi)$ . The power of  $t$  in (4.1.5) is

$$k = j + 1 + n_1 + n_2 + \cdots + n_{j+1} = j + 1 + \nu(\pi) - 1 = j + \nu(\pi),$$

Hence,

$$2k + 1 = 2j + 2\nu(\pi) + 1 = \text{largest part} + 2\nu(\pi).$$

Thus, we have shown that Theorem 4.1.2 and Theorem 4.1.3 are equivalent.  $\square$

**Example 4.1.4.** Let  $A_k(n)$  and  $B_k(n)$  denote the coefficients of  $t^k q^n$  on the left and right sides of (4.1.4), respectively. Let  $k = 8$  and  $n = 14$ . We see that  $A_8(14) = 4 = B_8(14)$ , with the respective representations being

$$\begin{aligned} &8 + 6, 8 + 5 + 1, 8 + 4 + 2, 8 + 3 + 2 + 1; \\ &13 + 1(17 = 13 + 2 \cdot 2), 9 + 3 + 1 + 1(17 = 9 + 2 \cdot 4), \\ &5 + 5 + 1 + 1 + 1 + 1(17 = 5 + 2 \cdot 6), 5 + 3 + 3 + 1 + 1 + 1(17 = 5 + 2 \cdot 6). \end{aligned}$$

**Proof of Theorem 4.1.3.** Using Corollary 3.1.3 twice below, we find that

$$\begin{aligned} \sum_{j=0}^{\infty} (-q; q)_j (tq)^{j+1} &= \sum_{j=0}^{\infty} \frac{(q^2; q^2)_j (tq)^{j+1}}{(q; q)_j} \\ &= tq(q^2; q^2)_{\infty} \sum_{j=0}^{\infty} \frac{(tq)^j}{(q; q)_j} \frac{1}{(q^{2j+2}; q^2)_{\infty}} \\ &= tq(q^2; q^2)_{\infty} \sum_{j=0}^{\infty} \frac{(tq)^j}{(q; q)_j} \sum_{m=0}^{\infty} \frac{q^{(2j+2)m}}{(q^2; q^2)_m} \\ &= tq(q^2; q^2)_{\infty} \sum_{m=0}^{\infty} \frac{q^{2m}}{(q^2; q^2)_m} \sum_{j=0}^{\infty} \frac{t^j q^{j(2m+1)}}{(q; q)_j} \\ &= tq(q^2; q^2)_{\infty} \sum_{m=0}^{\infty} \frac{q^{2m}}{(q^2; q^2)_m (tq^{2m+1}; q)_{\infty}} \\ &= \frac{tq(q^2; q^2)_{\infty}}{(tq; q)_{\infty}} \sum_{m=0}^{\infty} \frac{(tq; q^2)_m (tq^2; q^2)_m q^{2m}}{(q^2; q^2)_m}, \end{aligned} \quad (4.1.6)$$

where we have used the fact that  $(tq; q)_{2m}(tq^{2m+1}; q)_{\infty} = (tq; q)_{\infty}$ . Applying Heine's transformation, Theorem 3.2.1, to the far right side of (4.1.6), with first  $q$  replaced by  $q^2$ , and then with  $a = tq$ ,  $b = tq^2$ ,  $c = 0$ , and  $z = q^2$ , we deduce that

$$\begin{aligned} &\sum_{j=0}^{\infty} (-q; q)_j (tq)^{j+1} \\ &= \frac{tq(q^2; q^2)_{\infty}}{(tq; q)_{\infty}} \frac{(tq^2; q^2)_{\infty}}{(q^2; q^2)_{\infty}} \frac{(tq^3; q^2)_{\infty}}{(q^2; q^2)_{\infty}} \sum_{m=0}^{\infty} \frac{(q^2; q^2)_m t^m q^{2m}}{(tq^3; q^2)_m (q^2; q^2)_m} \\ &= \frac{tq(tq^2; q^2)_{\infty} (tq^3; q^2)_{\infty}}{(tq; q)_{\infty}} \sum_{m=0}^{\infty} \frac{t^m q^{2m}}{(tq^3; q^2)_m} \\ &= \frac{tq}{1-tq} \sum_{m=0}^{\infty} \frac{t^m q^{2m}}{(tq^3; q^2)_m} \\ &= \sum_{m=0}^{\infty} \frac{t^{m+1} q^{2m+1}}{(tq; q^2)_{m+1}}, \end{aligned}$$

and this completes the proof.  $\square$

**Definition 4.1.5.** Let  $P(r, m, n)$  denote the number of partitions of  $n$  into exactly  $m$  parts with the largest part  $\leq r$ .


Note that earlier we had defined  $p(r, m, n)$  to be the number of partitions of  $n$  into exactly  $m$  parts with the largest part equal to  $r$ .

**Theorem 4.1.6.** The number of partitions of  $a - c$  into exactly  $b - 1$  parts, none exceeding  $c$  equals the number of partitions of  $a - b$  into exactly  $c - 1$  parts, none exceeding  $b$ , or

$$P(c, b - 1, a - c) = P(b, c - 1, a - b).$$

We provide two proofs. The first is combinatorial; the second is analytic. We shall accompany our proof by an example.

**First Proof of Theorem 4.1.6.** Consider the Ferrers graph of a partition of the first type. For example, let  $a = 21$ ,  $c = 7$ , and  $b = 5$ . Thus, below we have a Ferrers graph of a partition of  $a - c = 14$  into exactly  $5 - 1 = 4$  parts, with none exceeding the largest part  $6 < 7$ .



**Figure 1.** A Partition of  $a - c$

Now add a row of  $c$  nodes to the top of the Ferrers graph above. So now we have a partition of  $a - c + c$ , in the Example 21.



**Figure 2.** A Partition of  $a - c + c$

We now delete the first column. Thus, we have a Ferrers graph of  $a - b$ . The top row has  $c - 1$  nodes. And, there are less than or equal to  $b$  rows, in this example,  $\leq 5$  rows.

Now take the conjugate of the partition above. Thus, we have a partition of  $a - b$  into exactly  $c - 1$  parts, with the biggest part less than or equal to  $b$ .

All of these processes are reversible, and so we have shown simply with the use of Ferrers graphs that

$$P(c, b - 1, a - c) = P(b, c - 1, a - b).$$

□




Figure 3. A Partition of  $a - b$




Figure 4. Conjugate Partition of  $a - b$

**Second Proof of Theorem 4.1.6.** While our first proof of Theorem 4.1.6 was purely combinatorial, our second proof depends on the theory of  $q$ -series. We form a generating function

$$S(a, b, c) := \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{c=0}^{\infty} P(a, b - 1, a - c) x^b y^c q^a. \tag{4.1.7}$$

We can write

$$S(a, b, c) = 1 + \sum_{n=1}^{\infty} x(yq)^n \prod_{j=1}^n (1 + xq^j + x^2q^{2j} + x^3q^{3j} + \dots). \tag{4.1.8}$$

To see that (4.1.8) holds, first note that we have extracted  $x$ , because the exact number of parts is  $b - 1$ . Second, observe that  $y^c q^a = (yq)^c q^{a-c}$ , and so the power  $c$  of  $yq$  is the upper bound for parts. The running index  $n = c$  then stands for the largest possible part in the partitions being generated. The product in (4.1.8) generates the remaining parts. Note that indeed all parts are less than or equal to  $n$ . Because we had previously extracted  $b$ , we acquire an  $x$  a total of  $b - 1$  times, as required. Lastly, note that we are generating partitions of  $a - c$  in the product. Summing each of the  $n$  geometric series on the right side of (4.1.8), we find that

$$S(a, b, c) = 1 + \sum_{n=1}^{\infty} \frac{x(yq)^n}{(xq; q)_n} =: f(x, y). \tag{4.1.9}$$

We see that the generating function for  $P(b, c - 1, a - b)$  is identical to that in (4.1.7), except that the roles of  $b$  and  $c$  are switched. Thus, to complete the proof, it suffices to show that

$$f(x, y) = f(y, x). \tag{4.1.10}$$

After a reformulation of the definition of  $f(x, y)$  in (4.1.9), we apply Heine's transformation (3.2.1) with  $a = 0$ ,  $b = q$ ,  $c = xq$ , and  $z = yq$ . Hence,

$$\begin{aligned} f(x, y) - 1 + x &= \sum_{n=0}^{\infty} \frac{x(yq)^n}{(xq; q)_n} = x \sum_{n=0}^{\infty} \frac{(0)_n(q)_n}{(xq)_n(q)_n} (yq)^n \\ &= x \frac{(q)_{\infty}}{(xq)_{\infty}(yq)_{\infty}} \sum_{n=0}^{\infty} \frac{(x)_n(yq)_n}{(0)_n(q)_n} q^n. \end{aligned} \quad (4.1.11)$$

Apply Heine's transformation (3.2.1) once again, now with  $a = yq$ ,  $b = x$ ,  $c = 0$ , and  $z = q$ . Hence, from (4.1.11),

$$\begin{aligned} f(x, y) - 1 + x &= x \frac{(q)_{\infty}}{(xq)_{\infty}(yq)_{\infty}} \frac{(yq^2)_{\infty}(x)_{\infty}}{(q)_{\infty}} \sum_{n=0}^{\infty} \frac{(0)_n(q)_n}{(q)_n(yq^2)_n} x^n \\ &= x(1-x) \sum_{n=0}^{\infty} \frac{x^n}{(yq)_{n+1}} \\ &= (1-x) \sum_{n=1}^{\infty} \frac{x^n}{(yq)_n}, \end{aligned} \quad (4.1.12)$$

where we replaced  $n$  by  $n-1$ . Therefore, rearranging (4.1.12), we deduce that

$$\begin{aligned} f(x, y) &= (1-x) \sum_{n=0}^{\infty} \frac{x^n}{(yq)_n} \\ &= \sum_{n=0}^{\infty} \frac{x^n}{(yq)_n} - \sum_{n=0}^{\infty} \frac{x^{n+1}}{(yq)_n} \\ &= 1 + \sum_{n=1}^{\infty} \frac{x^n}{(yq)_n} - \sum_{n=1}^{\infty} \frac{x^n(1-yq^n)}{(yq)_n} \\ &= 1 + \sum_{n=1}^{\infty} \frac{y(xq)^n}{(yq)_n} \\ &= f(y, x). \end{aligned} \quad (4.1.13)$$

Our last identity (4.1.13) is precisely (4.1.10), and so the proof is complete.  $\square$

We come now to one of the highlights of these lecture notes, our first proof of the epic Rogers–Ramanujan identities. All of the groundwork has been laid in Chapter 3. We first derive a corollary of Corollary 3.3.10.

**Corollary 4.1.7.** *We have*

$$1 + \sum_{k=1}^{\infty} \frac{(-1)^k(1-aq^{2k})(aq)_{k-1}a^{2k}q^{k(5k-1)/2}}{(q)_k} = (aq)_{\infty} \sum_{k=0}^{\infty} \frac{a^k q^{k^2}}{(q)_k}. \quad (4.1.14)$$

**Proof.** Return to (3.3.43) and let  $b, d, e \rightarrow \infty$ . If we use

$$\lim_{\alpha \rightarrow \infty} \frac{(\alpha)_k}{\alpha^k} = (-1)^k q^{k(k-1)/2}$$

on both the left and right sides of (3.3.43), we easily deduce the truth of (4.1.14) to complete the proof.  $\square$

**Theorem 4.1.8** (Rogers–Ramanujan Identities). *We have*

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q)_n} = \frac{1}{(q; q^5)_{\infty} (q^4; q^5)_{\infty}}, \quad (4.1.15)$$

$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_n} = \frac{1}{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty}}. \quad (4.1.16)$$

Before proving (4.1.15) and (4.1.16), we provide a brief history and examples. Ramanujan discovered these identities while in England, but at first he did not have proofs. One day, he was perusing old issues of the *Proceedings of the London Mathematical Society* and came across a paper [94] by L. J. Rogers from 1894 in which he found the identities that he recently had proven. It should be remarked that at that time Rogers was considered to be eccentric both personally and mathematically, and so his work was completely ignored. Shortly thereafter, Ramanujan found his own proofs of these identities.

Let us examine the partition-theoretic interpretation of the first identity (4.1.15). We note that  $1/(q)_n$  generates partitions into less than or equal to  $n$  parts. Take such a partition and arrange the terms in decreasing order. If the number of terms is less than  $n$ , then add 0's so that we have exactly  $n$  parts. Noting that  $n^2 = 1 + 3 + \cdots + (2n - 1)$ , take the previous partition and add to its parts in order,  $(2n - 1), (2n - 3), \dots, 3, 1$ . We thus have a partition in which the parts differ by at least 2. On the right-hand side of (4.1.15), we have the generating function for partitions into parts congruent to 1 and 4 modulo 5. In conclusion, we have the following partition-theoretic interpretation.

**Theorem 4.1.9.** *The number of partitions of an integer  $n$  into distinct parts such that the difference between any two parts is at least 2 is equinumerous with the number of partitions of  $n$  into parts congruent to either 1 or 4 modulo 5.*

We examine (4.1.16). As above,  $1/(q)_n$  generates partitions into at most  $n$  parts. Take such a partition and arrange the parts in decreasing order, where we add 0's if necessary to obtain a partition into exactly  $n$  parts. Now,  $n(n+1) = 2+4+\cdots+2n$ , and so we take the previous partition and add to its parts in decreasing order  $2n, 2n-2, \dots, 4, 2$ . We thus obtain a partition in which the parts differ by at least 2 and there are no 1's in the partition. Hence, we have the following theorem.

**Theorem 4.1.10.** *The number of partitions of an integer  $n$  into distinct parts such that the difference between any two parts is at least 2 and there are no 1's in the partitions is equinumerous with the number of partitions of  $n$  into parts congruent to either 2 or 3 modulo 5.*

**Example 4.1.11.** We give an example to illustrate each theorem. The partitions of 8 into distinct parts differing by at least two are the four partitions

$$8 = 7 + 1 = 6 + 2 = 5 + 3,$$

while the four partitions of 8 into parts congruent to 1 or 4 modulo 5 are

$$6 + 1 + 1 = 4 + 4 = 4 + 1 + 1 + 1 + 1 = 1 + 1 + \cdots + 1.$$

The three partitions of 8 into parts differing by at least 2 with no 1's allowed are

$$8 = 6 + 2 = 5 + 3,$$

while the three partitions of 8 into parts congruent to either 2 or 3 modulo 5 are

$$8 = 3 + 3 + 2 = 2 + 2 + 2 + 2.$$

**Proof.** In Corollary 4.1.7, set  $a = 1$ . Then

$$\begin{aligned} (q)_\infty \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q)_n} &= 1 + \sum_{n=1}^{\infty} \frac{(-1)^n (1 - q^{2n})(q)_{n-1} q^{n(5n-1)/2}}{(q)_n} \\ &= 1 + \sum_{n=1}^{\infty} (-1)^n (1 + q^n) q^{n(5n-1)/2} \\ &= \sum_{n=0}^{\infty} (-1)^n q^{n(5n-1)/2} + \sum_{n=-\infty}^{-1} (-1)^n q^{-n} q^{n(5n+1)/2} \\ &= \sum_{n=-\infty}^{\infty} (-1)^n q^{n(5n-1)/2} \\ &= f(-q^2, -q^3) \\ &= (q^2; q^5)_\infty (q^3; q^5)_\infty (q^5; q^5)_\infty, \end{aligned}$$

by the Jacobi triple product identity, (1.1.7) or (3.1.19). Dividing both sides above by  $(q)_\infty$ , and simplifying, we complete the proof of (4.1.15).

Next, in Corollary 4.1.7, set  $a = q$ . Then

$$\begin{aligned} (q^2)_\infty \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_n} &= 1 + \sum_{n=1}^{\infty} \frac{(-1)^n (1 - q^{2n+1})(q^2)_{n-1} q^{2n+n(5n-1)/2}}{(q)_n} \\ &= 1 + \sum_{n=1}^{\infty} \frac{(-1)^n (1 - q^{2n+1}) q^{n(5n+3)/2}}{1 - q}. \end{aligned}$$

Multiplying both sides above by  $(1 - q)$  and setting  $n = -m - 1$  in the second sum on the right side below, we arrive at

$$\begin{aligned} (q)_\infty \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_n} &= 1 - q + \sum_{n=1}^{\infty} (-1)^n q^{n(5n+3)/2} - \sum_{n=1}^{\infty} (-1)^n q^{n(5n+7)/2+1} \\ &= 1 - q + \sum_{n=1}^{\infty} (-1)^n q^{n(5n+3)/2} + \sum_{m=-\infty}^{-2} (-1)^m q^{(m+1)(5m-2)/2+1} \end{aligned}$$



$$\begin{aligned}
&= \sum_{n=-\infty}^{\infty} (-1)^n q^{n(5n+3)/2} \\
&= f(-q^4, -q) \\
&= (q; q^5)_{\infty} (q^4; q^5)_{\infty} (q^5; q^5)_{\infty},
\end{aligned}$$

by the Jacobi triple product identity (3.1.19). Dividing both sides above by  $(q)_{\infty}$ , we finish the proof of (4.1.16).  $\square$

We next derive another corollary of Watson's transformation.

**Corollary 4.1.12.** *We have*

$$1 + \sum_{k=1}^{\infty} \frac{(1 - aq^{2k})(d)_k (aq)_{k-1}}{(q)_k (aq/d)_k} \left(\frac{a^2}{d}\right)^k q^{2k^2} = \frac{(aq)_{\infty}}{(aq/d)_{\infty}} \sum_{k=0}^{\infty} \frac{(-1)^k (d)_k}{(q)_k} \left(\frac{a}{d}\right)^k q^{k(k+1)/2}. \quad (4.1.17)$$

**Proof.** Return to (3.3.42) and let  $b$  and  $e$  tend to infinity. Hence, we find that

$$1 + \sum_{k=1}^{\infty} \frac{(1 - aq^{2k})(d)_k (a)_k}{(1-a)(q)_k (aq/d)_k} \left(\frac{a^2}{d}\right)^k q^{2k^2} = \frac{(aq)_{\infty}}{(aq/d)_{\infty}} \sum_{k=0}^{\infty} \frac{(-1)^k (d)_k}{(q)_k} \left(\frac{a}{d}\right)^k q^{k(k+1)/2}. \quad (4.1.18)$$

Minor simplification completes the proof.  $\square$

If we replace  $q$  by  $q^2$  and then set  $d = -q$  in (4.1.17), we find that

$$1 + \sum_{k=1}^{\infty} \frac{(1 - aq^{4k})(-1)^k (-q; q^2)_k (aq^2; q^2)_{k-1}}{(q^2; q^2)_k (-aq; q^2)_k} a^{2k} q^{4k^2 - k} = \frac{(aq^2; q^2)_{\infty}}{(-aq; q^2)_{\infty}} \sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} a^k q^{k^2}. \quad (4.1.19)$$

The next two theorems are the analytic versions of the two famous Göllnitz–Gordon identities [61]. However, the identities can be found in Ramanujan's lost notebook [92, p. 41], [18, pp. 36–37]. They can also be found in Slater's list [98, equations (36), (34)], but with  $q$  replaced by  $-q$ . The Göllnitz–Gordon identities have played a seminal role in the subsequent development in the theory of partition identities. They were first studied in this regard by H. Göllnitz [60] and by B. Gordon [62], [63]. A generalization by Andrews [6] led to a number of further discoveries culminating in [8]. After we prove the two identities, we shall discuss their combinatorial implications and give an example.

**Theorem 4.1.13** (The Göllnitz–Gordon Identities). *We have*

$$\sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2} = \frac{1}{(q; q^8)_{\infty} (q^4; q^8)_{\infty} (q^7; q^8)_{\infty}}, \quad (4.1.20)$$

$$\sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2 + 2k} = \frac{1}{(q^3; q^8)_{\infty} (q^4; q^8)_{\infty} (q^5; q^8)_{\infty}}. \quad (4.1.21)$$

**Proof.** Letting  $a = 1$  in (4.1.19), we deduce that

$$\begin{aligned}
\frac{(q^2; q^2)_\infty}{(-q; q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2} &= 1 + \sum_{k=1}^{\infty} \frac{(1 - q^{4k})(-1)^k (-q; q^2)_k (q^2; q^2)_{k-1}}{(q^2; q^2)_k (-q; q^2)_k} q^{4k^2 - k} \\
&= 1 + \sum_{k=1}^{\infty} (-1)^k (1 + q^{2k}) q^{4k^2 - k} \\
&= \sum_{k=1}^{\infty} (-1)^k q^{4k^2 - k} + \sum_{k=0}^{\infty} (-1)^k q^{4k^2 + k} \\
&= \sum_{k=-1}^{-\infty} (-1)^k q^{4k^2 + k} + \sum_{k=0}^{\infty} (-1)^k q^{4k^2 + k} \\
&= \sum_{k=-\infty}^{\infty} (-1)^k q^{4k^2 + k} \\
&= f(-q^5, -q^3) \\
&= (q^3; q^8)_\infty (q^5; q^8)_\infty (q^8; q^8)_\infty, \tag{4.1.22}
\end{aligned}$$

by the Jacobi triple product identity (3.1.19). A slight rearrangement of (4.1.22) yields

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2} &= \frac{(-q; q^2)_\infty (q^3; q^8)_\infty (q^5; q^8)_\infty (q^8; q^8)_\infty}{(q^2; q^2)_\infty} \\
&= \frac{(q^2; q^4)_\infty (q^3; q^8)_\infty (q^5; q^8)_\infty (q^8; q^8)_\infty}{(q; q^2)_\infty (q^2; q^2)_\infty} \\
&= \frac{1}{(q; q^8)_\infty (q^4; q^8)_\infty (q^7; q^8)_\infty}.
\end{aligned}$$

Thus, the proof of (4.1.20) is complete.

To prove (4.1.21), first set  $a = q^2$  in (4.1.19) to find that

$$\begin{aligned}
\frac{(q^4; q^2)_\infty}{(-q^3; q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2 + 2k} &= 1 + \sum_{k=1}^{\infty} \frac{(-1)^k (1 - q^{4k+2})(-q; q^2)_k (q^4; q^2)_{k-1}}{(q^2; q^2)_k (-q^3; q^2)_k} q^{4k^2 + 3k} \\
&= 1 + \sum_{k=1}^{\infty} \frac{(-1)^k (1 - q^{2k+1})(1 + q^{2k+1})(1 + q)}{(1 - q^2)(1 + q^{2k+1})} q^{4k^2 + 3k}. \tag{4.1.23}
\end{aligned}$$

Rearranging (4.1.23), and then setting  $k = -n - 1$  in the second sum below, we see that

$$\begin{aligned}
\frac{(q^2; q^2)_\infty}{(-q; q^2)_\infty} \sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2 + 2k} &= 1 - q + \sum_{k=1}^{\infty} (-1)^k (1 - q^{2k+1}) q^{4k^2 + 3k} \\
&= 1 - q + \sum_{k=1}^{\infty} (-1)^k q^{4k^2 + 3k} - \sum_{k=1}^{\infty} q^{4k^2 + 5k + 1}
\end{aligned}$$

$$\begin{aligned}
&= 1 - q + \sum_{k=1}^{\infty} (-1)^k q^{4k^2+3k} + \sum_{n=-2}^{-\infty} (-1)^n q^{4n^2+3n} \\
&= \sum_{n=-\infty}^{\infty} (-1)^n q^{4n^2+3n} \\
&= f(-q^7, -q) \\
&= (q; q^8)_{\infty} (q^7; q^8)_{\infty} (q^8; q^8)_{\infty},
\end{aligned}$$

by an appeal to the Jacobi triple product identity (3.1.19). Hence,

$$\begin{aligned}
\sum_{k=0}^{\infty} \frac{(-q; q^2)_k}{(q^2; q^2)_k} q^{k^2+2k} &= \frac{(q^2; q^4)_{\infty} (q; q^8)_{\infty} (q^7; q^8)_{\infty} (q^8; q^8)_{\infty}}{(q; q^2)_{\infty} (q^2; q^2)_{\infty}} \\
&= \frac{1}{(q^3; q^8)_{\infty} (q^4; q^8)_{\infty} (q^5; q^8)_{\infty}},
\end{aligned}$$

after cancellation. Thus, the proof of (4.1.21) is complete.  $\square$

**Theorem 4.1.14** (Combinatorial Versions of the Göllnitz–Gordon Identities). *(a) The first identity in Theorem 4.1.13 is equivalent to the statement: The number of partitions of  $n$  into distinct parts, with at least 2 between parts, and with no consecutive even parts, is equal to the number of partitions of  $n$  into parts congruent to 1, 4, or 7 modulo 8.*

*(b) The second identity in Theorem 4.1.13 is equivalent to the statement: The number of partitions of  $n$  into distinct parts, with at least 2 between parts, with no consecutive even parts, and with all parts  $\geq 3$ , is equal to the number of partitions of  $n$  into parts congruent to 3, 4, or 5 modulo 8.*

**Proof.** On the left-hand side of (4.1.20),  $1/(q^2; q^2)_k$  generates partitions into less than or equal to  $k$  even parts. Consider the Ferrers graph of such a partition. Next,  $(-q; q^2)_k$  generates partitions into less than or equal to  $k$  distinct odd parts. Begin with the largest odd part, say it is  $2m_1 + 1$ . Add 2 to each of the first  $m_1$  rows of the Ferrers graph, and then add 1 to the  $m_1$  plus first row. Consider the next largest odd part, say  $2m_2 + 1$ . Repeat the process, adding 2 to each of the first  $m_2$  rows and 1 to the following row. Note that if we examine two successive parts that are obtained by adjoining 1, then these parts differ by at least two, because the odd parts are distinct. Now recall that  $k^2 = 1 + 3 + \cdots + (2k - 1)$ . Take the Ferrers graph above and add to it, in decreasing order,  $(2k - 1), (2k - 3), \dots, 1$ . Thus, we now have a partition into exactly  $k$  parts. We see that in those instances where we had two successive odd parts before our last additions, that now we will have even parts, and that the even parts will differ by at least 4. In other words, there are no consecutive even parts. The partition-theoretic interpretation of the right-hand side of (4.1.20) is clear.

The proof of the second identity is very similar. The first two steps in constructing a Ferrers graph are the same. Now we note that, referring to the left-hand side of (4.1.21),  $k^2 + 2k = (2k + 1) + (2k - 1) + \cdots + 3$ . Thus, when we add these odd numbers to the

Ferrers graph obtained after the first two steps, the last part of the graph is at least 3. The partition-theoretic interpretation of the right-hand side of (4.1.21) is obvious.  $\square$

**Example 4.1.15.** Let  $n = 8$ . The partitions of 8 into parts differing by at least 2 and with no consecutive even integers are:

$$8, 7 + 1, 6 + 2, 5 + 3.$$

The partitions of 8 into parts congruent to 1, 4, 7 modulo 8 are

$$1 + 1 + 1 + 1 + 1 + 1 + 1 + 1, 4 + 1 + 1 + 1 + 1 + 1, 7 + 1, 4 + 4.$$

The partitions of 8 into parts differing by at least 2, with no consecutive even integers are, and with all parts at least 3 are:

$$8, 5 + 3.$$

The partitions of 8 into parts congruent to 3, 4, 5 modulo 8 are

$$5 + 3, 4 + 4.$$

## 4.2. Weak Form of Bailey's Lemma

**Definition 4.2.1.** Suppose that two sequences  $\alpha_n$  and  $\beta_n$  satisfy the condition

$$\beta_n = \sum_{j=0}^n \frac{\alpha_j}{(q; q)_{n-j} (aq; q)_{n+j}}. \quad (4.2.1)$$

Then  $(\alpha_n, \beta_n)$  called a Bailey pair.

**Theorem 4.2.2** (Bailey's Lemma (Weak form)). Let  $(\alpha_n, \beta_n)$  be a Bailey pair. Then (subject to convergence conditions)

$$\sum_{n=0}^{\infty} q^{n^2} a^n \beta_n = \frac{1}{(aq; q)_{\infty}} \sum_{n=0}^{\infty} q^{n^2} a^n \alpha_n. \quad (4.2.2)$$

**Proof.** Using the definition (4.2.1), we find that

$$\begin{aligned} \sum_{n=0}^{\infty} q^{n^2} a^n \sum_{j=0}^n \frac{\alpha_j}{(q; q)_{n-j} (aq; q)_{n+j}} &= \sum_{j=0}^{\infty} \alpha_j \sum_{n=j}^{\infty} \frac{q^{n^2} a^n}{(q; q)_{n-j} (aq; q)_{n+j}} \\ &= \sum_{j=0}^{\infty} q^{j^2} a^j \alpha_j \sum_{n=0}^{\infty} \frac{q^{n(n+2j)} a^n}{(q; q)_n (aq; q)_{n+2j}} \\ &= \sum_{j=0}^{\infty} q^{j^2} a^j \alpha_j \frac{1}{(aq; q)_{\infty}}, \end{aligned}$$

which gives (4.2.2).  $\square$

**Example 4.2.3.** Let us take the example when  $a = 1$  and  $\beta_n = 1/(q; q)_n$ . Then, trivially,  $\alpha_0 = \beta_0 = 1$ . Next,

$$\beta_1 = \frac{1}{1-q} = \frac{\alpha_0}{(1-q)(1-q)} + \frac{\alpha_1}{(1-q)(1-q^2)}.$$

Thus,

$$\alpha_1 = -q - q^2.$$

We can continue these calculations to find that

$$\begin{aligned}\alpha_2 &= q^5 + q^7 \\ \alpha_3 &= -q^{12} - q^{15} \\ \alpha_4 &= q^{22} + q^{26}.\end{aligned}$$

It appears that that we always obtain two powers of  $q$ , that the signs are alternating, that the difference in powers is equal to the index, and that maybe the exponents are growing quadratically in the index. We are thus led to the conjecture

$$\alpha_n = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{n(3n-1)/2} (1 + q^n), & \text{otherwise.} \end{cases}$$

If we can prove that this pattern persists, then by substituting into Bailey's lemma, we conclude that

$$\begin{aligned}\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} &= \frac{1}{(q; q)_{\infty}} \left( 1 + \sum_{n=1}^{\infty} (-1)^n q^{n(3n-1)/2+n^2} (1 + q^n) \right) \\ &= \frac{1}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} (-1)^n q^{n(5n-1)/2} \\ &= \frac{(q^5; q^5)_{\infty} (q^2; q^5)_{\infty} (q^3; q^5)_{\infty}}{(q; q)_{\infty}} \\ &= \frac{1}{(q; q^5)_{\infty} (q^4; q^5)_{\infty}}.\end{aligned}$$

Thus, we obtain the first Rogers-Ramanujan identity.

Rather than calculate  $\alpha_n$  recursively, it would be far better to calculate  $\alpha_n$  directly in terms of  $\{\beta_n\}$ . We show how to do this in the next lemma.

**Lemma 4.2.4.** If  $(\alpha_n, \beta_n)$  is a Bailey pair, then

$$\alpha_n = \frac{(1 - aq^{2n})}{1 - a} \sum_{j=0}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}} \beta_j}{(q; q)_{n-j}}. \quad (4.2.3)$$

**Proof.** To prove this, we recall the defining relationship (4.2.1). Hence,

$$\begin{aligned}
& \frac{1 - aq^{2n}}{1 - a} \sum_{j=0}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}} \beta_j}{(q; q)_{n-j}} \\
&= \frac{(1 - aq^{2n})}{1 - a} \sum_{j=0}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}}}{(q; q)_{n-j}} \sum_{r=0}^j \frac{\alpha_r}{(q; q)_{j-r} (aq; q)_{j+r}} \\
&= \frac{1 - aq^{2n}}{1 - a} \sum_{r=0}^n \alpha_r \sum_{j=r}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}}}{(q; q)_{n-j} (q; q)_{j-r} (aq; q)_{j+r}} \\
&= \frac{1 - aq^{2n}}{1 - a} \sum_{r=0}^n \alpha_r \sum_{j=0}^{n-r} \frac{(a; q)_{n+j+r} (-1)^{n-j-r} q^{\binom{n-j-r}{2}}}{(q; q)_{n-j-r} (q; q)_j (aq; q)_{j+2r}} \\
&= \frac{1 - aq^{2n}}{1 - a} \sum_{r=0}^n \alpha_r \frac{(a; q)_{n+r} (-1)^{n-r} q^{\binom{n-r}{2}}}{(q; q)_{n-r} (aq; q)_{2r}} \\
&\quad \times \sum_{j=0}^{n-r} \frac{(q^{n-r-j+1}; q)_j (aq^{n+r}; q)_j (-1)^j q^{-j(n-r) + \binom{j+1}{2}}}{(q; q)_j (aq^{2r+1}; q)_j}. \tag{4.2.4}
\end{aligned}$$

Denote the inner sum on the right side of (4.2.4) by  $I(r, q)$ . Then

$$\begin{aligned}
I(r, q) &= \sum_{j=0}^{n-r} \frac{(q^{(n-r)-j+1}; q)_j (aq^{n+r}; q)_j (-1)^j q^{-(n-r)j + \binom{j+1}{2}}}{(q; q)_j (aq^{2r+1}; q)_j} \\
&= \sum_{j=0}^{n-r} \frac{(q^{-(n-r)}; q)_j (aq^{n+r}; q)_j}{(q; q)_j (aq^{2r+1}; q)_j} q^j \\
&= {}_2\phi_1(q^{-(n-r)}, aq^{n+r}; aq^{2r+1}; q, q) \\
&= \frac{(q^{-n+r+1}; q)_{n-r}}{(aq^{2r+1}; q)_{n-r}} \\
&= \delta_{n,r},
\end{aligned}$$

where the penultimate equality follows from the alternative formulation of the  $q$ -Chu–Vandermonde theorem given in Exercise 7 of Chapter 3. Therefore,

$$\frac{1 - aq^{2n}}{1 - a} \sum_{j=0}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}} \beta_j}{(q; q)_{n-j}} = \frac{1 - aq^{2n}}{1 - a} \alpha_n \frac{(a; q)_{2n}}{(aq; q)_{2n}} = \alpha_n,$$

which completes the proof.  $\square$

The proof of Lemma 4.2.4 is unmotivated; it depends on knowing the formula that we want to prove. Another proof of Lemma 4.2.4 arising out of the theory little  $q$ -Jacobi polynomials has been given by Andrews [10].

We now show how the Rogers–Ramanujan identities in Theorem 4.1.8 follow from Lemma 4.2.4. Letting

$$\beta_n = \frac{1}{(q; q)_n},$$

we see that

$$\begin{aligned} \alpha_n &= \frac{1 - aq^{2n}}{1 - a} \sum_{j=0}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}}}{(q; q)_{n-j} (q; q)_j} \\ &= \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{\binom{n}{2}}}{(q; q)_n} \sum_{j=0}^n \frac{(aq^n; q)_j (-1)^j q^{-nj + \binom{j+1}{2}} (q^{n-j+1}; q)_j}{(q; q)_j} \\ &= \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{\binom{n}{2}}}{(q; q)_n} \sum_{j=0}^n \frac{(aq^n; q)_j (q^{-n}; q)_j q^j}{(q; q)_j} \\ &= \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{\binom{n}{2}}}{(q; q)_n} {}_2\phi_1(q^{-n}, aq^n; 0; q, q) \\ &= \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{\binom{n}{2}} a^n q^{n^2}}{(q; q)_n} \\ &= \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{n(3n-1)/2} a^n}{(q; q)_n}, \end{aligned}$$

where the second last equality follows from the  $q$ -Chu–Vandermonde, as given in Exercise 7 of Chapter 3. Hence, by the weak form of Bailey’s Lemma, Theorem 4.2.2,

$$\sum_{n=0}^{\infty} \frac{q^{n^2} a^n}{(q; q)_n} = \frac{1}{(aq; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(1 - aq^{2n})(a; q)_n (-1)^n q^{n(5n-1)/2} a^{2n}}{(1 - a)(q; q)_n}. \quad (4.2.5)$$

Setting  $a = 1$  in (4.2.5), we find that

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} = \frac{1}{(q; q)_{\infty}} \left( 1 + \sum_{n=1}^{\infty} (-1)^n (1 + q^n) q^{n(5n-1)/2} \right).$$

As we have previously seen, this readily reduces to the first Rogers–Ramanujan identity (4.1.15). Setting  $a = q$  in (4.2.5), we arrive at

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(q; q)_n} &= \frac{1}{(q^2; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(1 - q^{2n+1})}{1 - q} (-1)^n q^{n(5n+3)/2} \\ &= \frac{1}{(q; q)_{\infty}} \sum_{n=0}^{\infty} (-1)^n (1 - q^{2n+1}) q^{n(5n+3)/2}, \end{aligned}$$

and the second Rogers–Ramanujan identity (4.1.16) follows as before.

### 4.3. Strong Form of Bailey's Lemma

**Theorem 4.3.1** (Bailey's transform). *Under suitable convergence conditions, if*

$$\beta_n = \sum_{r=0}^n \alpha_r u_{n-r} v_{n+r}$$

and

$$\gamma_n = \sum_{r=n}^{\infty} \delta_r u_{r-n} v_{n+r},$$

then

$$\sum_{n=0}^{\infty} \alpha_n \gamma_n = \sum_{n=0}^{\infty} \beta_n \delta_n.$$

**Proof.** We have

$$\begin{aligned} \sum_{n=0}^{\infty} \alpha_n \gamma_n &= \sum_{n=0}^{\infty} \alpha_n \sum_{r=n}^{\infty} \delta_r u_{r-n} v_{r+n} \\ &= \sum_{r=0}^{\infty} \delta_r \sum_{n=0}^r \alpha_n u_{r-n} v_{r+n} \\ &= \sum_{r=0}^{\infty} \beta_r \delta_r. \end{aligned}$$

(This argument is purely formal and suitable convergence conditions are precisely those that make the series and rearrangements valid.)  $\square$

**Theorem 4.3.2** (Bailey's Lemma). *Suppose that  $(\alpha_n, \beta_n)$  form a Bailey pair. Then, if  $\rho_1$  and  $\rho_2$  are arbitrary non-zero real numbers,  $(\alpha'_n, \beta'_n)$  form a Bailey pair, where*

$$\begin{aligned} \alpha'_n &= \frac{(\rho_1; q)_n (\rho_2; q)_n}{(aq/\rho_1; q)_n (aq/\rho_2; q)_n} \left( \frac{aq}{\rho_1 \rho_2} \right)^n \alpha_n, \\ \beta'_n &= \frac{1}{(aq/\rho_1; q)_n (aq/\rho_2; q)_n} \sum_{j=0}^n \frac{(\rho_1; q)_j (\rho_2; q)_j (aq/(\rho_1 \rho_2); q)_{n-j}}{(q; q)_{n-j}} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \beta_j. \end{aligned}$$

Let us note that this procedure allows us to produce an infinite family of Bailey pairs

$$(\alpha_n, \beta_n) \rightarrow (\alpha'_n, \beta'_n) \rightarrow (\alpha''_n, \beta''_n) \rightarrow \cdots$$

This sequence can be reversed, because given  $(\alpha'_n, \beta'_n)$ , it is clearly possible to write  $\beta_n$  in terms of a sum of  $\beta'_j$ ,  $0 \leq j \leq n$ , and  $\alpha_n$  in terms of  $\alpha'_n$ , so that

$$\cdots \rightarrow (\alpha''_n, \beta''_n) \rightarrow (\alpha'_n, \beta'_n) \rightarrow (\alpha_n, \beta_n).$$

**Proof.** From the definition of Bailey pairs, we want to show that

$$\beta'_n = \sum_{j=0}^n \frac{\alpha'_j}{(q; q)_{n-j} (aq; q)_{n+j}}.$$



First, take

$$\delta_j := \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq^{n-j+1}/(\rho_1 \rho_2); q)_j} \left( \frac{aq}{\rho_1 \rho_2} \right)^j$$

and

$$\gamma_j := \sum_{r=j}^{\infty} \frac{\delta_r}{(q; q)_{r-j} (aq; q)_{r+j}}.$$

Then

$$\begin{aligned} \gamma_j &= \sum_{r=j}^{\infty} \frac{(\rho_1; q)_r (\rho_2; q)_r (q^{n-r+1}; q)_r}{(q; q)_{r-j} (aq; q)_{r+j} (aq^{n-r+1}/\rho_1 \rho_2; q)_r} \left( \frac{aq}{\rho_1 \rho_2} \right)^r \\ &= \sum_{r=0}^{\infty} \frac{(\rho_1; q)_{j+r} (\rho_2; q)_{j+r} (q^{n-j-r+1}; q)_{j+r}}{(q; q)_r (aq; q)_{r+2j} (aq^{n-j-r+1}/\rho_1 \rho_2; q)_{j+r}} \left( \frac{aq}{\rho_1 \rho_2} \right)^{j+r} \\ &= \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq; q)_{2j} (aq^{n-j+1}/(\rho_1 \rho_2); q)_j} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \\ &\quad \times \sum_{r=0}^{\infty} \frac{(\rho_1 q^j; q)_r (\rho_2 q^j; q)_r (q^{(n-j)-r+1}; q)_r}{(q; q)_r (aq^{2j+1}; q)_r (aq^{(n-j)-r+1}/\rho_1 \rho_2; q)_r} \left( \frac{aq}{\rho_1 \rho_2} \right)^r \\ &= \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq; q)_{2j} (aq^{n-j+1}/(\rho_1 \rho_2); q)_j} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \\ &\quad \times {}_3\phi_2(q^{-(n-j)}, \rho_1 q^j, \rho_2 q^j; aq^{2j+1}, \rho_1 \rho_2 q^{-(n-j)}; a; q, q) \\ &= \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq; q)_{2j} (aq^{n-j+1}/(\rho_1 \rho_2); q)_j} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \frac{(aq^{j+1}/\rho_1; q)_{n-j} (aq^{j+1}/\rho_2; q)_{n-j}}{(aq^{2j+1}; q)_{n-j} (aq/\rho_1 \rho_2; q)_{n-j}} \\ &= \frac{(aq/\rho_1; q)_n (aq/\rho_2; q)_n}{(aq; q)_{n+j} (aq/(\rho_1 \rho_2); q)_n} \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq/\rho_1; q)_j (aq/\rho_2; q)_j} \left( \frac{aq}{\rho_1 \rho_2} \right)^j, \end{aligned} \quad (4.3.1)$$

where in the penultimate line we used the Pfaff–Saalschütz theorem, Theorem 3.3.1.

Next, using Bailey's transform, Theorem 4.3.1, and (4.3.1), below, we find that

$$\begin{aligned} &\sum_{j=0}^n \frac{\alpha'_j}{(q; q)_{n-j} (aq; q)_{n+j}} \\ &= \sum_{j=0}^n \frac{(\rho_1; q)_j (\rho_2; q)_j \alpha_j}{(aq/\rho_1; q)_j (aq/\rho_2; q)_j (q; q)_{n-j} (aq; q)_{n+j}} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \\ &= \frac{1}{(q; q)_n} \sum_{j=0}^n \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq/\rho_1; q)_j (aq/\rho_2; q)_j (aq; q)_{n+j}} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \alpha_j \\ &= \frac{(aq/(\rho_1 \rho_2); q)_n}{(aq/\rho_1; q)_n (aq/\rho_2; q)_n (q; q)_n} \\ &\quad \times \sum_{j=0}^{\infty} \frac{(aq/\rho_1; q)_n (aq/\rho_2; q)_n}{(aq/(\rho_1 \rho_2); q)_n (aq; q)_{n+j}} \frac{(\rho_1; q)_j (\rho_2; q)_j (q^{n-j+1}; q)_j}{(aq/\rho_1; q)_j (aq/\rho_2; q)_j} \left( \frac{aq}{\rho_1 \rho_2} \right)^j \alpha_j \end{aligned}$$

$$\begin{aligned}
&= \frac{(aq/(\rho_1\rho_2); q)_n}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n(q; q)_n} \sum_{j=0}^{\infty} \alpha_j \gamma_j \\
&= \frac{(aq/(\rho_1\rho_2); q)_n}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n(q; q)_n} \sum_{j=0}^{\infty} \beta_j \delta_j \\
&= \frac{(aq/(\rho_1\rho_2); q)_n}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n(q; q)_n} \sum_{j=0}^{\infty} \frac{(\rho_1; q)_j(\rho_2; q)_j(q^{n-j+1}; q)_j}{(aq^{n-j+1}/(\rho_1\rho_2); q)_j} \left(\frac{aq}{\rho_1\rho_2}\right)^j \beta_j \\
&= \frac{1}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n} \sum_{j=0}^{\infty} \frac{(\rho_1; q)_j(\rho_2; q)_j(aq/(\rho_1\rho_2); q)_{n-j}}{(q; q)_{n-j}} \left(\frac{aq}{\rho_1\rho_2}\right)^j \beta_j \\
&= \beta'_n,
\end{aligned}$$

which is what we wanted to prove.  $\square$

#### 4.4. Applications of Bailey's Lemma

Recall from (4.2.3) that if  $(\alpha_n, \beta_n)$  is a Bailey pair, then

$$\alpha_n = \frac{1 - aq^{2n}}{1 - a} \sum_{j=0}^n \frac{(a; q)_{n+j} (-1)^{n-j} q^{\binom{n-j}{2}} \beta_j}{(q; q)_{n-j}}.$$

It follows that a simple Bailey pair is given by

$$\beta_n = \begin{cases} 1, & \text{if } n = 0, \\ 0, & \text{if } n > 0, \end{cases} \quad (4.4.1)$$

$$\alpha_n = \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{\binom{n}{2}}}{(q; q)_n}. \quad (4.4.2)$$

We now apply Bailey's Lemma, Theorem 4.3.2, to deduce a second Bailey pair

$$\alpha'_n = \frac{(\rho_1; q)_n(\rho_2; q)_n}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n} \left(\frac{aq}{\rho_1\rho_2}\right)^n \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n (-1)^n q^{\binom{n}{2}}}{(q; q)_n}, \quad (4.4.3)$$

$$\beta'_n = \frac{(aq/(\rho_1\rho_2); q)_n}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n(q; q)_n}, \quad (4.4.4)$$

where  $\rho_1\rho_2 \neq 0$ . If we repeat this process, again using Bailey's Lemma, i.e., Theorem 4.3.2, we find another Bailey pair, provided that  $\sigma_1\sigma_2 \neq 0$ ,

$$\alpha_n'' = \frac{(\sigma_1; q)_n(\sigma_2; q)_n}{(aq/\sigma_1; q)_n(aq/\sigma_2; q)_n} \left(\frac{aq}{\sigma_1\sigma_2}\right)^n \frac{(\rho_1; q)_n(\rho_2; q)_n}{(aq/\rho_1; q)_n(aq/\rho_2; q)_n} \left(\frac{aq}{\rho_1\rho_2}\right)^n \\ \times \frac{1 - aq^{2n}}{1 - a} \frac{(a; q)_n(-1)^n q^{\binom{n}{2}}}{(q; q)_n}, \quad (4.4.5)$$

$$\beta_n'' = \frac{1}{(aq/\sigma_1; q)_n(aq/\sigma_2; q)_n} \sum_{j=0}^n \frac{(\sigma_1; q)_j(\sigma_2; q)_j(aq/(\sigma_1\sigma_2); q)_{n-j}}{(q; q)_{n-j}} \left(\frac{aq}{\sigma_1\sigma_2}\right)^j \\ \times \frac{(aq/(\rho_1\rho_2); q)_j}{(aq/\rho_1; q)_j(aq/\rho_2; q)_j(q; q)_j}. \quad (4.4.6)$$

Let  $a \rightarrow 1$  and  $\rho_1, \rho_2 \rightarrow \infty$  in (4.4.1)–(4.4.6). Then, we readily deduce that

$$\alpha_n = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{n(n-1)/2} (1 + q^n), & \text{otherwise,} \end{cases}$$

$$\alpha_n' = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{n(3n-1)/2} (1 + q^n), & \text{otherwise,} \end{cases} \\ \beta_n' = \frac{1}{(q; q)_n},$$

$$\alpha_n'' = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{n(5n-1)/2} (1 + q^n), & \text{otherwise,} \end{cases} \\ \beta_n'' = \sum_{j=0}^n \frac{q^{j^2}}{(q; q)_{n-j}(q; q)_j}.$$

Substituting  $(\alpha_n'', \beta_n'')$  into the weak form of Bailey's Lemma, Theorem 4.2.2, and employing the Jacobi triple product identity (1.1.7), we deduce that

$$\sum_{n=0}^{\infty} \sum_{j=0}^n \frac{q^{n^2+j^2}}{(q; q)_{n-j}(q; q)_j} = \frac{1}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} (-1)^n q^{n(7n-1)/2} \\ = \frac{1}{(q; q)_{\infty}} f(-q^3, -q^4) \\ = \frac{1}{(q; q^7)_{\infty}(q^2; q^7)_{\infty}(q^5; q^7)_{\infty}(q^6; q^7)_{\infty}}. \quad (4.4.7)$$

If we continue in this fashion and calculate  $k$  Bailey pairs, we find that

$$\begin{aligned} \sum_{n_1 \geq n_2 \geq \dots \geq n_k \geq 0} \frac{q^{n_1^2 + \dots + n_k^2}}{(q; q)_{n_1 - n_2} \cdots (q; q)_{n_{k-1} - n_k} (q; q)_{n_k}} &= \frac{1}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n q^{n((2k+3)n-1)/2} \\ &= \frac{(q^{k+1}; q^{2k+3})_\infty (q^{k+2}; q^{2k+3})_\infty (q^{2k+3}; q^{2k+3})_\infty}{(q; q)_\infty} \\ &= \prod_{r \neq 0, \pm(k+1) \pmod{2k+3}} \frac{1}{(q^r; q^{2k+3})_\infty}. \end{aligned} \quad (4.4.8)$$

The identity (4.4.8) is equivalent to the following partition theorem.

**Theorem 4.4.1** (Andrews–Gordon). *Let  $A_k(n)$  denote the number of partitions  $\lambda$  of  $n$  with parts satisfying the difference conditions*

$$\lambda_i - \lambda_{i+k} \geq 2. \quad (4.4.9)$$

Then

$$\sum_{n=0}^{\infty} A_k(n) q^n = \prod_{r \neq 0, \pm(k+1) \pmod{2k+3}} \frac{1}{(q^r; q^{2k+3})_\infty}$$

Gordon proved this theorem combinatorially. Later, Andrews showed that the generating function of  $A_k(n)$  is the left-hand side of (4.4.8).

## 4.5. Exercises

1. Give a proof of Lebesgue's Identity, Theorem 4.1.1, by using Heine's transformation, Theorem 3.2.1.
2. Give a combinatorial proof of Lebesgue's Identity, Theorem 4.1.1.
3. Find a combinatorial interpretation for the left side of (4.4.7).
4. If successful in the previous exercise, find a combinatorial proof of (4.4.7).

## Analogues of Euler's Theorem; Partitions with Gaps

### 5.1. Motivation

Before proceeding further, let us introduce a standard, more economical notation for the partition of an integer. Instead of writing, for example, a partition with seven 5's in it by  $\cdots 5 + 5 + 5 + 5 + 5 + 5 + 5 + \cdots$ , we shall write  $5^7$ . We give further examples below.

Recall Euler's fundamental theorem: The number of partitions of a positive integer  $n$  into distinct parts is identical to the number of partitions of  $n$  into odd parts. If we arrange the parts of a partition of  $n$  in descending order, say,  $a_1 > a_2 > \cdots > a_j$ , then the distance between successive parts is at least 1. On the other hand, if we have a partition of  $n$  into odd parts, say,  $b_1 > b_2 > \cdots > b_r$ , then  $b_i - b_j \equiv 0 \pmod{2}$ , or in other words,  $b_i \equiv 1 \pmod{2}$ . Recall the definitions of the Rogers–Ramanujan functions,

$$G(q) := \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} \quad \text{and} \quad H(q) := \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q; q)_n}.$$

The famous Rogers–Ramanujan identities are given by

$$G(q) = \frac{1}{(q; q^5)_{\infty} (q^4; q^5)_{\infty}} \quad \text{and} \quad H(q) = \frac{1}{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty}}. \quad (5.1.1)$$

The first identity is equivalent to: The set of partitions of  $n$  wherein the difference between any two parts is at least 2 is equinumerous with the set of partitions into parts congruent to either 1 or 4 modulo 5, i.e. the parts are  $\equiv \pm 1 \pmod{5}$ .

The natural question that one should ask now is: Suppose the parts of a partition have a distance of at least 3 between successive parts. Is this set of partitions equinumerous with a set of partitions in which the parts belong to prescribed residue classes?

Suppose we have a class of partitions such that the gap between any two parts is at least 3. If we perform some calculations, it appears that partitions with parts congruent to  $1, 5 \pmod{6}$  are equinumerous with partitions with a distance of at least 3 between parts. In fact, the sizes of these sets of partitions are the same for  $1 \leq n \leq 8$ . However, when we reach 9, we find a discrepancy. The partitions of 9 with a gap of at least 3 between parts are:  $9, 8+1, 7+2, 6+3$ . The partitions with parts congruent to 1 or 5 modulo 6 are:  $71^2, 51^4, 1^9$ . The first set has four representations, and the second set has 3 representations. For  $n = 10, 11, 12, 13, 14$ , the two aforementioned classes of partitions are equinumerous. Consider now  $n = 15$ . Those partitions with gaps of at least 3 between successive parts are:  $15, 14+1, 13+2, 12+3, 11+4, 10+5, 10+4+1, 9+6, 9+5+1, 8+5+2$ . Those partitions of 15 with all parts congruent to 1 or 5 modulo 6 are:  $1^2 13, 1^4 11, 17^2, 1^3 57, 1^8 7, 5^3, 1^5 5^2, 1^{10} 5, 1^{15}$ . There are 10 elements in the first class, and 9 in the second class. Is there a requirement that would eliminate some of the partitions in the first class? Readers should refrain here from peeking at the answer, which will be given at the beginning of the next paragraph.

What we need to eliminate are those representations with consecutive multiples of 3, i.e., remove  $6 + 3$  and  $9 + 6$ , respectively, from the two sets of partitions with gaps of at least 3 above. If we do so in the two examples above, or in any of the other cases, we will find that the sets are equinumerous.

We offer a remark here to indicate that, generally, if we prescribe the congruence conditions for the parts of partitions, then it is more difficult to find, if possible, the corresponding partitions with gaps. Normally, it is easier to begin with a description of the partitions with gaps. Depending on the starting conditions, we thus want to compute  $b_n$  or  $a_n$  in the generating function  $F(q)$ , which we write in the form

$$F(q) := \sum_{n=0}^{\infty} b_n q^n = \prod_{n=1}^{\infty} (1 - q^n)^{-a_n}, \quad b_0 = 1. \quad (5.1.2)$$

In our example,  $a_n = 0$ , when  $n \equiv 0, 2, 3, 4 \pmod{6}$ ;  $a_n = 1$ , when  $n \equiv 1, 5 \pmod{6}$ .

Our goal is to compute a generating function for the sequence or sequences satisfying the aforementioned gap and congruence conditions. Taking the logarithmic derivative of (5.1.2), we find that

$$\frac{F'(q)}{F(q)} = -\frac{d}{dq} \sum_{n=1}^{\infty} a_n \log(1 - q^n) = \sum_{n=1}^{\infty} \frac{a_n n q^{n-1}}{1 - q^n},$$

from which it follows that

$$\begin{aligned} qF'(q) &= F(q) \sum_{n=1}^{\infty} \frac{a_n n q^n}{1 - q^n} = F(q) \sum_{n=1}^{\infty} a_n n \sum_{m=1}^{\infty} q^{mn} \\ &= F(q) \sum_{N=1}^{\infty} q^N \sum_{n|N} n a_n = F(q) \sum_{N=1}^{\infty} D_N q^N, \end{aligned} \quad (5.1.3)$$

where

$$D_N = \sum_{n|N} na_n.$$

Equating coefficients of  $q^M$ ,  $M \geq 1$ , on both sides of (5.1.3) and using (5.1.2), we find that

$$Mb_M = \sum_{j=0}^M b_j D_{M-j}, \quad \text{with } D_0 = 0.$$

Thus, we have found a recurrence relation for  $\{b_M\}$  in terms of  $\{a_j\}$ . Conversely, given  $\{b_j\}$ , we can calculate  $\{a_j\}$ . As an example, if  $a_n \equiv 1$ , then  $b_n = p(n)$ , and furthermore

$$Mp(M) = \sum_{j=0}^{M-1} p(j)\sigma(M-j).$$

which is in the spirit of several results that we had derived in the last portions of Chapter 1.

As with our example,  $a_n = 0$ , when  $n \equiv 0, 2, 3, 4 \pmod{6}$ ;  $a_n = 1$ , when  $n \equiv 1, 5 \pmod{6}$ , it is usually more difficult to impose congruence conditions than it is to impose gap conditions. Let  $\sigma(\pi)$  denote a partition, and let  $\nu(\pi)$  denote the number of parts of  $\pi$ . Consider

$$f(z, q) := \sum_{\substack{\pi \\ \text{min. dif.} \geq 3 \\ \text{no consec. mult. of } 3}} z^{\nu(\pi)} q^{\sigma(\pi)} = \sum_{m, n=0}^{\infty} c(m, n) z^m q^n. \quad (5.1.4)$$

Let  $f_i(z, q)$  denote the same function as in (5.1.4), but with the added condition that the smallest part of  $\pi$  is  $> i$ . Thus,

$$f_0(z, q) = f(z, q),$$

and

$$f_0(z, q) - f_1(z, q)$$

generates partitions with minimal part equal to 1. Now delete 1 from all the partitions  $\pi$ . Subtract 3 from each of the remaining parts. Note that before subtracting 3 from each part, each part now is  $\geq 4$ . Thus,

$$f_0(z, q) - f_1(z, q) = zqf_0(zq^3, q). \quad (5.1.5)$$

Now observe that  $f_1(z, q) - f_2(z, q)$  is a general function of admissible partitions with 2 appearing. We repeat the argument above by subtracting 2 from each part of each partition. Hence, deleting 2 from each part of each partition  $\pi$  and subtracting 3 from each part, we find that

$$f_1(z, q) - f_2(z, q) = zq^2f_1(zq^3, q). \quad (5.1.6)$$

Repeat the argument, but now recall that no consecutive multiples of 3 appear. So instead of all the parts being greater than 2, they really are all greater than 3. Hence,

$$f_2(z, q) - f_3(z, q) = zq^3f_3(zq^3, q). \quad (5.1.7)$$

If we now subtract 3 from each part, we are back to our original setting, i.e.,

$$f_3(z, q) = f_0(zq^3, q). \quad (5.1.8)$$

Rewrite (5.1.5) in the form

$$f_1(z, q) = f_0(z, q) - zqf_0(zq^3, q). \quad (5.1.9)$$

Substitute (5.1.8) into (5.1.7) twice, once with  $z$  replaced by  $zq^3$ , to obtain

$$f_2(z, q) = f_0(zq^3, q) + zq^3f_0(zq^6, q). \quad (5.1.10)$$

Now, substitute (5.1.9) and (5.1.10) into (5.1.6) to deduce that

$$\begin{aligned} \{f_0(z, q) - zqf_0(zq^3, q)\} - \{f_0(zq^3, q) + zq^3f_0(zq^6, q)\} \\ = zq^2\{f_0(zq^3, q) - zq^4f_0(zq^6, q)\}, \end{aligned}$$

which we solve for  $f_0(z, q)$ . We now delete the second argument, so that we have

$$f_0(z) = (1 + zq + zq^2)f_0(zq^3) + (zq^3 - z^2q^6)f_0(zq^6). \quad (5.1.11)$$

Now define

$$\varphi(z) = \frac{f_0(z)}{(z; q^3)_\infty}. \quad (5.1.12)$$

Thus, if we divide both sides by  $(zq^3; q^3)_\infty$ , (5.1.11) can be rewritten in the form

$$\begin{aligned} (1 - z)\varphi(z) &= (1 + zq + zq^2)\varphi(zq^3) + zq^3(1 - zq^3)\frac{f_0(zq^6)}{(zq^3; q^3)_\infty} \\ &= (1 + zq + zq^2)\varphi(zq^3) + zq^3\varphi(zq^6). \end{aligned} \quad (5.1.13)$$

Let

$$\varphi(z) = \sum_{n=0}^{\infty} \alpha_n z^n, \quad \varphi(0) = \alpha_0 = f_0(0) = 1. \quad (5.1.14)$$

Substitute (5.1.14) into (5.1.13) and equate coefficients of  $z^n$  to deduce that

$$\alpha_n - \alpha_{n-1} = \alpha_n q^{3n} + \alpha_{n-1} q^{3n-2} + \alpha_{n-1} q^{3n-1} + \alpha_{n-1} q^{6n-3},$$

which can be rewritten in the shape

$$\alpha_n = \frac{(1 + q^{3n-2})(1 + q^{3n-1})}{1 - q^{3n}} \alpha_{n-1}. \quad (5.1.15)$$

Iterating (5.1.15)  $n$  times and employing the initial condition  $\alpha_0 = 1$ , we deduce that

$$\alpha_n = \frac{(-q; q^3)_n (-q^2; q^3)_n}{(q^3; q^3)_n}. \quad (5.1.16)$$

We now can conclude from (5.1.16) and (5.1.12) that

$$\begin{aligned} f(z, q) = f_0(z, q) = f_0(z) &= (z; q^3)_\infty \sum_{n=0}^{\infty} \frac{(-q; q^3)_n (-q^2; q^3)_n}{(q^3; q^3)_n} z^n \\ &= (z; q^3)_\infty {}_2\phi_1(-q, -q^2; 0; q^3; z). \end{aligned} \quad (5.1.17)$$



We now apply Heine's transformation, Theorem 3.2.1, with  $q$  replaced by  $q^3$ ,  $a = -q$ ,  $b = -q^2$ , and  $c = 0$ , on the right-hand side of (5.1.17) to deduce that

$$f(z, q) = (-q^2; q^3)_\infty (-qz; q^3)_\infty {}_2\phi_1(0, z; -qz; q^3; -q^2). \quad (5.1.18)$$

Let  $z = 1$  and note that  $(1)_n = 0$ ,  $n \geq 1$ . Thus,

$${}_2\phi_1(0, 1; -q; q^3; -q^2) = 1.$$

Thus, from (5.1.18), we can conclude that

$$\begin{aligned} f(1, q) &= (-q^2; q^3)_\infty (-q; q^3)_\infty \\ &= \frac{(-q; q^3)_\infty (-q^2; q^3)_\infty (q; q^3)_\infty (q^2; q^3)_\infty}{(q; q^3)_\infty (q^2; q^3)_\infty} \\ &= \frac{(q^2; q^6)_\infty (q^4; q^6)_\infty}{(q; q^3)_\infty (q^2; q^3)_\infty} \\ &= \frac{(q^2; q^6)_\infty (q^4; q^6)_\infty}{(q; q^6)_\infty (q^4; q^6)_\infty (q^2; q^6)_\infty (q^5; q^6)_\infty} \\ &= \frac{1}{(q; q^6)_\infty (q^5; q^6)_\infty}. \end{aligned} \quad (5.1.19)$$

We have just proved the following theorem, originally due to I. Schur in 1926 [95].

**Theorem 5.1.1** (Schur). *The set of partitions of  $n$  with minimal difference at least equal to 3, and with no consecutive multiples of 3 is equinumerous with the set of partitions of  $n$  into parts that are congruent to 1 or 5 modulo 6.*

**Definition 5.1.2.** *Let  $S_1(n)$  denote the number of partitions of  $n$  into parts which differ by at least 3 and with no consecutive multiples of 3.*

*Let  $S_2(n)$  denote the number of partitions of  $n$  into distinct parts, with none of them divisible by 3.*

*Let  $S_3(n)$  denote the number of partitions of  $n$  into parts congruent to either 1 or 5 modulo 6.*

From the far right side of (5.1.19) and our construction, we have proved the following theorem.

**Theorem 5.1.3.** *We have*

$$\sum_{n=0}^{\infty} S_1(n)q^n = \frac{1}{(q; q^6)_\infty (q^5; q^6)_\infty}. \quad (5.1.20)$$

The next theorem is trivial.

**Theorem 5.1.4.** *We have*

$$\sum_{n=0}^{\infty} S_2(n)q^n = (-q; q^3)_\infty (-q^2; q^3)_\infty.$$

$n$	partitions, $R_2(n)$	partitions in $S$	conclusion
1	1	1	$1 \in S$
2	2	$1 + 1$	2 not $\in S$
3	3	$1 + 1 + 1$	3 not $\in S$
4	$4, 3 + 1$	$1^4, 4$	$4 \in S$
5	$5, 4 + 1$	$1^5, 4 + 1$	5 not $\in S$
6	$6, 5 + 1$	$1^6, 4 + 1 + 1$	6 not $\in S$
7	$7, 6 + 1, 5 + 2$	$1^7, 41^3, 7$	$7 \in S$
8	$8, 71, 62, 53$	$1^8, 4^2, 41^4, 71$	8 not $\in S$
9	$9, 81, 72, 63, 531$	$1^9, 4^2 1, 41^5, 71^2, 9$	$9 \in S$
10	$(10), 91, 82, 73, 631$	$1^{10}, 4^2 1^2, 41^6, 71^3, 91$	10 not $\in S$
11	$(11), (10)1, 92, 83, 74, 731$	$1^{11}, 4^2 1^3, 41^7, 71^4, 91^2, 74$	11 not $\in S$

**Table 1.** Partitions Enumerated by  $R_2(n)$

From the far left side of (5.1.18) and the two previous theorems, we have established the following corollary.

**Corollary 5.1.5.** *We have*

$$S_1(n) = S_2(n) = S_3(n).$$

**Definition 5.1.6.** *Let  $R_d(n)$  denote the number of partitions of  $n$  into parts that differ by at least  $d$  with no consecutive multiples of  $d$  allowed.*

Note that, by Theorem 5.1.3, for  $d = 3$ , we have proved that these partitions are equinumerous with the set of partitions into parts congruent to 1 or 5 modulo 6. For  $d = 1$ , we cannot have partitions into parts with no consecutive multiples of 1. However, if we drop the requirement that no consecutive multiples of 2 are allowed but require that the parts differ by at least 2, then these partitions, by the first Rogers–Ramanujan identity, are equinumerous with the set of partitions into parts that are congruent to 1 or 4 modulo 5. Let us see if we can find a generating function for  $R_2(n)$ , or find a set of congruences, say,  $S = S(n)$ , that is identical in number with the cardinality of  $R_2(n)$ .

Let us explain the reasoning in the right-hand column of Table 1. We see that 1 must belong to the targeted set  $S$ , for otherwise we would have no partitions in  $S$ , and the

desired equal cardinalities could not happen. If  $2 \in S$ , then we would have two partitions in  $S$ , and so equal cardinalities would be impossible. The same reasoning holds for 3. There are two partitions enumerated by  $R_2(4)$ , and so to get two partitions in  $S$ , 4 must belong to  $S$ . Proceeding in this manner, up to  $n = 11$ , we find that  $1, 4, 7, 9 \in S$ . We see that  $S$  cannot be described by congruence classes modulo 2, 3, 5, 6, or 7. However, congruence classes modulo 8 might be possible from this limited table. Indeed, this conjecture is correct.

**Theorem 5.1.7.** *The cardinality of  $R_2(n)$ ,  $n \geq 1$ , is equal to that of the subset  $S$ , where*

$$S = \{n : n \equiv 1, 4, 7 \pmod{8}; n \geq 1\}.$$

**Proof.** Recall that  $\sigma(\pi)$  denotes a partition and  $\nu(\pi)$  denotes the number of parts of  $\pi$ . Define

$$g_i(z, q) := \sum_{\substack{\pi \\ \text{parts differ by } \geq 2 \\ \text{no consec. mult. of } 2 \\ \text{each part} > i}} z^{\nu(\pi)} q^{\sigma(\pi)}. \quad (5.1.21)$$

Deleting the argument  $q$  from our notation, we proceed exactly in the same manner as we did in the proof of Theorem 5.1.3. The partitions in the difference  $g_0(z) - g_1(z)$  have minimal part equal to 1. We delete 1 from each partition, and so

$$g_0(z) - g_1(z) = zqg_0(zq^2), \quad \text{or} \quad g_1(z) = g_0(z) - zqg_0(zq^2). \quad (5.1.22)$$

Deleting 2 from each partition and remembering that no consecutive multiples of 2 are allowed, we find that

$$g_1(z) - g_2(z) = zq^2g_2(zq^2). \quad (5.1.23)$$

Furthermore,

$$g_2(z) = g_0(zq^2). \quad (5.1.24)$$

Put (5.1.22) and (5.1.24) into (5.1.23) to find that

$$\{g_0(z) - zqg_0(zq^2)\} - g_0(zq^2) = zq^2g_0(zq^4),$$

which we rearrange in the form

$$g_0(z) = (1 + zq)g_0(zq^2) + zq^2g_0(zq^4). \quad (5.1.25)$$

Now let

$$g_0(z) = \sum_{n=0}^{\infty} a_n z^n, \quad g_0(0) = a_0 = 1. \quad (5.1.26)$$

Putting (5.1.26) into (5.1.25) and equating coefficients of  $z^n$ ,  $n \geq 1$ , on both sides, we find that

$$a_n = a_n q^{2n} + a_{n-1} q^{2n-1} + a_{n-1} q^{4n-2},$$

which we reformulate in the recurrence relation

$$a_n = \frac{q^{2n-1}(1 + q^{2n-1})}{1 - q^{2n}} a_{n-1}, \quad n \geq 1. \quad (5.1.27)$$

Iterating (5.1.27)  $n$  times and recalling that  $a_0 = 1$ , we deduce that

$$a_n = \frac{q^{n^2}(-q; q^2)_n}{(q^2; q^2)_n}, \quad n \geq 0, \quad (5.1.28)$$

which, when put in (5.1.26), yields

$$g_0(z) = \sum_{n=0}^{\infty} \frac{(-q; q^2)_n}{(q^2; q^2)_n} q^{n^2} z^n.$$

To complete the proof of the theorem that we sought, rewritten combinatorially below, we need to evaluate  $g_0(1)$ , which is not easy. One needs to use Watson's  $q$ -analogue of Whipple's theorem. In fact,

$$g_0(1) = \sum_{n=0}^{\infty} \frac{(-q; q^2)_n}{(q^2; q^2)_n} q^{n^2} = \frac{1}{(q; q^8)_{\infty} (q^4; q^8)_{\infty} (q^7; q^8)_{\infty}}. \quad (5.1.29)$$

□

We have just then “almost proved” the following theorem.

**Theorem 5.1.8.** *The number of partitions of  $n$  with parts differing by at least 2 and with no consecutive multiples of 2 is equal to the number of partitions of  $n$  into parts congruent to 1, 4, 7 modulo 8.*

It is natural to ask if Theorem 5.1.8 has an analogue if we replace “no consecutive multiples of 2” by “no successive odd parts.” The next result provides an affirmative answer.

**Theorem 5.1.9.** *The number of partitions of  $n$  into distinct parts, with a difference of at least 2 between parts, with no successive odd parts, and with each part at least 2 is equal to the number of partitions of  $n$  into parts congruent to 2, 3, 7 modulo 8.*

**Proof.** Let

$$g_i(z, q) := g_i(z) := \sum_{\substack{\pi \\ \text{parts differ by at least 2} \\ \text{no consecutive odd parts} \\ \text{each part} > i}} z^{\nu(\pi)} q^{\sigma(\pi)}.$$

We want to determine  $g_1(1)$ . First, observe that

$$g_3(z) = g_1(zq^2). \quad (5.1.30)$$

Second, observe that if we want to determine those partitions with smallest part equal to 2, then

$$g_1(z) - g_2(z) = zq^2 g_1(zq^2). \quad (5.1.31)$$

If we want to determine those partitions with smallest part precisely equal to 3, then

$$g_2(z) - g_3(z) = zq^3 g_3(zq^2). \quad (5.1.32)$$

One might think that on the right side of (5.1.32), we should have  $g_2(zq^2)$  instead of  $g_3(zq^2)$ . But remember that we are forbidden to have the successive pair of parts, 3, 5, and for this reason  $g_3$  appears, and not  $g_2$ . Using (5.1.30)–(5.1.32), we see that

$$\begin{aligned} g_1(z) &= \{g_3(z) + zq^3g_3(zq^2)\} + zq^2g_1(zq^2) \\ &= g_1(zq^2) + zq^3g_1(zq^4) + zq^2g_1(zq^2) \\ &= (1 + zq^2)g_1(zq^2) + zq^3g_1(zq^4). \end{aligned} \quad (5.1.33)$$

If we set

$$g_1(z) = \sum_{n=0}^{\infty} a_n z^n, \quad g_1(0) = a_0 = 1,$$

then equating coefficients of  $q^n$  on both sides of (5.1.33), we deduce that

$$a_n(1 - q^{2n}) = q^{2n}(1 + q^{2n-1})a_{n-1}, \quad n \geq 1,$$

or

$$a_n = \frac{q^{2n}(1 + q^{2n-1})}{1 - q^{2n}} a_{n-1}.$$

If we iterate the recurrence relation above and note that  $a_0 = 1$ , we find that

$$a_n = \frac{(-q; q^2)_n}{(q^2; q^2)_n} q^{n^2+n}, \quad n \geq 1.$$

Hence,

$$g_1(z) = \sum_{n=0}^{\infty} \frac{(-q; q^2)_n}{(q^2; q^2)_n} q^{n^2+n} z^n.$$

To evaluate  $g_1(1)$ , we apply Lebesgue's theorem, Theorem 4.1.1, with  $q$  replaced by  $q^2$  and with  $b = 1/q$ . Hence,

$$\begin{aligned} g_1(1) &= \sum_{n=0}^{\infty} \frac{(-q; q^2)_n}{(q^2; q^2)_n} q^{n^2+n} = \frac{(-q^3; q^4)_{\infty}}{(q^2; q^4)_{\infty}} \\ &= \frac{(q^6; q^8)_{\infty}}{(q^2; q^4)_{\infty} (q^3; q^4)_{\infty}} \\ &= \frac{1}{(q^2; q^8)_{\infty} (q^3; q^8)_{\infty} (q^7; q^8)_{\infty}}, \end{aligned} \quad (5.1.34)$$

which completes the proof of Theorem 5.1.9.  $\square$

**Definition 5.1.10.** Let  $q_d(n)$  denote the number of partitions of  $n$  with parts differing by at least  $d$ . Let  $Q_d(n)$  be the number of partitions of  $n$  with parts congruent to  $\pm 1 \pmod{d+3}$ .

In 1956, H. L. Alder [4] made the following conjecture.

**Conjecture 5.1.11.**

$$\Delta_d(n) := q_d(n) - Q_d(n) \geq 0, \quad \text{for all } d \text{ and } n. \quad (5.1.35)$$

Let us examine some special cases. Suppose that  $d = 1$ . Then  $q_1(n)$  denotes the number of partitions into distinct parts, while  $Q_1(n)$  denotes the number of partitions into parts  $\equiv \pm 1 \pmod{4}$ , i.e., odd parts. By Euler's Theorem,  $\Delta_1(n) = 0$ . Thus (5.1.35) is valid for  $d = 1$ .

Take  $d = 2$ . Thus,  $q_2(n)$  stands for the number of partitions of  $n$  into parts differing by at least 2, while  $Q_2(n)$  denotes the number of partitions of  $n$  into parts congruent to  $\pm 1 \pmod{5}$ . By the first Rogers–Ramanujan identity,  $q_2(n) = Q_2(n)$ , and so  $\Delta_2(n) = 0$ . Therefore, Alder's conjecture is valid for  $d = 2$ .

Let  $d = 3$ . Then  $q_3(n)$  denotes the number of partitions of  $n$  into parts differing by at least 3, and  $Q_3(n)$  denotes the number of partitions of  $n$  into parts congruent to  $\pm 1 \pmod{6}$ . Let  $q_{c3}(n)$  denote the number of partitions of  $n$  into parts differing by at least 3 and with at least one consecutive multiple of 3. Then Schur's Theorem can be cast in the form  $q_3(n) - q_{c3}(n) = Q_3(n)$ . Thus,

$$\Delta_3(n) = q_3(n) - Q_3(n) = q_3(n) - \{q_3(n) - q_{c3}(n)\} = q_{c3}(n) \geq 0.$$

In 1971, Andrews [7] proved Alder's Conjecture for  $d = 2^r - 1$ ,  $r \geq 4$ . No further progress was made for over 30 years until A. J. Yee [105], [106] proved Alder's Conjecture for all  $d \geq 31$  and for  $d = 7$ . Finally, C. Alfes, M. Jameson, and R. Lemke Oliver [5] used the *circle method* and computation to handle the remaining cases.

The next theorem, at first glance, does not appear to have an aesthetic appeal, and the proof does not seem to be enchanting either. It is due to J. J. Sylvester [99], and will serve as motivation to one of the most elementary proofs of the Rogers–Ramanujan identities, due to Andrews. The proof itself also has interesting consequences, which we will point out along the way. However, the method is deceptively elegant and powerful.

**Theorem 5.1.12.** *Let*

$$\begin{aligned} S(N; x, q) = 1 + \sum_{j=1}^{\infty} \begin{bmatrix} N+1-j \\ j \end{bmatrix} (-xq; q)_{j-1} q^{j(3j-1)/2} x^j \\ + \sum_{j=1}^{\infty} \begin{bmatrix} N-j \\ j \end{bmatrix} (-xq; q)_{j-1} q^{3j(j+1)/2} x^{j+1}. \end{aligned} \quad (5.1.36)$$

*Then, for each positive integer  $N$ ,*

$$S(N; x, q) = (-xq; q)_N. \quad (5.1.37)$$

Observe that both sums in (5.1.36) are finite.

**Proof.** We first combine together the two sums in (5.1.36), while also using the definition of the Gaussian binomial coefficients. Then we add and subtract  $q^j$  within the expression in curly brackets. Next, we simplify the sums, and replace the summation index  $j$  by

$j + 1$  in the first sum. Consequently, in order, we deduce that

$$\begin{aligned}
S(N; x, q) &= 1 + \sum_{j=1}^{\infty} \frac{(q)_{N-j}(-xq)_{j-1}q^{j(3j-1)/2}x^j}{(q)_j(q)_{N+1-2j}} \{(1 - q^{N+1-j}) + (1 - q^{N+1-2j})xq^{2j}\} \\
&= 1 + \sum_{j=1}^{\infty} \frac{(q)_{N-j}(-xq)_{j-1}q^{j(3j-1)/2}x^j}{(q)_j(q)_{N+1-2j}} \{(1 - q^j) + q^j(1 + xq^j)(1 - q^{N+1-2j})\} \\
&= 1 + \sum_{j=0}^{\infty} \frac{(q)_{N-j-1}(-xq)_j q^{(j+1)(3j+2)/2} x^{j+1}}{(q)_j(q)_{N-1-2j}} \\
&\quad + \sum_{j=1}^{\infty} \frac{(q)_{N-j}(-xq)_j q^{j(3j+1)/2} x^j}{(q)_j(q)_{N-2j}}. \tag{5.1.38}
\end{aligned}$$

If we now let  $N \rightarrow \infty$  in (5.1.38) and assume, for the moment, that (5.1.37) holds, then we deduce Theorem 1.2.25. Thus, (5.1.36) can be considered to be a finite analogue of (1.2.20). We continue with the proof of Theorem 5.1.12.

Extract the term with  $j = 0$  from the first series on the right-hand side of (5.1.38). Then factor out  $(1 + xq)$  from both series that remain. Also, change the order of the two infinite series. Hence,

$$\begin{aligned}
S(N; x, q) &= 1 + xq + (1 + xq) \left\{ \sum_{j=1}^{\infty} \begin{bmatrix} (N-1) + 1 - j \\ j \end{bmatrix} (-xq^2)_{j-1} q^{j(3j-1)/2} (xq)^j \right. \\
&\quad \left. + \sum_{j=1}^{\infty} \begin{bmatrix} (N-1) - j \\ j \end{bmatrix} (-xq^2)_{j-1} q^{j(3j+3)/2} (xq)^{j+1} \right\} \\
&= (1 + xq)S(N-1; xq; q) \\
&= (1 + xq)(1 + xq^2)S(N-2; xq^2; q) \\
&= \dots \\
&= (-xq; q)_N S(0; xq^N, q). \tag{5.1.39}
\end{aligned}$$

Return to the definition of  $S(N; x, q)$  in (5.1.36), and set  $N = 0$ . Recall that  $\begin{bmatrix} n \\ m \end{bmatrix} = 0$ , for  $m > n$ . Thus, we see that in (5.1.36)

$$\begin{bmatrix} 1 - j \\ j \end{bmatrix}, \quad \begin{bmatrix} -j \\ j \end{bmatrix} = 0, \quad j \geq 1.$$

Thus,  $S(0; xq^N, q) = 1$ . Hence, from (5.1.39), we see that we have proved (5.1.37).  $\square$

Let us return to (5.1.38) and let  $N \rightarrow \infty$ . We then obtain the following corollary, which can be thought of as a generalization of the pentagonal number theorem.

**Corollary 5.1.13.** *We have*

$$(-xq; q)_\infty = \sum_{j=0}^{\infty} \frac{(-xq)_j}{(q)_j} q^{(3j^2+j)/2} x^j \{q^{2j+1}x + 1\}.$$

If we set  $x = -1$  in Corollary 5.1.13, we obtain the pentagonal number theorem, Theorem 1.2.26.

We next study, for nonnegative integers  $h, k$ ,

**Definition 5.1.14.** *For nonnegative integers  $h, k$ , and  $N$ , and a real number  $x$ , define*

$$\begin{aligned} C_{k,h}(N; x, q) &= \sum_{j=0}^{\infty} \begin{bmatrix} N+h-kj \\ j \end{bmatrix} (xq)_j (-1)^j x^{kj} q^{(2k+1)j(j+1)/2-hj} \\ &\quad - \sum_{j=0}^{\infty} \begin{bmatrix} N-kj \\ j \end{bmatrix} (xq)_j (-1)^j x^{kj+h} q^{(2k+1)j(j+1)/2+hj+h}. \end{aligned} \quad (5.1.40)$$

In particular, if  $h = k = 1$ , then (5.1.40) reduces to

$$\begin{aligned} C_{1,1}(N; x, q) &= \sum_{j=0}^{\infty} \begin{bmatrix} N+1-j \\ j \end{bmatrix} (xq)_j (-1)^j x^j q^{3j(j+1)/2-j} \\ &\quad - \sum_{j=0}^{\infty} \begin{bmatrix} N-j \\ j \end{bmatrix} (xq)_j (-1)^j x^{j+1} q^{3j(j+1)/2+j+1}. \end{aligned} \quad (5.1.41)$$

Extract the terms with  $j = 0$  in (5.1.41), namely,  $1 - xq$ . In the remaining series (beginning with  $j = 1$ ), factor out  $1 - xq$ . We then find that

$$C_{1,1}(N; x, q) = (1 - xq)S(N; -xq, q) = S(N + 1; -x, q), \quad (5.1.42)$$

by (5.1.36) and the second equality in (5.1.39).

We also note that

$$C_{k,0}(N; x, q) = 0, \quad (5.1.43)$$

since the terms cancel.

**Theorem 5.1.15.** *If  $k, h \geq 1$ , then*

$$\begin{aligned} C_{k,h}(N; x, q) - C_{k,h-1}(N; x, q) \\ = x^{h-1} q^{h-1} (1 - xq) C_{k,k-h+1}(N - (k - h + 1); xq, q). \end{aligned} \quad (5.1.44)$$



**Proof.** Combining the two series and invoking the second  $q$ -analogue of Pascal's formula (Chapter 2, Exercise 5), and noting that the terms with  $j = 0$  cancel, we find that

$$\begin{aligned}
& C_{k,h}(N; x, q) - C_{k,h-1}(N; x, q) \\
&= \sum_{j=0}^{\infty} \left\{ \begin{bmatrix} N+h-kj \\ j \end{bmatrix} - q^j \begin{bmatrix} N+h-1-kj \\ j \end{bmatrix} \right\} (xq)_j (-1)^j x^{kj} q^{(2k+1)j(j+1)/2-hj} \\
&\quad + \sum_{j=0}^{\infty} \begin{bmatrix} N-kj \\ j \end{bmatrix} (1-xq^{j+1})(xq)_j (-1)^j x^{kj+h-1} q^{(2k+1)j(j+1)/2+(h-1)j+h-1} \\
&= \sum_{j=1}^{\infty} \begin{bmatrix} N+h-1-kj \\ j-1 \end{bmatrix} (xq)_j (-1)^j x^{kj} q^{(2k+1)j(j+1)/2-hj} \\
&\quad + \sum_{j=0}^{\infty} \begin{bmatrix} N-kj \\ j \end{bmatrix} (xq)_{j+1} (-1)^j x^{kj+h-1} q^{(2k+1)j(j+1)/2+(h-1)j+h-1} \\
&= \sum_{j=0}^{\infty} \begin{bmatrix} N+h-1-kj-k \\ j \end{bmatrix} (xq)_{j+1} (-1)^{j+1} x^{k(j+1)} q^{(2k+1)(j+1)(j+2)/2-h(j+1)} \\
&\quad + \sum_{j=0}^{\infty} \begin{bmatrix} N-kj \\ j \end{bmatrix} (xq)_{j+1} (-1)^j x^{kj+h-1} q^{(2k+1)j(j+1)/2+(h-1)j+h-1}, \tag{5.1.45}
\end{aligned}$$

where we replaced  $j$  by  $j+1$  in the first sum. Returning to (5.1.45), we switch the order of the two series above, extract common factors from both sums, and refer to (5.1.40) to conclude that

$$\begin{aligned}
& C_{k,h}(N; x, q) - C_{k,h-1}(N; x, q) \\
&= x^{h-1} q^{h-1} (1-xq) \left\{ \sum_{j=0}^{\infty} \begin{bmatrix} N-kj \\ j \end{bmatrix} (xq^2)_j (-1)^j (xq)^{kj} q^{(2k+1)j(j+1)/2-(k-h+1)j} \right. \\
&\quad \left. - \sum_{j=0}^{\infty} \begin{bmatrix} N-(k-h+1)-kj \\ j \end{bmatrix} (xq^2)_j (-1)^j (xq)^{kj+k-h+1} q^{(2k+1)j(j+1)/2+(k-h+1)j+k-h+1} \right\} \\
&= x^{h-1} q^{h-1} (1-xq) C_{k,k-h+1}(N-(k-h+1); xq, q). \tag{5.1.46}
\end{aligned}$$

Equation (5.1.46) is in agreement with (5.1.44), and so the proof is complete.  $\square$

As we shall see, the preceding theorems inspired Andrews to give a new proof of the Rogers–Ramanujan identities [13].

**Theorem 5.1.16.** *We have*

$$G(q) := \sum_{n=0}^{\infty} \frac{q^{n^2}}{(q; q)_n} = \frac{1}{(q; q^5)_{\infty} (q^4; q^5)_{\infty}}, \tag{5.1.47}$$

$$H(q) := \sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q; q)_n} = \frac{1}{(q^2; q^5)_{\infty} (q^3; q^5)_{\infty}}. \tag{5.1.48}$$

These identities were first proved by L. J. Rogers in 1894 [94], but his paper was hardly noticed. Ramanujan rediscovered them in India sometime probably between 1912 and early 1914 before he left for England. However, he did not possess a proof of the identities. He had communicated them in his first letter to G. H. Hardy, but unfortunately the page of the letter containing the identities has been lost. Published with Ramanujan's lost notebook [92] is a short manuscript that Ramanujan wrote containing four reasons why Ramanujan believed that the identities were true [17, Chapter 10]. After his arrival in Cambridge, the identities became famous, and their combinatorial interpretations were first found by P. A. MacMahon, who in his epic two volumes [77] presented them as open problems.

**Corollary 5.1.17.** (a) *The number of partitions of a positive integer  $n$  into parts differing by at least 2 is equal to the number of partitions of  $n$  into parts congruent to either 1 or 4 modulo 5.*

(b) *The number of partitions of a positive integer  $n$  into parts differing by at least 2 and with no 1's is equal to the number of partitions of  $n$  into parts congruent to either 2 or 3 modulo 5.*

**Proof.** We first prove (a). It is clear that the right-hand side of (5.1.47) generates all partitions into parts congruent to either 1 or 4 modulo 5. On the left-hand side, consider the Ferrers graph of a partition  $\pi$  of an integer  $N$  arranged into less than or equal to  $n$  decreasing parts. We now recall that  $1 + 3 + \cdots + (2n - 1) = n^2$ ,  $1 \leq n < \infty$ . Now add to the Ferrers graph of  $\pi$   $2n - 1$  nodes in the first row,  $2n - 3$  nodes in the second row, etc. and 1 node in the  $n$ th row, irrespective of whether the partition  $\pi$  actually has an  $n$ th part. We thus obtain a partition into exactly  $n$  distinct parts, and the difference between any two parts is at least equal to 2.

Consider next part (b). The combinatorial interpretation of the right-hand side of (5.1.48) should be clear. As in the proof of (a), consider the Ferrers graph of a partition  $\pi$  with less than or equal to  $n$  parts arranged in descending order. Recall that  $2 + 4 + \cdots + 2n = n(n + 1)$ . Adjoin the parts  $2n, 2n - 2, \dots, 2$  in descending order to the Ferrers graph of  $\pi$ . We then obtain a partition into exactly  $n$  parts, each differing from any other part by at least 2. However, the last part is at least equal to 2, because we added 2 to the last part of  $\pi$ . Hence, this establishes the combinatorial interpretation of the second Rogers–Ramanujan identity.  $\square$

**Example 5.1.18.** *To illustrate the first Rogers–Ramanujan identity, consider the partitions of 8 into the two types:*

$$8 = 7 + 1 = 6 + 2 = 5 + 3,$$

$$6 + 1 + 1 = 4 + 4 = 4 + 1 + 1 + 1 + 1 = 1 + 1 + 1 + 1 + 1 + 1 + 1.$$

*To illustrate the second identity, again consider 8 with the following partitions:*

$$8 = 6 + 2 = 5 + 3,$$

$$8 = 3 + 3 + 2 = 2 + 2 + 2 + 2.$$

One day during his stay at Cambridge, Ramanujan was perusing old issues of the *Proceedings of the London Mathematical Society*, and he saw that Rogers [94] had proved the identities that he had conjectured. Shortly thereafter, Ramanujan found his own proof. Rogers was contacted, and he himself found another proof, whereupon Hardy arranged for the two proofs to be published together [86]. It was first observed by R. A. Askey that, ironically, in India, Ramanujan had actually established a theorem from which the Rogers–Ramanujan identities could be deduced as corollaries. That identity, Entry 7 in Chapter 16 of Ramanujan’s second notebook [91], [27, p. 16], is given by

$$\sum_{n=0}^{\infty} \frac{(a)_n (d/b)_n (d/c)_n (d/q)_n (1-dq^{2n-1})(bc/a)^n}{(b)_n (c)_n (d/a)_n (q)_n (1-d/q)} q^{n(n-1)} = \frac{(a)_{\infty} (d)_{\infty}}{(b)_{\infty} (c)_{\infty}} \sum_{n=0}^{\infty} \frac{(b/a)_n (c/a)_n}{(d/a)_n (q)_n} a^n,$$

and the aforementioned proof by Askey is found in [27, pp. 77–78]. One can also find a short history of the Rogers–Ramanujan identities in [27, pp. 77–79]. A much more complete and informative survey of proofs of the Rogers–Ramanujan identities has been written by Andrews [14].

It is now time to give Andrews’s proof of the Rogers–Ramanujan identities [13] stimulated by the work of Sylvester.

**Proof.** Let  $h = k = 2$  in (5.1.44) in Theorem 5.1.15 to deduce that

$$C_{2,2}(N; x, q) - C_{2,1}(N; x, q) = xq(1-xq)C_{2,1}(N-1; xq, q). \quad (5.1.49)$$

Next, set  $h = 1$  and  $k = 2$  in (5.1.44) and recall (5.1.43). Thus,

$$C_{2,1}(N; x, q) = (1-xq)C_{2,2}(N-2; xq, q). \quad (5.1.50)$$

Eliminate  $C_{2,1}(N; x, q)$  from (5.1.49) and (5.1.50) to arrive at

$$C_{2,2}(N; x, q) = (1-xq)C_{2,2}(N-2; xq, q) + xq(1-xq)(1-xq^2)C_{2,2}(N-3; xq^2, q). \quad (5.1.51)$$

Define

$$D(N; x, q) := \sum_{0 \leq 2j \leq N} \begin{bmatrix} N-j \\ j \end{bmatrix} x^j q^{j^2} = \sum_{j=0}^{\infty} \begin{bmatrix} N-j \\ j \end{bmatrix} x^j q^{j^2}. \quad (5.1.52)$$

Note that if we set  $x = 1$  and  $x = q$  in (5.1.52), and let  $N \rightarrow \infty$ , we obtain, respectively, the Rogers–Ramanujan functions  $G(q)$  and  $H(q)$  in (5.1.47) and (5.1.48), respectively.

Next, define

$$\Delta(N; x, q) = C_{2,2}(N; x, q) - (xq)_{[N/2]+1} D\left(\left[\frac{N}{2}\right] + 2; x, q\right). \quad (5.1.53)$$

We now compute some polynomials related to  $C_{2,2}(i; a, q)$ ,  $D\left(\left[\frac{i}{2}\right] + 2; a, q\right)$ , and  $\Delta(i; a, q)$ .

On the basis of Table 2, we make the following conjecture, which we prove.

$i$	$C_{2,2}(i; a, q)$	$(aq)_{[i/2]+1} D\left(\left[\frac{i}{2}\right] + 2; a, q\right)$	$\Delta(i; a, q)$
0	$1 - a^2q^2$	$(1 - aq)(1 + aq)$	0
1	$1 - a^2q^2 - a^2q^3 + a^3q^4$	$(1 - aq)(1 + aq)$	$-a^2q^3 + a^3q^4$
2	$1 - a^2q^2 - a^2q^3 - a^2q^4$ $+ a^3q^4 + a^3q^5$	$(1 - aq)(1 - aq^2)$ $\times (1 + aq + aq^2)$	0
3	$1 - a^2q^2 - a^2q^3 - a^2q^4$ $- a^2q^5 + a^3q^4 + a^3q^5$ $+ a^3q^6 + a^4q^9 - a^5q^{10}$	$(1 - aq)(1 - aq^2)$ $\times (1 + aq + aq^2)$	$-a^2q^5 + a^3q^6$ $+ a^4q^9 - a^5q^{10}$

**Table 2.** Table of Polynomials

**Theorem 5.1.19.** For each nonnegative integer  $N$ ,

$$x^{-2}q^{-2N-1}\Delta(2N-1; x, q) \quad \text{and} \quad x^{-3}q^{-2N-4}\Delta(2N; x, q) \quad (5.1.54)$$

are polynomials in  $x$  and  $q$ .

**Proof.** It is clear from Table 2 that Theorem 5.1.19 is true for  $N = 0, 1, 2, 3$ . We use induction. Recall that  $\begin{bmatrix} N-1 \\ -1 \end{bmatrix} = 0$ . Using the second  $q$ -analogue of Pascal's formula from Exercise 5 in Chapter 2, we write

$$D(N; x, q) = \sum_{j=0}^{\infty} \left( \begin{bmatrix} N-j-1 \\ j-1 \end{bmatrix} + q^j \begin{bmatrix} N-j-1 \\ j \end{bmatrix} \right) x^j q^{j^2}.$$

We now switch the order of the two sums above and, noting that the term with  $j = 0$  equals 0 in what was formerly the first sum, replace  $j$  by  $j + 1$  in what was formerly the first sum to find that

$$\begin{aligned} D(N; x, q) &= D(N-1; xq, q) + \sum_{j=0}^{\infty} \begin{bmatrix} N-j-2 \\ j \end{bmatrix} x^{j+1} q^{(j+1)^2} \\ &= D(N-1; xq, q) + xqD(N-2; xq^2, q). \end{aligned} \quad (5.1.55)$$

Next, after using the distributive law, we observe that the terms with  $j = 0$  cancel in the first sum below. We appeal to the first  $q$ -analogue of Pascal's formula in Exercise 5 in Chapter 2. Thus,

$$\begin{aligned} &(1 - xq^N)D(N+1; x, q) - D(N; x, q) \\ &= \sum_{j=0}^{\infty} \left( \begin{bmatrix} N+1-j \\ j \end{bmatrix} x^j q^{j^2} - \begin{bmatrix} N-j \\ j \end{bmatrix} x^j q^{j^2} \right) \end{aligned}$$

$$\begin{aligned}
& -xq^N \sum_{j=0}^{\infty} \begin{bmatrix} N+1-j \\ j \end{bmatrix} x^j q^{j^2} \\
&= \sum_{j=1}^{\infty} \begin{bmatrix} N-j \\ j-1 \end{bmatrix} x^j q^{N+(j-1)^2} - xq^N \sum_{j=0}^{\infty} \begin{bmatrix} N+1-j \\ j \end{bmatrix} x^j q^{j^2} \\
&= \sum_{j=0}^{\infty} \begin{bmatrix} N-j-1 \\ j \end{bmatrix} x^{j+1} q^{N+j^2} - xq^N \sum_{j=0}^{\infty} \begin{bmatrix} N+1-j \\ j \end{bmatrix} x^j q^{j^2} \\
&= xq^N \sum_{j=0}^{\infty} \left( \begin{bmatrix} N-j-1 \\ j \end{bmatrix} - \begin{bmatrix} N+1-j \\ j \end{bmatrix} \right) x^j q^{j^2}. \tag{5.1.56}
\end{aligned}$$

We now apply the first of the two  $q$ -analogues of Pascal's formula from Exercise 5 of Chapter 2 to each of the  $q$ -binomial coefficients on the far right side of (5.1.56). Accordingly,

$$\begin{aligned}
\begin{bmatrix} N-j-1 \\ j \end{bmatrix} - \begin{bmatrix} N+1-j \\ j \end{bmatrix} &= \begin{bmatrix} N-j \\ j \end{bmatrix} - q^{N-2j} \begin{bmatrix} N-j-1 \\ j-1 \end{bmatrix} \\
&\quad - \left( \begin{bmatrix} N-j \\ j \end{bmatrix} + q^{N-2j+1} \begin{bmatrix} N-j \\ j-1 \end{bmatrix} \right) \\
&= -q^{N-2j+1} \begin{bmatrix} N-j \\ j-1 \end{bmatrix} - q^{N-2j} \begin{bmatrix} N-j-1 \\ j-1 \end{bmatrix}.
\end{aligned}$$

Substituting the foregoing calculation into (5.1.56), we deduce that

$$\begin{aligned}
& (1-xq^N)D(N+1; x, q) - D(N; x, q) \\
&= -xq^N \sum_{j=0}^{\infty} \left( q^{N-2j+1} \begin{bmatrix} N-j \\ j-1 \end{bmatrix} + q^{N-2j} \begin{bmatrix} N-j-1 \\ j-1 \end{bmatrix} \right) x^j q^{j^2}.
\end{aligned}$$

The term with  $j=0$  is equal to 0, and so we replace  $j$  by  $j+1$  above and recall the definition of  $D(N; x, q)$  in (5.1.52) to deduce that

$$\begin{aligned}
& (1-xq^N)D(N+1; x, q) - D(N; x, q) \\
&= -xq^{2N} \sum_{j=0}^{\infty} \left( q^{-2j-1} \begin{bmatrix} N-1-j \\ j \end{bmatrix} + q^{-2j-2} \begin{bmatrix} N-2-j \\ j \end{bmatrix} \right) x^{j+1} q^{j^2+2j+1} \\
&= -x^2 q^{2N} D(N-1; x, q) - x^2 q^{2N-1} D(N-2; x, q). \tag{5.1.57}
\end{aligned}$$

Assume now that Theorem 5.1.19 is valid for each subscript  $< 2N$ . Return to the definition of  $D$  in (5.1.53) and use (5.1.51) and (5.1.55) to deduce that

$$\begin{aligned}
\Delta(2N; x, q) &= C_{2,2}(2N; x, q) - (xq)_{N+1} D(N+2; x, q) \\
&= (1-xq)C_{2,2}(2N-2; xq, q) + xq(1-xq)(1-xq^2)C_{2,2}(2N-3; xq^2, q) \\
&\quad - (xq)_{N+1} \{ D(N+1; xq, q) + xqD(N; xq^2, q) \} \\
&= (1-xq)\Delta(2(N-1); xq, q) + xq(1-xq)(1-xq^2)\Delta(2(N-1)-1; xq^2, q). \tag{5.1.58}
\end{aligned}$$

Multiplying both sides of (5.1.58) by  $x^{-3}q^{-2N-4}$ , we arrive at

$$\begin{aligned} & x^{-3}q^{-2N-4}\Delta(2N; x, q) \\ &= q(1-xq)(xq)^{-3}q^{-2(N-1)-4}\Delta(2(N-1); xq, q) \\ & \quad + (1-xq)(1-xq^2)(xq^2)^{-2}q^{-2(N-1)-1}\Delta(2(N-1)-1; xq^2, q). \end{aligned} \quad (5.1.59)$$

Applying the induction hypothesis, we complete the proof in the case that the index is even.

We next consider the case when the index is odd and assume that the theorem is valid for all smaller indices. We once again use (5.1.51) and (5.1.55). However, in this case, we must add and subtract  $xq(xq)_{N+2}D(N+1; xq^2, q)$  in order to bring us to  $\Delta(2N-2; xq^2, q)$ . Hence,

$$\begin{aligned} \Delta(2N+1; x, q) &= C_{2,2}(2N+1; x, q) - (xq)_{N+1}D(N+2; x, q) \\ &= (1-xq)C_{2,2}(2N-1; xq, q) + xq(1-xq)(1-xq^2)C_{2,2}(2N-2; xq^2, q) \\ & \quad - (xq)_{N+1} \{D(N+1; xq, q) + xqD(N; xq^2, q)\} \\ &= (1-xq)\Delta(2N-1; xq, q) + xq(1-xq)(1-xq^2)\Delta(2N-2; xq^2, q) \\ & \quad + xq(xq)_{N+2}D(N+1; xq^2, q) - xq(xq)_{N+1}D(N; xq^2, q) \\ &= (1-xq)\Delta(2N-1; xq, q) + xq(1-xq)(1-xq^2)\Delta(2N-2; xq^2, q) \\ & \quad + xq(xq)_{N+1} \{-x^2q^4q^{2N}D(N-1; xq^2, q) - x^2q^4q^{2N-1}D(N-2; xq^2, q)\}, \end{aligned} \quad (5.1.60)$$

where we have employed (5.1.57). Multiply both sides of (5.1.60) by  $x^{-2}q^{-2N-3}$  to see that

$$\begin{aligned} & x^{-2}q^{-2N-3}\Delta(2N+1; x, q) \\ &= (1-xq)(xq)^{-2}q^{-2N-1}\Delta(2N-1; xq, q) \\ & \quad + x^2q^6(1-xq)(1-xq^2)(xq^2)^{-3}q^{-2N-2}\Delta(2N-2; xq^2, q) \\ & \quad - xq^2(xq)_{N+1}D(N-1; xq^2, q) - xq(xq)_{N+1}D(N-2; xq^2, q). \end{aligned} \quad (5.1.61)$$

By induction and inspection, we see that each expression on the right-hand side of (5.1.61) is a polynomial in  $x$  and  $q$ . Thus, we have completed the proof for odd index. Hence, with our conclusions from (5.1.59) and (5.1.61), we have completed the proof of Theorem 5.1.19.  $\square$

We now complete Andrews's proof of the Rogers–Ramanujan identities.

Let us fix  $r$  and  $n$  and examine the coefficients of  $x^r q^n$  in  $D(N; x, q)$  and  $C_{2,2}(N; x, q)$ . If  $N$  is sufficiently large, say  $N \geq N_0(r, n)$ , then these coefficients of  $x^r q^n$  are constant. To see this, we observe that as  $N$  increases, the new terms in  $q$  have powers larger than  $n$ .

Let us examine carefully  $x^{-2}q^{-2N-1}\Delta(2N-1; x, q)$ , and put  $m = 2N-1$ . From Theorem 5.1.19, we see that if  $-2N-1+n < 0$ , or equivalently,  $m \geq n-1$ , then this constant coefficient of  $q^n$  in  $\Delta(2N-1; x, q)$  must be equal to 0. Suppose next that

$m = 2N$ . Then, by Theorem 5.1.19, if  $-2N - 4 + n < 0$ , or equivalently,  $m \geq n - 3$ , this constant coefficient of  $q^n$  in  $\Delta(2N; x, q)$  must also be equal to 0. In general, if  $m \geq n - 1$ , the coefficient of  $q^n$  in

$$C_{2,2}(m; x, q) - (xq)_{[m/2]+1} D(\lfloor \frac{m}{2} \rfloor + 2; x, q)$$

must be equal to 0. In particular, if we let  $m \rightarrow \infty$ , we must conclude that the coefficient of  $q^n$  in

$$C_{2,2}(\infty; x, q) - (xq)_\infty D(\infty; x, q)$$

is equal to 0. In other words,

$$C_{2,2}(\infty; x, q) - (xq)_\infty D(\infty; x, q) \equiv 0. \quad (5.1.62)$$

Returning to the definition (5.1.52), we see that

$$\begin{aligned} \lim_{N \rightarrow \infty} D(N; 1, q) &= \lim_{N \rightarrow \infty} \sum_{0 \leq 2j \leq N} \begin{bmatrix} N - j \\ j \end{bmatrix} q^{j^2} \\ &= \sum_{j=0}^{\infty} \frac{q^{j^2}}{(q; q)_j}, \end{aligned} \quad (5.1.63)$$

where we have used the notation from (5.1.47). Next, from the definition of  $C_{2,2}$  in (5.1.40),

$$\begin{aligned} \lim_{N \rightarrow \infty} C_{2,2}(N; 1, q) &= \lim_{N \rightarrow \infty} \left( \sum_{j=0}^{\infty} \begin{bmatrix} N + 2 - 2j \\ j \end{bmatrix} (q)_j (-1)^j q^{5j(j+1)/2 - 2j} \right. \\ &\quad \left. - \sum_{j=0}^{\infty} \begin{bmatrix} N - 2j \\ j \end{bmatrix} (q)_j (-1)^j q^{5j(j+1)/2 + 2j + 2} \right) \\ &= \sum_{j=0}^{\infty} (-1)^j q^{j(5j+1)/2} - \sum_{j=0}^{\infty} (-1)^j q^{j(5j+9)/2 + 2} \\ &= \sum_{j=0}^{\infty} (-1)^j q^{j(5j+1)/2} + \sum_{n=-1}^{-\infty} (-1)^n q^{n(5n+1/2)} \\ &= \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j+1)/2} \\ &= f(-q^3, -q^2) \\ &= (q^3; q^5)(q^2; q^5)(q^5; q^5) \\ &= \frac{(q; q)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty}, \end{aligned} \quad (5.1.64)$$

where in the anti-penultimate line we set  $j = -n - 1$ , and lastly invoked the Jacobi triple product identity (3.1.19).

Now put (5.1.63) and (5.1.64) in (5.1.62), with  $x = 1$ , to finally deduce that

$$\frac{(q; q)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty} = (q; q)_\infty \sum_{j=0}^{\infty} \frac{q^{j^2}}{(q; q)_j},$$

which proves the first Rogers–Ramanujan identity (5.1.47).

To prove the second identity, return to (5.1.50) and let  $N \rightarrow \infty$ , and then use (5.1.62). Hence,

$$\begin{aligned} C_{2,1}(\infty; x, q) &= (1 - xq)C_{2,2}(\infty; xq, q) \\ &= (1 - xq)(xq^2)_\infty D(\infty; xq, q) \\ &= (xq)_\infty D(\infty; xq, q). \end{aligned} \tag{5.1.65}$$

If  $x = 1$ , then (5.1.65) reduces to

$$C_{2,1}(\infty; 1, q) = (q)_\infty D(\infty; q, q) = (q)_\infty \sum_{j=0}^{\infty} \frac{q^{j(j+1)}}{(q; q)_j}. \tag{5.1.66}$$

Now, setting  $k = 2$  and  $h = 1$  in (5.1.40), and replacing  $j$  by  $-j - 1$  in the second sum in the third equality below, we find that

$$\begin{aligned} \lim_{N \rightarrow \infty} C_{2,1}(N; 1, q) &= \lim_{N \rightarrow \infty} \left( \sum_{j=0}^{\infty} \begin{bmatrix} N+1-2j \\ j \end{bmatrix} (q)_j (-1)^j q^{5j(j+1)/2-j} \right. \\ &\quad \left. - \sum_{j=0}^{\infty} \begin{bmatrix} N-2j \\ j \end{bmatrix} (q)_j (-1)^j q^{5j(j+1)/2+j+1} \right) \\ &= \sum_{j=0}^{\infty} (-1)^j q^{5j(j+1)/2-j} - \sum_{j=0}^{\infty} (-1)^j q^{5j(j+1)/2+j+1} \\ &= \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j+3)/2} \\ &= f(-q^4, -q) \\ &= (q; q^5)_\infty (q^4; q^5)_\infty (q^5; q^5)_\infty = \frac{(q; q)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty}, \end{aligned} \tag{5.1.67}$$

upon the invoking of the Jacobi triple product identity (3.1.19). The second Rogers–Ramanujan identity (5.1.47) follows immediately from (5.1.66) and (5.1.67).  $\square$

The ideas of Sylvester and Andrews have not been as fully exploited as they should be. In an email to the lecturer on February 9, 2014, Andrews wrote, “I do think that there is a lot left to do.” Other papers by Andrews relevant to his proof and that of Sylvester are [9], [15], and [16].



## 5.2. The Rogers–Ramanujan Continued Fraction

A continued fraction is an expression of the sort

$$b_0 + \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \frac{a_4}{b_4 + \cdots}}}}, \quad (5.2.1)$$

which is commonly written in the more compact form

$$b_0 + \frac{a_1}{b_1} + \frac{a_2}{b_2} + \frac{a_3}{b_3} + \frac{a_4}{b_4} + \cdots. \quad (5.2.2)$$

Suppose that we define the sequences  $P_n$  and  $Q_n$ ,  $n \geq -1$ , by

$$\begin{aligned} P_n &= b_n P_{n-1} + a_n P_{n-2}, & n \geq 1, \\ Q_n &= b_n Q_{n-1} + a_n Q_{n-2}, & n \geq 1, \\ P_{-1} &= 1, \quad Q_{-1} = 0, \quad P_0 = b_0, \quad Q_0 = 1. \end{aligned}$$

One can readily check that

$$\frac{P_n}{Q_n} := b_0 + \frac{a_1}{b_1} + \frac{a_2}{b_2} + \cdots + \frac{a_n}{b_n}. \quad (5.2.3)$$

Then if

$$\lim_{n \rightarrow \infty} \frac{P_n}{Q_n}$$

exists, we say that the continued fraction (5.2.2) converges; otherwise it diverges. For theorems providing criteria for the convergence or divergence of (5.2.2), we refer readers to the excellent text by L. Lorentzen and H. Waadeland [76, Chapter 1]. In particular, see [76, p. 35, Theorem 3].

Most mathematics students first encounter continued fractions in a course in elementary number theory. The first infinite continued fractions that students may be asked to evaluate are those in Exercise 6 below. These are, in fact, special cases of perhaps the most interesting continued fraction in mathematics, the Rogers–Ramanujan continued fraction, which first appeared in a paper by L. J. Rogers [94] in 1894.

**Definition 5.2.1.** *The Rogers–Ramanujan continued fraction  $R(q)$  is defined by*

$$R(q) := \frac{q^{1/5}}{1} + \frac{q}{1} + \frac{q^2}{1} + \frac{q^3}{1} + \cdots, \quad (5.2.4)$$

*provided that it converges. Furthermore, set*

$$T(q) := 1 + \frac{q}{1} + \frac{q^2}{1} + \frac{q^3}{1} + \cdots. \quad (5.2.5)$$

By Exercise 6, we see that  $R(\pm 1)$  converges. In general,  $R(q)$  converges for  $|q| < 1$ , but for  $|q| = 1$ , the problem of convergence is open. However, at roots of unity on the unit circle, by a theorem of I. Schur [96, pp. 319–321] and Ramanujan [91], [28, p. 35], we do know the behavior of  $R(q)$ .

**Theorem 5.2.2.** *Recall that  $T(q)$  is defined by (5.2.5). Let  $q$  be a primitive  $m$ th root of unity. If  $m$  is a multiple of 5,  $T(q)$  diverges. Otherwise,  $T(q)$  converges and*

$$T(q) = \alpha T(\alpha) q^{(1-\alpha\rho m)/5}, \quad (5.2.6)$$

where  $\alpha$  denotes the Legendre symbol  $(\frac{m}{5})$  and  $\rho$  is the least positive residue of  $m$  modulo 5.

In our completion of this chapter, our primary aim is to establish the connection between the Rogers–Ramanujan functions  $G(q)$  and  $H(q)$  and the Rogers–Ramanujan continued fraction  $R(q)$ . More precisely, Rogers [94] and Ramanujan [91, Vol. II, Chapter 16, Sect. 15], [27, p. 30] proved that

$$R(q) = q^{1/5} \frac{H(q)}{G(q)}. \quad (5.2.7)$$

In fact, we first prove a *finite form* of (5.2.7), which can be found as Entry 16 in Chapter 16 of Ramanujan’s second notebook [91], [27, p. 31], and from which (5.2.7) follows as an immediate corollary.

**Theorem 5.2.3.** *For each nonnegative integer  $n$ , let*

$$\mu := \mu_n(a, q) := \sum_{k=0}^{[(n+1)/2]} \frac{(q)_{n-k+1} a^k q^{k^2}}{(q)_k (q)_{n-2k+1}}, \quad (5.2.8)$$

$$\nu := \nu_n(a, q) := \sum_{k=0}^{[n/2]} \frac{(q)_{n-k} a^k q^{k(k+1)}}{(q)_k (q)_{n-2k}}, \quad (5.2.9)$$

where  $[x]$  denotes the greatest integer less than or equal to  $x$ . Then, for  $n \geq 1$ ,

$$\frac{\mu}{\nu} = 1 + \frac{aq}{1} + \frac{aq^2}{1} + \cdots + \frac{aq^n}{1}. \quad (5.2.10)$$

**Proof.** For each nonnegative integer  $r$ , define

$$F_r := F_r(a, q) := \sum_{k=0}^{[(n-r+1)/2]} \frac{(q)_{n-r-k+1} a^k q^{k(r+k)}}{(q)_k (q)_{n-r-2k+1}}.$$

Observe that

$$F_0 = \mu \quad \text{and} \quad F_1 = \nu. \quad (5.2.11)$$

Also note that

$$F_n = 1 \quad \text{and} \quad F_{n-1} = 1 + aq^n. \quad (5.2.12)$$

We now develop a recurrence relation for  $F_r$ . When we combine the two sums in the first step below, we conventionally set  $1/(q)_{-1} = 0$ . To that end,

$$\begin{aligned}
F_r - F_{r+1} &= \sum_{k=0}^{[(n-r+1)/2]} \frac{(q)_{n-r-k+1} a^k q^{k(r+k)}}{(q)_k (q)_{n-r-2k+1}} - \sum_{k=0}^{[(n-r)/2]} \frac{(q)_{n-r-k} a^k q^{k(r+1+k)}}{(q)_k (q)_{n-r-2k}} \\
&= \sum_{k=1}^{[(n-r+1)/2]} \frac{(q)_{n-r-k} a^k q^{k(r+k)}}{(q)_k (q)_{n-r-2k}} \left( \frac{1 - q^{n-r-k+1}}{1 - q^{n-r-2k+1}} - q^k \right) \\
&= \sum_{k=1}^{[(n-r+1)/2]} \frac{(q)_{n-r-k} a^k q^{k(r+k)}}{(q)_k (q)_{n-r-2k}} \frac{(1 - q^k)}{(1 - q^{n-r-2k+1})} \\
&= \sum_{k=1}^{[(n-r+1)/2]} \frac{(q)_{n-r-k} a^k q^{k(r+k)}}{(q)_{k-1} (q)_{n-r-2k+1}} \\
&= \sum_{j=0}^{[(n-r-1)/2]} \frac{(q)_{n-r-j-1} a^{j+1} q^{(j+1)(r+j+1)}}{(q)_j (q)_{n-r-2j-1}} \\
&= aq^{r+1} \sum_{j=0}^{[(n-r-1)/2]} \frac{(q)_{n-(r+2)-j+1} a^j q^{j(r+2+j)}}{(q)_j (q)_{n-(r+2)-2j+1}} \\
&= aq^{r+1} F_{r+2}. \tag{5.2.13}
\end{aligned}$$

Using (5.2.11), (5.2.13) repeatedly, and lastly (5.2.12), we conclude that

$$\begin{aligned}
\frac{\mu}{\nu} &= \frac{F_0}{F_1} = \frac{F_1 + aqF_2}{F_1} = 1 + \frac{aq}{F_1/F_2} \\
&= 1 + \frac{aq}{(F_2 + aq^2F_3)/F_2} = 1 + \frac{aq}{1} + \frac{aq^2}{F_2/F_3} \\
&= 1 + \frac{aq}{1} + \frac{aq^2}{1} + \cdots + \frac{aq^{n-1}}{F_{n-1}/F_n} \\
&= 1 + \frac{aq}{1} + \frac{aq^2}{1} + \cdots + \frac{aq^{n-1}}{1} + \frac{aq^n}{1}.
\end{aligned}$$

□

**Corollary 5.2.4.** For any complex number  $a$  and  $|q| < 1$ ,

$$\frac{\sum_{k=0}^{\infty} \frac{a^k q^{k^2}}{(q)_k}}{\sum_{k=0}^{\infty} \frac{a^k q^{k(k+1)}}{(q)_k}} = 1 + \frac{aq}{1} + \frac{aq^2}{1} + \cdots + \frac{aq^n}{1} + \cdots. \tag{5.2.14}$$

**Proof.** Let  $n \rightarrow \infty$  in (5.2.10). □

The continued fraction in (5.2.14) is called the *Generalized Rogers–Ramanujan Continued Fraction*.

Recall that the Rogers–Ramanujan identities are given by

$$G(q) = \frac{1}{(q; q^5)_\infty (q^4; q^5)_\infty} \quad \text{and} \quad H(q) = \frac{1}{(q^2; q^5)_\infty (q^3; q^5)_\infty}. \quad (5.2.15)$$

Using (5.2.15), we immediately deduce the elegant representation for  $R(q)$  in the next theorem.

**Theorem 5.2.5.** *We have*

$$R(q) = q^{1/5} \frac{(q; q^5)_\infty (q^4; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty}. \quad (5.2.16)$$

**Proof.** Set  $a = 1$  in Corollary 5.2.4, take the reciprocal of both sides, use the definitions of  $G(q)$  and  $H(q)$  in (5.2.15), and lastly use (5.1.1).  $\square$

We now establish a very important formula for the Rogers–Ramanujan continued fraction that has had many applications, including the explicit evaluation of the Rogers–Ramanujan continued fraction at certain arguments.

**Theorem 5.2.6.** *If  $T(q)$  is defined in (5.2.5). Then*

$$T(q^5) - q - \frac{q^2}{T(q^5)} = \frac{(q; q)_\infty}{(q^{25}; q^{25})_\infty}. \quad (5.2.17)$$

We now prove a more general theorem from which Theorem 5.2.6 follows by specialization. We employ the notation (1.1.3).

**Theorem 5.2.7.** *For any complex number  $a$ ,*

$$\begin{aligned} & (a, a^2, q/a, q/a^2, q; q)_\infty \\ &= (q^5; q^5)_\infty \left( \frac{(a^5 q; q^5)_\infty (a^{-5} q^4; q^5)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty} - a \frac{(a^5 q^2; q^5)_\infty (a^{-5} q^3; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty} \right. \\ & \quad \left. - a^2 \frac{(a^5 q^3; q^5)_\infty (a^{-5} q^2; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty} + a^3 \frac{(a^5 q^4; q^5)_\infty (a^{-5} q; q^5)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty} \right). \end{aligned} \quad (5.2.18)$$

Before proving Theorem 5.2.7, we show that Theorem 5.2.6 follows immediately from Theorem 5.2.7.

**Proof of Theorem 5.2.6.** If we replace  $q$  by  $q^5$  in (5.2.18), then set  $a = q$ , realize that  $(1; q^{25})_\infty = 0$ , and recall (5.2.16), we deduce (5.2.17) forthwith.  $\square$

The decompositions in Theorems 5.2.6 and 5.2.7 are called *5-dissections*, because in the former theorem, the series terms of  $(q; q)_\infty$  are separated out in powers of  $q$  according to their residue classes modulo 5, and in the latter theorem, the terms are separated out in powers of  $a$  according to their residue classes modulo 5.

**Proof of Theorem 5.2.7.** Using the Jacobi triple product identity (1.1.7) twice, we find that

$$(a, a^2, q/a, q/a^2, q; q)_\infty = \frac{(a, q/a, q; q)_\infty (a^2, q/a^2, q; q)_\infty}{(q; q)_\infty}$$

$$\begin{aligned}
&= \frac{1}{(q; q)_\infty} \sum_{r=-\infty}^{\infty} (-1)^r a^r q^{(r^2-r)/2} \sum_{s=-\infty}^{\infty} (-1)^s a^{2s} q^{(s^2-s)/2} \\
&= \frac{1}{(q; q)_\infty} \sum_{r,s=-\infty}^{\infty} (-1)^{r+s} a^{r+2s} q^{(r^2-r+s^2-s)/2} \\
&= \sum_{n=-\infty}^{\infty} a^n c_n(q), \tag{5.2.19}
\end{aligned}$$

where, for  $-\infty < n < \infty$ ,

$$c_n(q) := \frac{1}{(q; q)_\infty} \sum_{\substack{r,s=-\infty \\ r+2s=n}}^{\infty} (-1)^{r+s} q^{(r^2-r+s^2-s)/2}.$$

We now determine  $c_n(q)$  according to the residue class of  $n$  modulo 5.

First, consider the residue class  $0 \pmod{5}$ . Replace  $n$  by  $5n$  and make the change of variables  $r = n - 2t$  and  $s = 2n + t$ . Note that  $r + 2s = 5n$ . Then, simplifying and applying the Jacobi triple product identity (1.1.7), we find that

$$\begin{aligned}
c_{5n}(q) &= \frac{1}{(q; q)_\infty} \sum_{t=-\infty}^{\infty} (-1)^{n+t} q^{((n-2t)^2 - (n-2t) + (2n+t)^2 - (2n+t))/2} \\
&= \frac{(-1)^n q^{(5n^2-3n)/2}}{(q; q)_\infty} \sum_{t=-\infty}^{\infty} (-1)^t q^{(5t^2+t)/2} \\
&= \frac{(-1)^n q^{(5n^2-3n)/2}}{(q; q)_\infty} f(-q^3, -q^2) \\
&= \frac{(-1)^n q^{(5n^2-3n)/2}}{(q; q^5)_\infty (q^4; q^5)_\infty}. \tag{5.2.20}
\end{aligned}$$

Second, consider the residue class  $1 \pmod{5}$ . Set  $r = n - 2t + 1$  and  $s = 2n + t$ , so that  $r + 2s = 5n + 1$ . Then, upon simplification and the use of the Jacobi triple product identity (1.1.7), we see that

$$\begin{aligned}
c_{5n+1}(q) &= \frac{1}{(q; q)_\infty} \sum_{t=-\infty}^{\infty} (-1)^{n+t+1} q^{((n-2t+1)^2 - (n-2t+1) + (2n+t)^2 - (2n+t))/2} \\
&= \frac{(-1)^{n+1} q^{(5n^2-n)/2}}{(q; q)_\infty} \sum_{t=-\infty}^{\infty} (-1)^t q^{(5t^2-3t)/2} \\
&= \frac{(-1)^{n+1} q^{(5n^2-n)/2}}{(q; q)_\infty} f(-q, -q^4) \\
&= \frac{(-1)^{n+1} q^{(5n^2-n)/2}}{(q^2; q^5)_\infty (q^3; q^5)_\infty}. \tag{5.2.21}
\end{aligned}$$

It should now be clear how to calculate the three remaining cases,

$$c_{5n+2}(q) = \frac{(-1)^{n+1} q^{(5n^2+n)/2}}{(q^2; q^5)_\infty (q^3; q^5)_\infty}, \quad (5.2.22)$$

$$c_{5n+3}(q) = \frac{(-1)^n q^{(5n^2+3n)/2}}{(q; q^5)_\infty (q^4; q^5)_\infty}, \quad (5.2.23)$$

$$c_{5n+4}(q) = 0. \quad (5.2.24)$$

Substitute (5.2.20)–(5.2.24) in (5.2.19) and use the Jacobi triple product identity (1.1.7) four times to conclude that

$$\begin{aligned} & (a, a^2, q/a, q/a^2, q; q)_\infty \\ &= \frac{1}{(q; q^5)_\infty (q^4; q^5)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n a^{5n} q^{(5n^2-3n)/2} \\ & \quad - \frac{1}{(q^2; q^5)_\infty (q^3; q^5)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n a^{5n+1} q^{(5n^2-n)/2} \\ & \quad - \frac{1}{(q^2; q^5)_\infty (q^3; q^5)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n a^{5n+2} q^{(5n^2+n)/2} \\ & \quad + \frac{1}{(q; q^5)_\infty (q^4; q^5)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n a^{5n+3} q^{(5n^2+3n)/2} \\ &= (q^5; q^5)_\infty \left( \frac{(a^5 q; q^5)_\infty (a^{-5} q^4; q^5)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty} - a \frac{(a^5 q^2; q^5)_\infty (a^{-5} q^3; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty} \right. \\ & \quad \left. - a^2 \frac{(a^5 q^3; q^5)_\infty (a^{-5} q^2; q^5)_\infty}{(q^2; q^5)_\infty (q^3; q^5)_\infty} + a^3 \frac{(a^5 q^4; q^5)_\infty (a^{-5} q; q^5)_\infty}{(q; q^5)_\infty (q^4; q^5)_\infty} \right). \end{aligned}$$

Thus, the proof of Theorem 5.2.7 is complete.  $\square$

We now employ Theorem 5.2.6 to prove another identity involving  $T(q^5)$ . We then will use this identity to prove two congruences for  $p(n)$ , in anticipation of our more general approach to congruences for  $p(n)$  in the following chapter.

**Theorem 5.2.8.** *We have*

$$T^5(q^5) - 11q^5 - \frac{q^{10}}{T^5(q^5)} = \frac{(q^5; q^5)_\infty^6}{(q^{25}; q^{25})_\infty^6}. \quad (5.2.25)$$

**Proof.** Return to (5.2.17) and replace  $q$  everywhere by  $\omega q$ , where  $\omega$  is any fifth root of unity. Then take the product of both sides of the equation over all five fifth roots of unity. Hence,

$$\prod_{\omega} \frac{(\omega q; \omega q)_\infty}{(q^{25}; q^{25})_\infty} = \prod_{\omega} \left( T(q^5) - \omega q - \frac{\omega^2 q^2}{T(q^5)} \right). \quad (5.2.26)$$

On the left-hand side of (5.2.26), if  $5 \nmid n$ , we obtain products of the form

$$(1 - q^n)(1 - \omega q^n)(1 - \omega^2 q^n)(1 - \omega^3 q^n)(1 - \omega^4 q^n) = (1 - q^{5n}). \quad (5.2.27)$$

If  $5|n$  and  $n = 5m$ , then on the left side of (5.2.26), we have products of the form

$$(1 - q^n)(1 - q^n)(1 - q^n)(1 - q^n)(1 - q^n) = (1 - q^n)^5 = (1 - q^{5m})^5, \quad (5.2.28)$$

for each positive integer  $m$ . Hence, by (5.2.27) and (5.2.28),

$$\prod_{\omega} \frac{(\omega q; \omega q)_{\infty}}{(q^{25}; q^{25})_{\infty}} = \frac{1}{(q^{25}; q^{25})_5} \prod_{n=1}^{\infty} (1 - q^{5n}) \prod_{m=1}^{\infty} (1 - q^{5m})^5 = \frac{(q^5; q^5)_{\infty}^6}{(q^{25}; q^{25})_{\infty}^6}. \quad (5.2.29)$$

Since the left side of (5.2.26) is real, then the right side of (5.2.26) must be real as well. If we consider a symmetric sum of five terms containing products of  $k$  roots of unity,  $1 \leq k \leq 4$ , then the contribution of such a sum will be equal to 0, simply because the sum of 5 distinct fifth roots of unity equals 0. Thus, the only possible symmetric sums of five terms that may yield non-zero contributions will be a sum in which each term has a product of five fifth roots of unity. Hence,

$$\prod_{\omega} \left( T(q^5) - \omega q - \frac{\omega^2 q^2}{T(q^5)} \right) = T^5(q^5) - \frac{q^{10}}{T^5(q^5)} - q^5 + C_1 q^5 + C_2 q^5, \quad (5.2.30)$$

where  $C_1$  and  $C_2$  are certain constants.

First,  $C_1$  denotes the sum of  $\binom{5}{1,3,1}$  terms of the form

$$C_1 = \sum T(q^5) \cdot \omega_1 \omega_2 \omega_3 \cdot \frac{\omega_4^2}{T(q^5)} = \sum \omega_1 \omega_2 \omega_3 \omega_4^2 \quad (5.2.31)$$

Let us fix  $\omega_5$  and sum on  $\omega_j$ ,  $1 \leq j \leq 4$ . Since  $\omega_5$  can be any of five fifth roots of unity, by symmetry, clearly, by (5.2.31),  $C_1$  will be equal to 5 times this sum. Hence,

$$C_1 = 5 \sum \omega_1 \omega_2 \omega_3 \omega_4^2 = 5 \sum_{\omega_4 \neq \omega_5} \omega_4 \omega_5^{-1} = -5 \omega_5 \omega_5^{-1} = -5. \quad (5.2.32)$$

Second,  $C_2$  denotes the sum of  $\binom{5}{2,1,2}$  terms of the form

$$C_2 = - \sum T^2(q^5) \cdot \omega_1 \cdot \frac{(\omega_2 \omega_3)^2}{T^2(q^5)} = - \sum \omega_1 \omega_2^2 \omega_3^2. \quad (5.2.33)$$

Let us determine that partial subset of  $C_2$  in which we sum over  $\omega$  different from  $\omega_2$  and  $\omega_3$ , i.e.,

$$\begin{aligned} - \sum_{\omega \neq \omega_2, \omega_3} \omega \omega_2^2 \omega_3^2 &= -(\omega_1 + \omega_4 + \omega_5) \omega_2^2 \omega_3^2 \\ &= (\omega_2 + \omega_3) \omega_2^2 \omega_3^2 = \omega_2^3 \omega_3^2 + \omega_2^2 \omega_3^3. \end{aligned} \quad (5.2.34)$$

Next, we sum the expressions above over all possible  $\omega_2$  in (5.2.34). Hence,

$$\sum_{\omega_2 \neq \omega_3} (\omega_2^3 \omega_3^2 + \omega_2^2 \omega_3^3) = -\omega_3^2 \omega_3^2 - \omega_3^2 \omega_3^3 = -2 \quad (5.2.35)$$

Now let us summarize our calculations. Note that in (5.2.35), because of the symmetry of  $\omega_2$  and  $\omega_3$ , we have counted each contribution twice. Also, there are five possibilities

for  $\omega_1$ . Hence, from (5.2.33),

$$C_2 = 5 \cdot \frac{1}{2}(-2) = -5. \quad (5.2.36)$$

In summary, using the values of  $C_1$  and  $C_2$  from (5.2.32) and (5.2.36), respectively, in (5.2.30), we find that

$$\prod_{\omega} \left( T(q^5) - \omega q - \frac{\omega^2 q^2}{T(q^5)} \right) = T^5(q^5) - \frac{q^{10}}{T^5(q^5)} - q^5 - 5q^5 - 5q^5 = T^5(q^5) - \frac{q^{10}}{T^5(q^5)} - 11q^5,$$

which, combined with (5.2.29), completes the proof.  $\square$

We now demonstrate how Theorem 5.2.8 can be utilized to prove the following congruence for  $p(n)$ .

**Theorem 5.2.9.** *For each nonnegative integer  $n$ ,*

$$p(5n + 4) \equiv 0 \pmod{5}. \quad (5.2.37)$$

**Proof.** We first write (5.2.25) in the form

$$1 = \frac{(q^{25}; q^{25})_{\infty}^6}{(q^5; q^5)_{\infty}^6} \left( T^5(q^5) - 11q^5 - \frac{q^{10}}{T^5(q^5)} \right). \quad (5.2.38)$$

Secondly, we write (5.2.17) in the form

$$(q; q)_{\infty} = (q^{25}; q^{25})_{\infty} \left( T(q^5) - q - \frac{q^2}{T(q^5)} \right). \quad (5.2.39)$$

For brevity, set  $x = T(q^5)$ . Divide (5.2.38) by (5.2.39) to deduce that

$$\begin{aligned} \frac{1}{(q; q)_{\infty}} &= \frac{(q^{25}; q^{25})_{\infty}^5}{(q^5; q^5)_{\infty}^6} \frac{x^5 - 11q^5 - q^{10}/x^5}{x - q - q^2/x} \\ &= \frac{(q^{25}; q^{25})_{\infty}^5}{(q^5; q^5)_{\infty}^6} \left( x^4 + qx^3 + 2q^2x^2 + 3q^3x + 5q^4 - 3\frac{q^5}{x} + 2\frac{q^6}{x^2} - \frac{q^7}{x^3} + \frac{q^8}{x^4} \right). \end{aligned} \quad (5.2.40)$$

Now equate all terms with powers congruent to 4 modulo 5 to arrive at

$$\sum_{n=0}^{\infty} p(5n + 4)q^{5n+4} = 5q^4 \frac{(q^{25}; q^{25})_{\infty}^5}{(q^5; q^5)_{\infty}^6}. \quad (5.2.41)$$

The congruence (5.2.37) is now obvious.  $\square$

We can utilize (5.2.41) to derive a congruence for  $p(n)$  modulo 25.

**Theorem 5.2.10.** *For each nonnegative integer  $n$ ,*

$$p(25n + 24) \equiv 0 \pmod{25}. \quad (5.2.42)$$



**Proof.** Return to (5.2.41), divide both sides by  $q^4$ , and replace  $q^5$  by  $q$  to obtain the alternative form

$$\sum_{n=0}^{\infty} p(5n+4)q^n = 5 \frac{(q^5; q^5)_{\infty}^5}{(q; q)_{\infty}^6}. \quad (5.2.43)$$

By the binomial theorem,

$$(q; q)_{\infty}^5 \equiv (q^5; q^5)_{\infty} \pmod{5},$$

and so, from (5.2.43),

$$\begin{aligned} \sum_{n=0}^{\infty} p(5n+4)q^n &\equiv 5 \frac{(q^5; q^5)_{\infty}^4}{(q; q)_{\infty}} \pmod{5} \\ &\equiv (q^5; q^5)_{\infty}^4 \sum_{n=0}^{\infty} p(n)q^n \pmod{5}. \end{aligned} \quad (5.2.44)$$

By Theorem 5.2.9, the coefficients of  $q^{5n+4}$  on the right side of (5.2.44) are multiples of 5. Hence, the coefficients of  $q^{5n+4}$  on the left side of (5.2.44) are multiples of 5, i.e., the coefficients

$$p(5(5n+4)+4) = p(25n+24) \equiv 0 \pmod{5}.$$

This concludes the proof of Theorem 5.2.10.  $\square$

We conclude this chapter with one further application of Theorem 5.2.9. Define the Ramanujan  $\tau$ -function by

$$q(q; q)_{\infty}^{24} =: \sum_{n=1}^{\infty} \tau(n)q^n. \quad (5.2.45)$$

**Theorem 5.2.11.** *For each positive integer  $n$ ,*

$$\tau(5n) \equiv 0 \pmod{5}. \quad (5.2.46)$$

**Proof.** Using the definition (5.2.45) and the congruence (5.2.37), we find that

$$\begin{aligned} \sum_{n=1}^{\infty} \tau(n)q^n &= q(q; q)_{\infty}^{24} \\ &= q \frac{(q; q)_{\infty}^{25}}{(q; q)_{\infty}} \equiv q(q^5; q^5)_{\infty}^5 \sum_{n=0}^{\infty} p(n)q^n \pmod{5}. \end{aligned} \quad (5.2.47)$$

Equating coefficients of  $q^{5n}$  on both sides of (5.2.47), we find that  $\tau(5n)$  is a linear combination of terms of the form  $p(5m+4)$ . Hence, by Theorem 5.2.9,  $\tau(5n)$  is a multiple of 5. This concludes the proof of (5.2.46).  $\square$

The results in this section can be greatly generalized in that many more general continued fractions can be represented as quotients of two  $q$ -series. See, in particular, [17, Chapter 6] and [27, pp. 30–31].

**5.3. Exercises**

1. Prove combinatorially that  $S_2(n) = S_3(n)$ .
2. Use Lebesgue's Identity to evaluate  $g_0(q)$ .
3. Find an evaluation of (5.1.29) that is simpler than using Watson's  $q$ -analogue of Whipple's theorem.
4. Give a bijective proof of Corollary 5.1.13.
5. Prove (5.2.3).

6. Prove that

$$1 + \frac{1}{1} + \frac{1}{1} + \frac{1}{1} + \dots = \frac{\sqrt{5} + 1}{2} \quad (5.3.1)$$

and

$$1 - \frac{1}{1} + \frac{1}{1} - \frac{1}{1} + \dots = \frac{\sqrt{5} - 1}{2}. \quad (5.3.2)$$

7. Prove (5.2.22)–(5.2.24).

## The Frobenius and Generalized Frobenius Symbols

### 6.1. First Generalization of Frobenius Partitions

Recall that in Chapter 3 we defined the Frobenius symbol in Definition 3.1.10.

**Definition 6.1.1.** Consider the Ferrers graph of a positive integer  $n$ . Let  $r$  be the size of a Durfee square. Form the diagonal of the Durfee square, which will have  $r$  nodes. To the right of the diagonal is a graphical representation of a partition of no more than  $r$  parts, reading from top to bottom, say  $a_1, a_2, \dots, a_r$ . To the left of the diagonal is a graphical representation of another partition of no more than  $r$  parts, reading from left to right, namely  $b_1, b_2, \dots, b_r$ , say. Thus,  $n = r + \sum_{j=1}^r (a_j + b_j)$ . A matrix representation corresponding to these two partitions can be given by the Frobenius symbol

$$\begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}.$$

From the constructive definition that we gave, it is clear that  $a_1 > a_2 > \dots > a_r$  and that  $b_1 > b_2 > \dots > b_r$ . Note that it possible that  $a_r$  or  $b_r$  could be equal to 0. Recall that the number of such Frobenius symbols is  $p(n)$ . We will examine a couple generalizations.

Suppose that we allow repetition in any row at most  $k$  times. We keep the requirement that  $n = r + \sum_{j=1}^r (a_j + b_j)$ .

**Example 6.1.2.** Let  $k = 2$  and  $n = 3$ . Then the possible generalized Frobenius symbols are

$$\begin{pmatrix} 2 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$$

with the respective values of  $r = 1, 1, 1, 2, 2$ .

**Definition 6.1.3.** *If we allow  $k$  repetitions in the Frobenius symbol, then we let  $\varphi_k(n)$  denote the number of these Generalized Frobenius Symbols.*

Clearly,  $\varphi_1(n) = p(n)$ . Let

$$\Phi_k(q) := \sum_{n=0}^{\infty} \varphi_k(n) q^n. \quad (6.1.1)$$

At the beginning of Chapter 1, we showed that  $\Phi_1(q) = 1/(q; q)_{\infty}$ . We also showed in Chapter 3 that  $\Phi_1(q)$  was the constant term in the product

$$\prod_{n=1}^{\infty} (1 + zq^n)(1 + q^{n-1}/z) = (-zq; q)_{\infty}(-1/z; q)_{\infty}.$$

By the same reasoning, we see that  $\Phi_k(q)$  is the constant term

$$\begin{aligned} c_0 &:= [z^0] \prod_{n=1}^{\infty} (1 + zq^n + \cdots + z^k q^{kn})(1 + z^{-1}q^{n-1} + \cdots + z^{-k}q^{k(n-1)}) \\ &= [z^0] \prod_{n=1}^{\infty} \frac{(1 - z^{k+1}q^{(k+1)n})}{(1 - zq^n)} \frac{(1 - z^{-(k+1)}q^{(k+1)(n-1)})}{(1 - z^{-1}q^{n-1})}. \end{aligned} \quad (6.1.2)$$

In the next theorem, we prove that  $c_0$  can be written as a product of theta functions. However, even with this representation, it remains very difficult to determine  $\phi_k(n)$ . With a little effort, we shall explicitly determine  $\phi_2(n)$ .

**Theorem 6.1.4.** *Recall that  $f(a, b)$  is defined in (1.1.6). Let  $\zeta = \exp(2\pi i/(k+1))$ . Then*

$$c_0 = [z^0] \frac{1}{(q; q)_{\infty}^k} \prod_{j=1}^k f(\zeta^j zq, \zeta^{-j} z^{-1}). \quad (6.1.3)$$

**Proof.** First observe that

$$\prod_{j=0}^k (1 - \zeta^j x) = 1 - x^{k+1}. \quad (6.1.4)$$

Using (6.1.4) twice in (6.1.2), replacing  $j$  by  $k+1-j$  in the second product in the third equality below, and using the Jacobi triple product identity (3.1.18), we find that

$$\begin{aligned} c_0 &= [z^0] \prod_{n=1}^{\infty} \frac{1}{(1 - zq^n)(1 - z^{-1}q^{n-1})} \prod_{j=0}^k (1 - \zeta^j zq^n)(1 - \zeta^j z^{-1}q^{n-1}) \\ &= [z^0] \prod_{n=1}^{\infty} \prod_{j=1}^k (1 - \zeta^j zq^n)(1 - \zeta^j z^{-1}q^{n-1}) \\ &= [z^0] \prod_{n=1}^{\infty} \prod_{j=1}^k (1 - \zeta^j zq^n)(1 - \zeta^{-j} z^{-1}q^{n-1}) \\ &= [z^0] \frac{1}{(q; q)_{\infty}^k} \prod_{j=1}^k \prod_{n=1}^{\infty} (1 - \zeta^j zq^n)(1 - \zeta^{-j} z^{-1}q^{n-1})(1 - q^n) \end{aligned}$$

$$= [z^0] \frac{1}{(q; q)_\infty^k} \prod_{j=1}^k f(\zeta^j z q, \zeta^{-j} z^{-1}),$$

and so the proof of (6.1.3) is complete.  $\square$

Let  $k = 1$  in Theorem 6.1.4. Then  $\Phi_1(q)$  is the constant term in

$$\begin{aligned} \Phi_1(q) &= [z^0] \frac{1}{(q; q)_\infty} \sum_{n_1=0}^{\infty} (-1)^{n_1} (-1)^{n_1} z^{n_1} q^{n_1(n_1+1)/2} \\ &= [z^0] \frac{1}{(q; q)_\infty} \sum_{n_1=0}^{\infty} z^{n_1} q^{n_1(n_1+1)/2} \\ &= \frac{1}{(q; q)_\infty} \cdot 1 = \frac{1}{(q; q)_\infty}, \end{aligned}$$

in agreement with our previous observation in Chapters 1 and 3.

**Theorem 6.1.5.** *We have*

$$\Phi_2(q) = \frac{(q^6; q^{12})_\infty}{(q; q)_\infty (q^2; q^4)_\infty (q^3; q^6)_\infty}. \quad (6.1.5)$$

**Proof.** Let  $\zeta = e^{2\pi i/3}$ . If  $n_1$  and  $n_2$  are the two indices of summation for the theta functions in (6.1.3), we see that for the constant term,  $n_1 = -n_2$ . Hence, using the Jacobi product identity below (3.1.18), we find that

$$\begin{aligned} \Phi_2(q) &= \frac{1}{(q; q)_\infty^2} \sum_{n_2=-\infty}^{\infty} \zeta^{-1 \cdot n_2 + 2n_2} q^{-n_2(-n_2+1)/2 + n_2(n_2+1)/2} \\ &= \frac{1}{(q; q)_\infty^2} \sum_{n=-\infty}^{\infty} \zeta^n q^{n^2} \\ &= \frac{1}{(q; q)_\infty^2} f(\zeta q, \zeta^{-1} q) \\ &= \frac{1}{(q; q)_\infty^2} (-\zeta q; q^2)_\infty (-\zeta^{-1} q; q^2)_\infty (q^2; q^2)_\infty \\ &= \frac{(q^2; q^2)_\infty}{(q; q)_\infty^2} \prod_{n=1}^{\infty} (1 + (\zeta + \zeta^{-1})q^{2n-1} + q^{4n-2}) \\ &= \frac{(q^2; q^2)_\infty}{(q; q)_\infty^2} \prod_{n=1}^{\infty} (1 - q^{2n-1} + q^{4n-2}) \\ &= \frac{(q^2; q^2)_\infty}{(q; q)_\infty^2} \prod_{n=1}^{\infty} \frac{1 + q^{6n-3}}{1 + q^{2n-1}} \\ &= \frac{(q^2; q^2)_\infty (-q^3; q^6)_\infty}{(q; q)_\infty^2 (-q; q^2)_\infty} \\ &= \frac{(q^2; q^2)_\infty (q^6; q^{12})_\infty}{(q; q)_\infty^2 (-q; q^2)_\infty (q^3; q^6)_\infty} \end{aligned}$$

$$\begin{aligned}
&= \frac{(q^6; q^{12})_\infty}{(q; q)_\infty (q; q^2)_\infty (-q; q^2)_\infty (q^3; q^6)_\infty} \\
&= \frac{(q^6; q^{12})_\infty}{(q; q)_\infty (q^2; q^4)_\infty (q^3; q^6)_\infty}.
\end{aligned}$$

Thus, obtaining (6.1.5), we complete the proof.  $\square$

## 6.2. Second Generalization of Frobenius Partitions

We now consider  $k$  copies of the nonnegative integers with a total ordering as follows:

$$0_1 < 0_2 < \cdots < 0_k < 1_1 < 1_2 < \cdots < 1_k < 2_1 < 2_2 < \cdots. \quad (6.2.1)$$

**Definition 6.2.1.** *The generalized Frobenius symbol, associated with the natural number  $n$ , is given by*

$$\begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ b_1 & b_2 & \cdots & b_r \end{pmatrix},$$

where the elements are arranged in strictly decreasing order according to the convention (6.2.1). Moreover,

$$n = r + \sum_{j=1}^r (a_j + b_j).$$

Let  $c\varphi_k(n)$  denote the number of such partitions of  $n$ , and let

$$c\Phi_k(q) = \sum_{n=0}^{\infty} c\varphi_k(n)q^n \quad (6.2.2)$$

denote the generating function of  $c\varphi_k(n)$ . (The appendage  $c$  is an indication that we can regard these partitions of  $n$  as partitions in  $k$  colors.)

**Example 6.2.2.** *Let  $k = 2$  and  $n = 2$ . The possible generalized partitions of  $n$  are:*

$$\begin{pmatrix} 1_2 \\ 0_2 \end{pmatrix}, \begin{pmatrix} 1_2 \\ 0_1 \end{pmatrix}, \begin{pmatrix} 1_1 \\ 0_2 \end{pmatrix}, \begin{pmatrix} 1_1 \\ 0_1 \end{pmatrix}, \begin{pmatrix} 0_2 \\ 1_2 \end{pmatrix}, \begin{pmatrix} 0_1 \\ 1_2 \end{pmatrix}, \begin{pmatrix} 0_2 \\ 1_1 \end{pmatrix}, \begin{pmatrix} 0_1 \\ 1_1 \end{pmatrix}, \begin{pmatrix} 0_2 & 0_1 \\ 0_2 & 0_1 \end{pmatrix}.$$

Therefore,  $c\varphi_2(2) = 9$ .

Analogously to how we argued in (3.1.42) and Definition 3.1.10,  $c\Phi_k(q)$  is the constant term in

$$\prod_{n=1}^{\infty} (1 + zq^n)^k (1 + z^{-1}q^{n-1})^k. \quad (6.2.3)$$

Using the same kind of argument that we used above to deduce (6.1.3), except that now  $\zeta = 1$  and  $z$  is replaced by  $-z$ , and recalling the notation (2.2.3), we find that

$$\begin{aligned} c\Phi_k(q) &= [z^0] \frac{1}{(q; q)_\infty^k} \sum_{n_1, n_2, \dots, n_k=0}^{\infty} q^{n_1(n_1+1)/2+n_2(n_2+1)/2+\dots+n_k(n_k+1)/2} z_1^{n_1} z_1^{n_2} \dots z_1^{n_k} \\ &= \frac{1}{(q; q)_\infty^k} \sum_{\substack{n_1, n_2, \dots, n_k=0 \\ n_1+n_2+\dots+n_k=0}}^{\infty} q^{n_1(n_1+1)/2+n_2(n_2+1)/2+\dots+n_k(n_k+1)/2}. \end{aligned} \quad (6.2.4)$$

In particular,

$$c\Phi_1(q) = \frac{1}{(q; q)_\infty}$$

and, by the Jacobi triple product identity (3.1.18),

$$c\Phi_2(q) = \frac{1}{(q; q)_\infty^2} \sum_{n_1=-\infty}^{\infty} q^{n_1^2} = \frac{(-q; q^2)_\infty^2 (q^2; q^2)_\infty}{(q; q)_\infty^2}. \quad (6.2.5)$$

We close our discussion of  $c\varphi_k(n)$  by deriving a simple congruence for  $c\varphi_k(n)$  modulo any prime  $k = p$ . From (6.2.4) and (6.2.3),

$$\begin{aligned} \sum_{n=0}^{\infty} c\varphi_k(n) q^n &= [z^0] (-zq; q)_\infty^k (-z^{-1}; q)_\infty^k \\ &\equiv [z^0] (-z^k q^k; q^k)_\infty^k (-z^{-k}; q^k)_\infty^k \pmod{k} \\ &\equiv [z^0] \frac{1}{(q^k; q^k)_\infty^k} \sum_{n=0}^{\infty} z^{kn} q^{kn(n+1)/2} \pmod{k} \\ &= \frac{1}{(q^k; q^k)_\infty^k} \pmod{k} \\ &= \sum_{m=0}^{\infty} p(m) q^{km} \pmod{k}. \end{aligned}$$

Equating coefficients of  $q^n$  above, we deduce the following theorem.

**Theorem 6.2.3.** *We have for every pair of natural numbers  $k, n$ ,*

$$c\varphi_k(n) \equiv \begin{cases} 0 \pmod{k}, & \text{if } k \nmid n, \\ p(n/k) \pmod{k}, & \text{if } k \mid n. \end{cases}$$





---

## Chapter 7

# Congruences for $p(n)$

### 7.1. Introduction

In about 1916, P. A. MacMahon used the most common recurrence formula for  $p(n)$  to calculate  $p(n)$  for the first 200 values of  $n$ . He conveniently arranged the values in segmented columns with five values in each grouping. Ramanujan noticed that every fifth value was divisible by 5, i.e.,  $p(5n + 4) \equiv 0 \pmod{5}$ ,  $0 \leq n \leq 39$ . He also noticed further congruences modulo 7 and 11. More precisely, in 1919, Ramanujan [87], [90, pp. 210–213] announced that he had found three simple congruences satisfied by  $p(n)$ , namely,

$$p(5n + 4) \equiv 0 \pmod{5}, \tag{7.1.1}$$

$$p(7n + 5) \equiv 0 \pmod{7}, \tag{7.1.2}$$

$$p(11n + 6) \equiv 0 \pmod{11}. \tag{7.1.3}$$

He gave proofs of (7.1.1) and (7.1.2) in [87] and later in a short one page note [88], [90, p. 230] announced that he had also found a proof of (7.1.3). He also remarks in [88] that “It appears that there are no equally simple properties for any moduli involving primes other than these three.” It was not until 2003 that this speculative observation was proved by S. Ahlgren and M. Boylan [3]. In a posthumously published paper [89], [90, pp. 232–238], G. H. Hardy extracted different proofs of (7.1.1)–(7.1.3) from an unpublished manuscript of Ramanujan on  $p(n)$  and  $\tau(n)$  [92, pp. 133–177], [36], [19, Chapter 5].

In [87], Ramanujan offered a more general conjecture. Let  $\delta = 5^a 7^b 11^c$  and let  $\lambda$  be an integer such that  $24\lambda \equiv 1 \pmod{\delta}$ . Then

$$p(n\delta + \lambda) \equiv 0 \pmod{\delta}. \tag{7.1.4}$$

In more detail, Ramanujan conjectured:

$$\text{If } 24N \equiv 1 \pmod{5^n}, \quad \text{then } p(N) \equiv 0 \pmod{5^n}; \quad (7.1.5)$$

$$\text{if } 24N \equiv 1 \pmod{7^n}, \quad \text{then } p(N) \equiv 0 \pmod{7^n}; \quad (7.1.6)$$

$$\text{if } 24N \equiv 1 \pmod{11^n}, \quad \text{then } p(N) \equiv 0 \pmod{11^n}. \quad (7.1.7)$$

In his unpublished manuscript [92, pp. 133–177], [36], Ramanujan gave a proof of (7.1.4) for arbitrary  $a$  and  $b = c = 0$ . He also began a proof of his conjecture for arbitrary  $b$  and  $a = c = 0$ , but he did not complete it. If he had completed his proof, he would have noticed that his conjecture in this case needed to be modified. However, on the other hand, it is possible that while trying to construct a proof, he found that his conjecture was false, and so he abandoned his attempt at finding a proof. Recall that Ramanujan had formulated his conjectures after studying MacMahon's table of values of  $p(n)$ ,  $0 \leq n \leq 200$ . After Ramanujan died, H. Gupta [64], [65, pp. 47–53] extended MacMahon's table up to  $n = 300$ . Upon examining Gupta's table in 1934, S. Chowla [47] found that  $p(243) = 133978259344888$  is not divisible by  $7^3$ , despite the fact that  $24 \cdot 243 \equiv 1 \pmod{7^3}$ . To correct Ramanujan's conjecture, define  $\delta' = 5^a 7^{b'} 11^c$ , where  $b' = b$ , if  $b = 0, 1, 2$ , and  $b' = \lfloor (b+2)/2 \rfloor$ , if  $b > 2$ . Then

$$p(n\delta + \lambda) \equiv 0 \pmod{\delta'}. \quad (7.1.8)$$

In particular,

$$\text{If } 24N \equiv 1 \pmod{7^n}, \quad \text{then } p(N) \equiv 0 \pmod{7^{\lfloor (n+2)/2 \rfloor}}.$$

In 1938, G. N. Watson [100] published a proof of (7.1.8) for  $a = c = 0$  and gave a more detailed version of Ramanujan's proof of (7.1.8) in the case  $b = c = 0$ . It was not until 1967 that A. O. L. Atkin [24] proved (7.1.8) for arbitrary  $c$  and  $a = b = 0$ . M. D. Hirschhorn and D. C. Hunt [71] constructed a proof of (7.1.8) for arbitrary powers of 5, while F. Garvan [55] devised a proof of (7.1.8) for general powers of 7, both in the spirit of Ramanujan's proof. An account of the two general congruences for powers of 5 and 7 in the spirit of modular forms can be found in M. I. Knopp's excellent book [74].

## 7.2. Ramanujan's Congruence

$$p(5n + 4) \equiv 0 \pmod{5}$$

We shall give several proofs of Ramanujan's congruence for  $p(n)$  modulo 5. We offer three proofs in this section. The first and the second are more elementary than the third, but the third gives more information.

**Theorem 7.2.1.** *For each nonnegative integer  $n$ ,*

$$p(5n + 4) \equiv 0 \pmod{5}. \quad (7.2.1)$$

**First Proof of Theorem 7.2.1.** Our first proof is taken from Ramanujan's paper [87], [90, pp. 210–213] and is reproduced in Hardy's book [67, pp. 87–88].

We begin by writing

$$q(q; q)_\infty^4 \frac{(q^5; q^5)_\infty}{(q; q)_\infty^5} = q \frac{(q^5; q^5)_\infty}{(q; q)_\infty} = (q^5; q^5)_\infty \sum_{m=0}^{\infty} p(m)q^{m+1}. \quad (7.2.2)$$

By the binomial theorem,

$$(q; q)_\infty^5 \equiv (q^5; q^5)_\infty \pmod{5} \quad \text{or} \quad \frac{(q^5; q^5)_\infty}{(q; q)_\infty^5} \equiv 1 \pmod{5}. \quad (7.2.3)$$

Hence, by (7.2.2) and (7.2.3),

$$q(q; q)_\infty^4 \equiv (q^5; q^5)_\infty \sum_{m=0}^{\infty} p(m)q^{m+1} \pmod{5}. \quad (7.2.4)$$

We now see from (7.2.4) that in order to show that  $p(5n+4) \equiv 0 \pmod{5}$  we must show that the coefficients of  $q^{5n+5}$  on the left side of (7.2.4) are multiples of 5.

By the pentagonal number theorem, Corollary 1.2.26, and Jacobi's identity (3.1.31),

$$\begin{aligned} q(q; q)_\infty^4 &= q(q; q)_\infty (q; q)_\infty^3 \\ &= q \sum_{j=-\infty}^{\infty} (-1)^j q^{j(3j+1)/2} \sum_{k=0}^{\infty} (-1)^k (2k+1) q^{k(k+1)/2} \\ &= \sum_{j=-\infty}^{\infty} \sum_{k=0}^{\infty} (-1)^{j+k} (2k+1) q^{1+j(3j+1)/2+k(k+1)/2}. \end{aligned} \quad (7.2.5)$$

Our objective is to determine when the exponents on the right side are multiples of 5. Observe that

$$2(j+1)^2 + (2k+1)^2 = 8 \left\{ 1 + \frac{1}{2}j(3j+1) + \frac{1}{2}k(k+1) \right\} - 10j^2 - 5.$$

Thus,  $1 + \frac{1}{2}j(3j+1) + \frac{1}{2}k(k+1)$  is a multiple of 5 if and only if

$$2(j+1)^2 + (2k+1)^2 \equiv 0 \pmod{5}. \quad (7.2.6)$$

It is easily checked that  $2(j+1)^2 \equiv 0, 2$ , or  $3$  modulo 5 and that  $(2k+1)^2 \equiv 0, 1$ , or  $4$  modulo 5. We therefore see that (7.2.6) is true if and only if

$$2(j+1)^2 \equiv 0 \pmod{5} \quad \text{and} \quad (2k+1)^2 \equiv 0 \pmod{5}.$$

In particular,  $2k+1 \equiv 0 \pmod{5}$ , which, by (7.2.5), implies that the coefficient of  $q^{5n+5}$ ,  $n \geq 0$ , in  $q(q; q)_\infty^4$  is a multiple of 5. The coefficient of  $q^{5n+5}$  on the right side of (7.2.4) is therefore also a multiple of 5, i.e.,  $p(5n+4)$  is a multiple of 5.  $\square$

We next give a variant of Ramanujan's proof due to Mike Hirschhorn [70].

**Second Proof of Theorem 7.2.1.** Recall again Jacobi's Identity (3.1.31)

$$J := \sum_{k=0}^{\infty} (-1)^k (2k+1) q^{k(k+1)/2} = (q; q)_\infty^3. \quad (7.2.7)$$

If on the left side of (7.2.7) we collect together terms according to the residue classes of the powers of  $q$  modulo 5, we find that

$$J \equiv J_0 + J_1 \pmod{5}, \quad (7.2.8)$$

where  $J_j$  contains all of the terms in which the power of  $q$  is congruent to  $j$  modulo 5. On the other hand, by the generating function for  $p(n)$ , the binomial theorem, and (7.2.8),

$$\begin{aligned} \sum_{n=0}^{\infty} p(n)q^n &= \frac{1}{(q; q)_{\infty}} = \frac{(q; q)_{\infty}^9}{(q; q)_{\infty}^{10}} \\ &= \frac{J^3}{((q; q)_{\infty}^5)^2} \equiv \frac{(J_0 + J_1)^3}{(q^5; q^5)_{\infty}^2} \\ &= \frac{J_0^3 + 3J_0^2J_1 + 3J_0J_1^2 + J_1^3}{(q^5; q^5)_{\infty}^2} \pmod{5}. \end{aligned}$$

A careful inspection of the terms on the right-hand side above shows that none of the powers are congruent to 4 modulo 5. Thus,  $p(5n + 4) \equiv 0 \pmod{5}$ .  $\square$

We now give a third simple proof due to G. E. Andrews [12] and based on the simple lemma given below. See also an extensive generalization of this lemma by Andrews and R. Roy [23]. In particular, taking a special case of their general theorem, Andrews and Roy establish the congruence  $p(7n + 5) \equiv 0 \pmod{7}$ .

**Lemma 7.2.2.** *Let  $\{a_n\}$ ,  $n \geq 0$ , be any sequence of integers. Then the coefficient of  $q^{5n+3}$ ,  $n \geq 0$ , in*

$$L(q) := \frac{1}{(q; q)_{\infty}^2} \sum_{n=0}^{\infty} a_n q^{n^2} \quad (7.2.9)$$

is divisible by 5.

**Proof.** Write (7.2.9) in the form

$$L(q) = (q; q)_{\infty}^3 \frac{1}{(q; q)_{\infty}^5} \sum_{m=0}^{\infty} a_m q^{m^2} \equiv (q; q)_{\infty}^3 \frac{1}{(q^5; q^5)_{\infty}} \sum_{m=0}^{\infty} a_m q^{m^2} \pmod{5},$$

by the binomial theorem. Using Jacobi's identity (3.1.31), we thus see that it suffices to examine the coefficient of  $q^{5n+3}$  in

$$(q; q)_{\infty}^3 \sum_{m=0}^{\infty} a_m q^{m^2} = \sum_{j=0}^{\infty} (-1)^j (2j+1) q^{j(j+1)/2} \sum_{m=0}^{\infty} a_m q^{m^2}. \quad (7.2.10)$$

We want those terms above for which  $j(j+1)/2 + m^2 = 5n + 3$ , where  $n \geq 0$ . It is easy to see that this condition is equivalent to the congruence

$$(2j+1)^2 + 3m^2 \equiv 0 \pmod{5}. \quad (7.2.11)$$

Since  $(2j+1)^2 \equiv 0, \pm 1 \pmod{5}$  and  $3m^2 \equiv 0, 2, 3 \pmod{5}$ , we see that (7.2.11) holds only when

$$m \equiv 2j + 1 \equiv 0 \pmod{5}. \quad (7.2.12)$$

The coefficients of  $q^{5n+3}$  in (7.2.10) are then composed of terms of the sort  $(-1)^j(2j+1)a_m$ , which, by (7.2.12), are all multiples of 5.  $\square$

**Third Proof of Theorem 7.2.1.** Using (7.6.1), we find that

$$\begin{aligned} \sum_{k=0}^{\infty} p(k)q^{2k} &= \frac{1}{(q^2; q^2)_{\infty}} = \frac{1}{(q; q)_{\infty}(-q; q)_{\infty}} = \frac{1}{(q; q)_{\infty}^2} \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \\ &= \frac{1}{(q; q)_{\infty}^2} \left( 1 + 2 \sum_{m=1}^{\infty} (-1)^m q^{m^2} \right). \end{aligned}$$

By Lemma 7.2.2, the coefficients  $p(k)$  on the left side above are multiples of 5 whenever  $2k \equiv 5j+3 \pmod{5}$ , i.e., whenever  $k = 5n+4$ . This then completes our third proof.  $\square$

Recall that  $\varphi_2(n)$  is defined in Definition 6.1.3.

**Corollary 7.2.3.** *We have*

$$\varphi_2(5n+3) \equiv 0 \pmod{5}. \quad (7.2.13)$$

**Proof.** Recall from (6.1.1) and the third line of the proof of Theorem 6.1.5 that

$$\Phi_2(q) = \sum_{n=0}^{\infty} \varphi_2(n)q^n = \frac{1}{(q; q)_{\infty}^2} \sum_{n=-\infty}^{\infty} \zeta^n q^{n^2}, \quad \zeta = e^{2\pi i/3}. \quad (7.2.14)$$

Now,

$$\sum_{n=-\infty}^{\infty} \zeta^n q^{n^2} = \sum_{m=-\infty}^{\infty} \zeta^{3m} q^{9m^2} + \sum_{\substack{n=1 \\ n \not\equiv 0 \pmod{3}}}^{\infty} (\zeta^n + \zeta^{-n}) q^{n^2}. \quad (7.2.15)$$

But,

$$\zeta^n + \zeta^{-n} = \begin{cases} 2 \cos(2\pi/3) = -1, & n \equiv 1 \pmod{3}, \\ 2 \cos(4\pi/3) = -1, & n \equiv 2 \pmod{3}. \end{cases} \quad (7.2.16)$$

If we use (7.2.16) in (7.2.15) and then (7.2.15) in (7.2.14), we see that the hypotheses of Lemma 7.2.2 are satisfied. Hence, by the aforementioned lemma, the proof of (7.2.13) is complete.  $\square$

**Corollary 7.2.4.** *We have*

$$c\varphi_2(5n+3) \equiv 0 \pmod{5}. \quad (7.2.17)$$

**Proof.** Recall from (6.2.2) and (6.2.5) that

$$c\Phi_2(q) = \sum_{n=0}^{\infty} c\varphi_2(n)q^n = \frac{1}{(q; q)_{\infty}^2} \sum_{n=-\infty}^{\infty} q^{n^2} = \frac{1}{(q; q)_{\infty}^2} \left( 1 + 2 \sum_{n=0}^{\infty} q^{n^2} \right). \quad (7.2.18)$$

Thus, from (7.2.18), we see that (7.2.17) follows immediately to complete the proof.  $\square$

Although the proofs of the congruence  $p(5n+4) \equiv 0 \pmod{5}$  that we have given so far are attractive, they must take second place to the proof arising from the following beautiful identity.

**Theorem 7.2.5.** *We have*

$$\sum_{n=0}^{\infty} p(5n+4)q^n = 5 \frac{(q^5; q^5)_{\infty}^5}{(q; q)_{\infty}^6}. \quad (7.2.19)$$

It is obvious that the congruence  $p(5n+4) \equiv 0 \pmod{5}$  follows directly from (7.2.19).

In singling out an elegant formula of Ramanujan that characterizes Ramanujan's mathematics, Hardy remarked [90, p. xxxv], "and, if I had to select one formula from all Ramanujan's work, I would agree with Major MacMahon in selecting a formula from [87], viz.

$$p(4) + p(9)x + p(14)x^2 + \dots = 5 \frac{\{(1-x^5)(1-x^{10})(1-x^{15})\dots\}^5}{\{(1-x)(1-x^2)(1-x^3)\dots\}^6},$$

where  $p(n)$  is the number of partitions of  $n$ ."

The proof of Theorem 7.2.5 that we shall give is based on another beautiful formula of Ramanujan.

**Theorem 7.2.6.** *If  $\left(\frac{n}{5}\right)$  denotes the Legendre symbol, then*

$$\sum_{n=0}^{\infty} \left(\frac{n}{5}\right) \frac{q^n}{(1-q^n)^2} = q \frac{(q^5; q^5)_{\infty}^5}{(q; q)_{\infty}}. \quad (7.2.20)$$

We first demonstrate that Theorem 7.2.5 follows from Theorem 7.2.6.

**Proof.** From (7.2.20),

$$q(q^5; q^5)_{\infty}^5 \sum_{n=0}^{\infty} p(n)q^n = \sum_{n=0}^{\infty} \left(\frac{n}{5}\right) \frac{q^n}{(1-q^n)^2} = \sum_{n=0}^{\infty} \left(\frac{n}{5}\right) \sum_{k=1}^{\infty} kq^{nk}. \quad (7.2.21)$$

We now equate the terms in (7.2.21) in which the powers of  $q$  are multiples of 5. We note that these powers arise only when  $k$  is a multiple of 5, since  $\left(\frac{n}{5}\right) = 0$  when  $n$  is a multiple of 5. Hence,

$$q(q^5; q^5)_{\infty}^5 \sum_{n=0}^{\infty} p(5n+4)q^{5n+4} = 5 \sum_{n=0}^{\infty} \left(\frac{n}{5}\right) \sum_{k=1}^{\infty} kq^{5nk} = 5q^5 \frac{(q^{25}; q^{25})_{\infty}^5}{(q^5; q^5)_{\infty}}, \quad (7.2.22)$$

where we applied Theorem 7.2.6, or (7.2.21), once again, but now with  $q$  replaced by  $q^5$ . Rewriting (7.2.22) slightly, we see that

$$\sum_{n=0}^{\infty} p(5n+4)q^{5n+4} = 5q^4 \frac{(q^{25}; q^{25})_{\infty}^5}{(q^5; q^5)_{\infty}^6}.$$

Cancelling  $q^4$  on both sides above and replacing  $q^5$  by  $q$ , we complete the proof of Theorem 7.2.6.  $\square$

We next show that Theorem 7.2.6 is a special instance of the next theorem.

**Theorem 7.2.7.** For any complex numbers  $x$  and  $y$ ,

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \left\{ \frac{xq^n}{(1-xq^n)^2} - \frac{yq^n}{(1-yq^n)^2} \right\} \\ = \frac{(x-y)(1-xy)}{(1-x)^2(1-y)^2} \frac{(xyq)_{\infty}(q/(xy)_{\infty}(xq/y)_{\infty}(yq/x)_{\infty}(q; q)_{\infty}^4}{(xq)_{\infty}^2(q/x)_{\infty}^2(yq)_{\infty}^2(q/y)_{\infty}^2}. \end{aligned} \quad (7.2.23)$$

**Proof.** As promised, we show that Theorem 7.2.6 follows from Theorem 7.2.7. Replace  $q$  by  $q^5$  and set  $x = q$  and  $y = q^2$  in (7.2.23). Hence, the left-hand side of (7.2.23) becomes

$$\begin{aligned} & \sum_{n=-\infty}^{\infty} \left\{ \frac{q^{5n+1}}{(1-q^{5n+1})^2} - \frac{q^{5n+2}}{(1-q^{5n+2})^2} \right\} \\ &= \sum_{n=0}^{\infty} \left\{ \frac{q^{5n+1}}{(1-q^{5n+1})^2} - \frac{q^{5n+2}}{(1-q^{5n+2})^2} \right\} + \sum_{j=0}^{\infty} \left\{ \frac{q^{-5j-4}}{(1-q^{-5j-4})^2} - \frac{q^{-5j-3}}{(1-q^{-5j-3})^2} \right\} \\ &= \sum_{n=0}^{\infty} \left\{ \frac{q^{5n+1}}{(1-q^{5n+1})^2} - \frac{q^{5n+2}}{(1-q^{5n+2})^2} \right\} + \sum_{j=0}^{\infty} \left\{ \frac{q^{5j+4}}{(1-q^{5j+4})^2} - \frac{q^{5j+3}}{(1-q^{5j+3})^2} \right\} \\ &= \sum_{n=0}^{\infty} \binom{n}{5} \frac{q^n}{(1-q^n)^2}, \end{aligned} \quad (7.2.24)$$

where we set  $n = -j - 1$  in the sums over the negative indices.

On the other hand, the right-hand side of (7.2.23) becomes, under the above-mentioned substitutions,

$$\begin{aligned} & \frac{(q-q^2)(1-q^3)(q^8; q^5)_{\infty}(q^2; q^5)_{\infty}(q^4; q^5)_{\infty}(q^6; q^5)_{\infty}(q^5; q^5)_{\infty}^4}{(1-q)^2(1-q^2)^2(q^6; q^5)_{\infty}^2(q^4; q^5)_{\infty}^2(q^7; q^5)_{\infty}^2(q^3; q^5)_{\infty}^2} \\ &= q \frac{(q^3; q^5)_{\infty}(q^2; q^5)_{\infty}(q^5; q^5)_{\infty}^4}{(q; q^5)_{\infty}(q^4; q^5)_{\infty}(q^2; q^5)_{\infty}^2(q^3; q^5)_{\infty}^2} = q \frac{(q^5; q^5)_{\infty}^5}{(q; q)_{\infty}}. \end{aligned} \quad (7.2.25)$$

If we now put together (7.2.24) and (7.2.25), we deduce Theorem 7.2.6.  $\square$

**Proof of Theorem 7.2.7.** Let

$$F(x, y) := \frac{(x-y)(1-xy)}{(1-x)^2(1-y)^2} \frac{(xyq)_{\infty}(q/(xy)_{\infty}(xq/y)_{\infty}(yq/x)_{\infty}(q; q)_{\infty}^4}{(xq)_{\infty}^2(q/x)_{\infty}^2(yq)_{\infty}^2(q/y)_{\infty}^2}. \quad (7.2.26)$$

and

$$G(x, y) := \frac{(xyq)_{\infty}(q/(xy)_{\infty}(xq/y)_{\infty}(yq/x)_{\infty}(q; q)_{\infty}^4}{(xq)_{\infty}^2(q/x)_{\infty}^2(yq)_{\infty}^2(q/y)_{\infty}^2}. \quad (7.2.27)$$

Suppose that  $y \neq q^n$ ,  $-\infty < n < \infty$ . Regard  $F(x, y)$  as a function of  $x$ , with  $y$  constant for the time being. Observe that  $F(x, y)$  has double poles at  $x = q^n$ ,  $-\infty < n < \infty$ . Our first goal is to find the principal parts of  $F(x, y)$  at these poles. It will help us to first

derive some functional equations. By (7.2.26),

$$\begin{aligned}
F(qx, y) &= \frac{(qx - y)(1 - qxy)}{(1 - qx)^2(1 - y)^2} \frac{(xyq^2)_\infty (1/(xy))_\infty (xq^2/y)_\infty (y/x)_\infty (q; q)_\infty^4}{(xq^2)_\infty^2 (1/x)_\infty^2 (yq)_\infty^2 (q/y)_\infty^2} \\
&= \frac{(xq - y)(1 - 1/(xy))(1 - y/x)}{(1 - qx/y)(1 - y)^2(1 - 1/x)^2} G(x, y) \\
&= \frac{(x - y)(1 - xy)}{(1 - x)^2(1 - y)^2} G(x, y) \\
&= F(x, y).
\end{aligned} \tag{7.2.28}$$

Also, by (7.2.28),

$$F(q^{-1}x, y) = F(q(q^{-1}x), y) = F(x, y). \tag{7.2.29}$$

Observe that, from the definition (7.2.26),

$$\lim_{x \rightarrow 1} (1 - x)^2 F(x, y) = 1. \tag{7.2.30}$$

Thus, using the definition (7.2.27), we so far know that

$$F(x, y) - \frac{1}{(x - 1)^2} = \frac{1}{(x - 1)^2} \left( \frac{(x - y)(1 - xy)}{(1 - y)^2} G(x, y) - 1 \right). \tag{7.2.31}$$

We now need to calculate

$$\begin{aligned}
\lim_{x \rightarrow 1} (x - 1) \left( F(x, y) - \frac{1}{(x - 1)^2} \right) &= \lim_{x \rightarrow 1} \frac{1}{x - 1} \left( \frac{(x - y)(1 - xy)}{(1 - y)^2} G(x, y) - 1 \right) \\
&= \lim_{x \rightarrow 1} \left( \left\{ \frac{1 - xy}{(1 - y)^2} - \frac{y(x - y)}{(1 - y)^2} \right\} G(x, y) + \frac{(x - y)(1 - xy)}{(1 - y)^2} \frac{\partial}{\partial x} G(x, y) \right) \\
&= 1 + \lim_{x \rightarrow 1} \frac{(x - y)(1 - xy)}{(1 - y)^2} G(x, y) \frac{\partial}{\partial x} \log G(x, y) \\
&= 1 + \lim_{x \rightarrow 1} \frac{\partial}{\partial x} \log G(x, y),
\end{aligned} \tag{7.2.32}$$

because  $G(1, y) = 1$ . Now

$$\begin{aligned}
\frac{\partial}{\partial x} \log G(x, y) &= \sum_{n=1}^{\infty} \left( -\frac{yq^n}{1 - xyq^n} + \frac{q^n/(x^2y)}{1 - q^n/(xy)} - \frac{q^n/y}{1 - xq^n/y} \right. \\
&\quad \left. + \frac{yq^n/x^2}{1 - yq^n/x} + 2\frac{q^n}{1 - xq^n} - 2\frac{q^n/x^2}{1 - q^n/x} \right).
\end{aligned}$$

It follows from the calculation above that

$$\lim_{x \rightarrow 1} \frac{\partial}{\partial x} \log G(x, y) = 0. \tag{7.2.33}$$

We provide another argument suggested by Dan Schultz. From (7.2.27), it is easily seen that  $G(x, y) = G(1/x, y)$ . Thus, by the chain rule, if  $u = 1/x$ ,

$$\frac{\partial}{\partial x} G(x, y) = -\frac{1}{x^2} \frac{\partial}{\partial u} G(u, y).$$



If we now set  $x = 1$  (so that  $u = 1$ ) and replace  $x$  by  $u$  on the right-hand side, we see that

$$\frac{\partial}{\partial x} G(1, y) = -\frac{\partial}{\partial x} G(1, y).$$

Hence,

$$\frac{\partial}{\partial x} G(1, y) = 0. \quad (7.2.34)$$

So, using either (7.2.33) or (7.2.34) in (7.2.32), we conclude that

$$\lim_{x \rightarrow 1} (x-1) \left( F(x, y) - \frac{1}{(x-1)^2} \right) = 1. \quad (7.2.35)$$

Hence, by (7.2.30) and (7.2.32), the principal part of  $F(x, y)$  about  $x = 1$  equals

$$\frac{1}{(x-1)^2} + \frac{1}{x-1} = \frac{x}{(x-1)^2}.$$

Since, by (7.2.28) and (7.2.29),  $F(x, y) = F(qx, y) = F(q/x, y)$ , it follows that the principal part about the pole  $q^{-n}$ ,  $-\infty < n < \infty$ , equals

$$\frac{xq^n}{(1-xq^n)^2}.$$

Hence, we have shown so far that

$$\begin{aligned} F(x, y) &= \frac{x}{(1-x)^2} + \sum_{n=1}^{\infty} \left( \frac{xq^n}{(1-xq^n)^2} + \frac{x^{-1}q^n}{(1-x^{-1}q^n)^2} \right) + \cdots \\ &=: G(x) + H(x, y), \end{aligned} \quad (7.2.36)$$

say. Now write

$$H(x, y) = \sum_{n=-\infty}^{\infty} a_n(y)x^n. \quad (7.2.37)$$

Since we have subtracted all of the principal parts of  $F(x, y)$  (except around  $x = 0$ ) in (7.2.36) in defining  $H(x, y)$ , it follows that (7.2.37) is valid for  $0 < |x| < \infty$ . Now,  $F(qx, y) = F(x, y)$ , and clearly from above  $G(x) = G(qx)$ . It follows from (7.2.36) that  $H(x, y) = H(qx, y)$ . Hence,

$$H(qx, y) = \sum_{n=-\infty}^{\infty} a_n(y)(qx)^n = H(x, y) = \sum_{n=-\infty}^{\infty} a_n(y)x^n.$$

It follows that

$$a_n(y)q^n = a_n(y), \quad -\infty < n < \infty.$$

Hence,  $a_n(y) = 0$ ,  $n \neq 0$ , and so

$$H(x, y) = a_0(y).$$

It then follows from (7.2.36) that

$$F(x, y) = G(x) + a_0(y).$$

Putting  $x = y$  above, we conclude that

$$0 = F(y, y) = G(y) + a_0(y), \quad \text{or} \quad a_0(y) = -G(y).$$

Thus,

$$F(x, y) = G(x) - G(y),$$

which is what we wanted to prove.  $\square$

Theorem 7.2.5 can be utilized to provide a proof of Ramanujan's congruence for  $p(n)$  modulo 25.

**Theorem 7.2.8.** *For every nonnegative integer  $n$ ,*

$$p(25n + 24) \equiv 0 \pmod{25}. \quad (7.2.38)$$

**Proof.** Applying the binomial theorem on the right side of (7.2.19), we find that

$$\sum_{n=0}^{\infty} p(5n + 4)q^n \equiv 5 \frac{(q^5; q^5)_{\infty}^4}{(q; q)_{\infty}} = 5(q^5; q^5)_{\infty}^4 \sum_{n=0}^{\infty} p(n)q^n \pmod{25}. \quad (7.2.39)$$

From Theorem 7.2.1 we know that the coefficients of  $q^4, q^9, q^{14}, \dots, q^{5n+4}, \dots$  on the far right side of (7.2.39) are all multiples of 25. It follows that the coefficients of  $q^{5n+4}, n \geq 0$ , on the far left side of (7.2.39) are also multiples of 25, i.e.,

$$p(25n + 24) \equiv 0 \pmod{25}.$$

This completes the proof.  $\square$

### 7.3. Ramanujan's Congruence

$$p(7n + 5) \equiv 0 \pmod{7}$$

**Theorem 7.3.1.** *For each nonnegative integer  $n$ ,*

$$p(7n + 5) \equiv 0 \pmod{7}. \quad (7.3.1)$$

**Proof.** Our proof is again taken from Ramanujan's paper [87] and was sketched by Hardy [67, p. 88].

First, by the binomial theorem,

$$\begin{aligned} q^2(q^7; q^7)_{\infty} \sum_{n=0}^{\infty} p(n)q^n &= q^2 \frac{(q^7; q^7)_{\infty}}{(q; q)_{\infty}} = q^2(q; q)_{\infty}^6 \frac{(q^7; q^7)_{\infty}}{(q; q)_{\infty}^7} \\ &\equiv q^2(q; q)_{\infty}^6 \pmod{7}. \end{aligned} \quad (7.3.2)$$

Hence, if we can show that the coefficient of  $q^{7n+7}, n \geq 0$ , in  $q^2(q; q)_{\infty}^6$  is a multiple of 7, it will follow from (7.3.2) that the coefficient of  $q^{7n+7}$  on the far left side is a multiple of 7, i.e.,  $p(7n + 5) \equiv 0 \pmod{7}$ .

Applying Jacobi's identity, Theorem 7.2.7, we find that

$$\begin{aligned} q^2(q; q)_{\infty}^6 &= q^2 \{(q; q)_{\infty}^3\}^2 \\ &= \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} (-1)^{j+k} (2j+1)(2k+1) q^{2+j(j+1)/2+k(k+1)/2}. \end{aligned} \quad (7.3.3)$$

As we saw in the previous paragraph, we want to know when the exponents above are multiples of 7. Now observe that

$$(2j + 1)^2 + (2k + 1)^2 = 8\{2 + \frac{1}{2}j(j + 1) + \frac{1}{2}k(k + 1)\} - 14,$$

and so  $2 + \frac{1}{2}j(j + 1) + \frac{1}{2}k(k + 1)$  is a multiple of 7 if and only if

$$(2j + 1)^2 + (2k + 1)^2 \equiv 0 \pmod{7}. \quad (7.3.4)$$

We easily see that  $(2j + 1)^2, (2k + 1)^2 \equiv 0, 1, 2, 4 \pmod{7}$ , and so the only way (7.3.4) can hold is if both  $(2j + 1)^2, (2k + 1)^2 \equiv 0 \pmod{7}$ . In such cases, we trivially see that the coefficients on the right side of (7.3.3) are multiples of 7. Hence, the coefficient of  $q^{7n+7}, n \geq 1$ , on the left side of (7.3.3) is a multiple of 7. As we demonstrated in the foregoing paragraph, this implies that  $p(7n + 5) \equiv 0 \pmod{7}$ .  $\square$

The following variant of Ramanujan's proof is due to Mike Hirschhorn [70].

**Second Proof of Theorem 7.3.1.** Return to Jacobi's Identity (7.2.7) and gather together terms according to the residue classes of the powers of  $q$  modulo 7. We thus find that

$$J \equiv J_0 + J_1 + J_3 \pmod{7}, \quad (7.3.5)$$

where  $J_j$  contains all of the terms in which the power of  $q$  is congruent to  $j$  modulo 7. On the other hand, by the generating function for  $p(n)$ , the binomial theorem, and (7.3.5),

$$\begin{aligned} \sum_{n=0}^{\infty} p(n)q^n &= \frac{1}{(q; q)_{\infty}} = \frac{(q; q)_{\infty}^6}{(q; q)_{\infty}^7} \\ &= \frac{J^2}{(q; q)_{\infty}^7} \equiv \frac{(J_0 + J_1 + J_3)^2}{(q^7; q^7)_{\infty}} \\ &= \frac{J_0^2 + J_1^2 + J_3^2 + 2J_0J_1 + 2J_0J_3 + 2J_1J_3}{(q^7; q^7)_{\infty}} \pmod{7}. \end{aligned}$$

A careful examination of the terms on the right-hand side above shows that none of the powers are congruent to 5 modulo 7. Thus,  $p(7n + 5) \equiv 0 \pmod{7}$ .  $\square$

**Theorem 7.3.2.** *We have*

$$\sum_{n=0}^{\infty} p(7n + 5)q^n = 7 \frac{(q^7; q^7)_{\infty}^3}{(q; q)_{\infty}^4} + 49q \frac{(q^7; q^7)_{\infty}^7}{(q; q)_{\infty}^8}. \quad (7.3.6)$$

Theorem 7.3.2 is clearly an analogue of Theorem 7.2.6. Furthermore, it is clear that Theorem 7.3.1 is an immediate corollary of Theorem 7.3.2. Ramanujan's proof of Theorem 7.3.2 can be found in his unpublished manuscript on  $p(n)$  and  $\tau(n)$ , published for the first time with his lost notebook [92]. Ramanujan, however, provided almost no details, which were worked out for the first time by the present author, Ae Ja Yee, and Jinhee Yi [38]. The account of Ramanujan's proof of Theorem 7.3.2 that we give below is taken from [38].

**Proof of Theorem 7.3.2.** Using the pentagonal number theorem (1.2.23) in both the numerator and denominator and then separating the indices of summation in the numerator into residue classes modulo 7, we readily find that

$$\frac{(q^{1/7}; q^{1/7})_\infty}{(q^7; q^7)_\infty} = J_1 + q^{1/7} J_2 - q^{2/7} + q^{5/7} J_3, \quad (7.3.7)$$

where  $J_1, J_2$ , and  $J_3$  are power series in  $q$  with integral coefficients, and where the pentagonal number theorem was used to calculate the coefficient of  $q^{2/7}$ . Cubing both sides of (7.3.7), we find that

$$\begin{aligned} & \frac{(q^{1/7}; q^{1/7})_\infty^3}{(q^7; q^7)_\infty^3} \\ &= (J_1^3 + 3J_2^2 J_3 q - 6J_1 J_3 q) + q^{1/7} (3J_1^2 J_2 - 6J_2 J_3 q + J_3^2 q^2) \\ & \quad + 3q^{2/7} (J_1 J_2^2 - J_1^2 + J_3 q) + q^{3/7} (J_2^3 - 6J_1 J_2 + 3J_1 J_3^2 q) \\ & \quad + 3q^{4/7} (J_1 - J_2^2 + J_2 J_3^2 q) + 3q^{5/7} (J_2 + J_1^2 J_3 - J_3^2 q) \\ & \quad + q^{6/7} (6J_1 J_2 J_3 - 1). \end{aligned} \quad (7.3.8)$$

On the other hand, using Jacobi's identity, Corollary 3.1.9, and separating the indices of summation in the numerator on the left side of (7.3.8) into residue classes modulo 7, we easily find that

$$\frac{(q^{1/7}; q^{1/7})_\infty^3}{(q^7; q^7)_\infty^3} = G_1 + q^{1/7} G_2 + q^{3/7} G_3 - 7q^{6/7}, \quad (7.3.9)$$

where  $G_1, G_2$ , and  $G_3$  are power series in  $q$  with integral coefficients, and where Jacobi's identity, Corollary 3.1.9, was used to determine the coefficient of  $q^{6/7}$ . Comparing coefficients in (7.3.8) and (7.3.9), we conclude that

$$\begin{cases} J_1 J_2^2 - J_1^2 + J_3 q &= 0, \\ J_1 - J_2^2 + J_2 J_3^2 q &= 0, \\ J_2 + J_1^2 J_3 - J_3^2 q &= 0, \\ 6J_1 J_2 J_3 - 1 &= -7. \end{cases} \quad (7.3.10)$$

Replace  $q^{1/7}$  by  $\omega q^{1/7}$  in (7.3.7), where  $\omega$  is any seventh root of unity. Therefore,

$$\frac{(\omega q^{1/7}; \omega q^{1/7})_\infty}{(q^7; q^7)_\infty} = J_1 + \omega q^{1/7} J_2 - \omega^2 q^{2/7} + \omega^5 q^{5/7} J_3. \quad (7.3.11)$$

Taking the products of both sides of (7.3.11) over all seven seventh roots of unity, we find that

$$\frac{(q; q)_\infty^8}{(q^7; q^7)_\infty^8} = \prod_{\omega} (J_1 + \omega q^{1/7} J_2 - \omega^2 q^{2/7} + \omega^5 q^{5/7} J_3). \quad (7.3.12)$$

Using the generating function for  $p(n)$ , (7.3.7), and (7.3.12), we find that

$$\begin{aligned} \sum_{n=0}^{\infty} p(n)q^n &= \frac{1}{(q; q)_{\infty}} = \frac{(q^{49}; q^{49})_{\infty}^7 (q^7; q^7)_{\infty}^8 (q^{49}; q^{49})_{\infty}}{(q^7; q^7)_{\infty}^8 (q^{49}; q^{49})_{\infty}^8 (q; q)_{\infty}} \\ &= \frac{(q^{49}; q^{49})_{\infty}^7 \prod_{\omega} (J_1 + \omega q J_2 - \omega^2 q^2 + \omega^5 q^5 J_3)}{(q^7; q^7)_{\infty}^8 \prod_{\omega} (J_1 + \omega q J_2 - \omega^2 q^2 + \omega^5 q^5 J_3)} \\ &= \frac{(q^{49}; q^{49})_{\infty}^7}{(q^7; q^7)_{\infty}^8} \left\{ \prod_{\omega \neq 1} (J_1 + \omega q J_2 - \omega^2 q^2 + \omega^5 q^5 J_3) \right\}. \end{aligned} \quad (7.3.13)$$

We only need to compute the terms in  $\prod_{\omega \neq 1} (J_1 + \omega q J_2 - \omega^2 q^2 + \omega^5 q^5 J_3)$  where the powers of  $q$  are of the form  $7n + 5$  to complete the proof. In order to do this, we need to prove several identities using the identities of (7.3.10). More precisely, we need to prove that

$$J_1^7 + J_2^7 q + J_3^7 q^5 = \frac{(q; q)_{\infty}^8}{(q^7; q^7)_{\infty}^8} + 14q \frac{(q; q)_{\infty}^4}{(q^7; q^7)_{\infty}^4} + 57q^2, \quad (7.3.14)$$

$$J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 = -\frac{(q; q)_{\infty}^4}{(q^7; q^7)_{\infty}^4} - 8q, \quad (7.3.15)$$

$$J_1^2 J_2^3 + J_3^2 J_1^3 q + J_2^2 J_3^3 q^2 = -\frac{(q; q)_{\infty}^4}{(q^7; q^7)_{\infty}^4} - 5q. \quad (7.3.16)$$

Since  $J_2^2 = J_1 + J_2 J_3^2 q$ ,  $J_1^2 = J_1 J_2^2 + J_3 q$ ,  $J_3^2 q = J_2 + J_1^2 J_3$ , and  $J_1 J_2 J_3 = -1$  by (7.3.10), we find that

$$\begin{aligned} J_1^2 J_2^3 + J_3^2 J_1^3 q + J_2^2 J_3^3 q^2 &= J_1^3 J_2 + J_1^2 J_2^2 J_3^2 q + J_1^2 J_2^2 J_3^2 q + J_3^3 J_1 q^2 + J_2^3 J_3 q + J_1^2 J_2^2 J_3^2 q \\ &= J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 + 3q, \end{aligned} \quad (7.3.17)$$

$$\begin{aligned} J_1 J_2^5 + J_3 J_1^5 + J_2 J_3^5 q^3 &= J_1 J_2 (J_1 + J_2 J_3^2 q)^2 + J_3 J_1 (J_1 J_2^2 + J_3 q)^2 + J_2 J_3 (J_2 + J_1^2 J_3)^2 q \\ &= J_1^3 J_2 + 2J_1^2 J_2^2 J_3^2 q + J_1 J_2^3 J_3^4 q^2 + J_3 J_1^3 J_2^4 + 2J_1^2 J_2^2 J_3^2 q + J_3^3 J_1 q^2 \\ &\quad + J_2^3 J_3 q + 2J_1^2 J_2^2 J_3^2 q + J_2 J_3^3 J_1^4 q \\ &= J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 - (J_1^2 J_2^2 + J_2^2 J_1^2 q + J_2^2 J_3^3 q^2) + 6q \\ &= 3q, \end{aligned} \quad (7.3.18)$$

where (7.3.18) is obtained from (7.3.17). (Observe from (7.3.17) that it suffices to prove only (7.3.15) or (7.3.16).) By squaring the left side of (7.3.15) and using (7.3.10), (7.3.18),

and (7.3.17), we find that

$$\begin{aligned}
(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2)^2 &= J_1^6 J_2^2 + J_2^6 J_3^2 q^2 + J_3^6 J_1^2 q^4 \\
&\quad + 2(J_1^3 J_2^4 J_3 q + J_1 J_2^3 J_3^4 q^3 + J_1^4 J_2 J_3^3 q^2) \\
&= J_1^7 + J_1^6 J_2 J_3^2 q + J_2^7 q + J_1^2 J_2^6 J_3 q + J_3^7 q^5 + J_1 J_2^2 J_3^6 q^4 \\
&\quad - 2(J_1^2 J_2^3 q + J_2^2 J_1^3 q^2 + J_2^2 J_3^3 q^3) \\
&= J_1^7 + J_2^7 q + J_3^7 q^5 - (J_1 J_2^5 q + J_3 J_1^5 q + J_2 J_3^5 q^4) \\
&\quad - 2(J_1^2 J_2^2 q + J_2^2 J_3^3 q^3 + J_1^3 J_2^2 q^2) \\
&= J_1^7 + J_2^7 q + J_3^7 q^5 - 2q(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2) - 9q^2.
\end{aligned}$$

Thus,

$$(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 + q)^2 = (J_1^7 + J_2^7 q + J_3^7 q^5) - 8q^2. \quad (7.3.19)$$

Expanding the right side of (3.3.26) and using (3.1.37), (7.3.18), and (7.3.17), we obtain

$$\begin{aligned}
\frac{(q; q)_\infty^8}{(q^7; q^7)_\infty^8} &= J_1^7 + J_2^7 q + J_3^7 q^5 + 7(J_1 J_2^5 q + J_3 J_1^5 q + J_2 J_3^5 q^4) + 7(J_1^4 J_2^2 J_3 q + J_1 J_2^4 J_3^2 q^2 \\
&\quad + J_2 J_3^4 J_1^2 q^3) + 7(J_1^3 J_2 q + J_2^3 J_3 q^2 + J_3^3 J_1 q^3) \\
&\quad + 14(J_1^2 J_2^3 q + J_2^2 J_1^3 q^2 + J_2^2 J_3^3 q^3) + 7J_1^2 J_2^2 J_3^2 q^2 + 14J_1 J_2 J_3 q^2 - q^2 \\
&= J_1^7 + J_2^7 q + J_3^7 q^5 + 21q^2 - 7q(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2) + 7q(J_1^3 J_2 + J_2^3 J_3 q \\
&\quad + J_3^3 J_1 q^2) + 14q(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 + 3q) + 7q^2 - 14q^2 - q^2 \\
&= J_1^7 + J_2^7 q + J_3^7 q^5 + 14q(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2) + 55q^2. \quad (7.3.20)
\end{aligned}$$

Combining (7.3.19) and (7.3.20), we find that

$$\begin{aligned}
\frac{(q; q)_\infty^8}{(q^7; q^7)_\infty^8} &= (J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 + q)^2 + 8q^2 + 14(J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2)q + 55q^2 \\
&= (J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 + 8q)^2.
\end{aligned}$$

By (3.1.34), we see that for  $q$  sufficiently small and positive,  $J_2 < 0$ . Thus, taking the square root of both sides above, we find that

$$J_1^3 J_2 + J_2^3 J_3 q + J_3^3 J_1 q^2 = -\frac{(q; q)_\infty^4}{(q^7; q^7)_\infty^4} - 8q, \quad (7.3.21)$$

which proves (7.3.15). We now see that (7.3.14) follows from (7.3.20) and (7.3.21), and (7.3.16) follows from (7.3.17) and (7.3.21).

Returning to (3.3.27), we are now ready to compute the terms in  $\prod_{i=1}^6 (J_1 + \omega^i q J_2 - \omega^{2i} q^2 + \omega^{5i} q^5 J_3)$  where the powers of  $q$  are of the form  $7n + 5$ . Using the computer algebra system MAPLE, (7.3.15), (7.3.16), and (7.3.18), we find that the desired terms

with powers of the form  $q^{7n+5}$  are equal to

$$\begin{aligned}
& - (J_1 J_2^5 + J_3 J_1^5 + 3J_1^3 J_2 + 4J_1^2 J_2^3) q^5 - (3J_2^3 J_3 + 4J_3^2 J_1^3 - 8) q^{12} \\
& - (4J_2^2 J_3^3 + 3J_3^3 J_1) q^{19} - J_2 J_3^5 q^{26} \\
= & - 3(J_1^3 J_2 + J_2^3 J_3 q^7 + J_3^3 J_1 q^{14}) q^5 - 4(J_1^2 J_2^3 + J_3^2 J_1^3 q^7 + J_2^2 J_3^3 q^{14}) q^5 \\
& - (J_1 J_2^5 + J_3 J_1^5 + J_2 J_3^5 q^{21}) q^5 + 8q^{12} \\
= & 7 \frac{(q^7; q^7)_\infty^4}{(q^{49}; q^{49})_\infty^4} q^5 + 49q^{12}. \tag{7.3.22}
\end{aligned}$$

Choosing only those terms on each side of (7.3.13) where the powers of  $q$  are of the form  $7n + 5$  and using the omitted calculations, we find that

$$\sum_{\substack{n=0 \\ n \equiv 5 \pmod{7}}}^{\infty} p(n) q^n = q^5 \frac{(q^{49}; q^{49})_\infty^7}{(q^7; q^7)_\infty^8} \left( 7 \frac{(q^7; q^7)_\infty^4}{(q^{49}; q^{49})_\infty^4} + 49q^7 \right),$$

or

$$\sum_{n=0}^{\infty} p(7n + 5) q^{7n} = 7 \frac{(q^{49}; q^{49})_\infty^3}{(q^7; q^7)_\infty^4} + 49q^7 \frac{(q^{49}; q^{49})_\infty^7}{(q^7; q^7)_\infty^8}. \tag{7.3.23}$$

Replacing  $q^7$  by  $q$  in (7.3.23), we complete the proof of (7.3.6).  $\square$

By comparing (7.3.7) with Entry 17(v) in Chapter 19 of Ramanujan's second notebook [91], [27, p. 303], we see that

$$J_1 = \frac{f(-q^2, -q^5)}{f(-q, -q^6)}, \quad J_2 = -\frac{f(-q^3, -q^4)}{f(-q^2, -q^5)}, \quad \text{and} \quad J_3 = \frac{f(-q, -q^6)}{f(-q^3, -q^4)},$$

where

$$f(a, b) := \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \quad |ab| < 1.$$

In the notation of Section 18 of Chapter 19 in [91], [27, p. 306],

$$\alpha = u^{1/7} = q^{-2/7} J_1, \quad \beta = -v^{1/7} = q^{-1/7} J_2, \quad \text{and} \quad \gamma = w^{1/7} = q^{3/7} J_3. \tag{7.3.24}$$

Thus, the identity (7.3.7) is equivalent to an identity in Entry 18 in Chapter 19 of Ramanujan's second notebook [91], [27, p. 305, eq. (18.2)]. The proof of (7.3.7) given here is much simpler than that given in [27, pp. 306–312].

Recall that the identity of Theorem 7.2.5 yielded in Theorem 7.2.8 a congruence for  $p(n)$  modulo  $5^2$ . Similarly, the identity in Theorem 7.3.2 yields a congruence for  $p(n)$  modulo  $7^2$ , as we now demonstrate.

**Theorem 7.3.3.** *For each nonnegative integer  $n$ ,*

$$p(49n + 47) \equiv 0 \pmod{49}. \tag{7.3.25}$$

**Proof.** Write (7.3.6) in the form

$$\begin{aligned} \sum_{n=0}^{\infty} p(7n+5)q^n &= 7 \frac{(q^7; q^7)_{\infty}^3 (q; q)_{\infty}^3}{(q; q)_{\infty}^7} + 49q \frac{(q^7; q^7)_{\infty}^7}{(q; q)_{\infty}^8} \\ &\equiv 7(q^7; q^7)_{\infty}^2 \sum_{m=0}^{\infty} (-1)^m (2m+1) q^{m(m+1)/2} \pmod{49}, \end{aligned} \quad (7.3.26)$$

by the binomial theorem and Jacobi's identity, Corollary 3.1.9. We now examine the terms on the right side of (7.3.26) where the powers of  $q$  are of the form  $7n+6$ . Separating the summands into residue classes modulo 7, we see that the only terms yielding such exponents are when  $m \equiv 3 \pmod{7}$ . But then  $2m+1 \equiv 0 \pmod{7}$ . Thus, the coefficient of the power  $q^{7n+6}$ ,  $n \geq 1$ , on the right side of (7.3.26) is a multiple of 49. The same must be true, of course, on the left side of (7.3.26), i.e., the coefficient  $p(49n+47)$  must be a multiple of 49, i.e., (7.3.25) has been established.  $\square$

#### 7.4. Ramanujan's Congruence

$$p(11n+6) \equiv 0 \pmod{11}$$

The following lemma is a special case of Ramanujan's  ${}_1\psi_1$ -summation. It is also a special case of a considerably more general theorem on partial fraction decompositions of  $q$ -products [46]. Observe the symmetry in  $x$  and  $y$  in the lemma.

**Lemma 7.4.1.** For  $|q| < |x| < 1$ ,

$$\frac{[xy]_{\infty} (q)_{\infty}^2}{[x, y]_{\infty}} = \sum_{n=-\infty}^{\infty} \frac{x^n}{1 - yq^n}. \quad (7.4.1)$$

**Proof.** We regard

$$F(y) := \frac{[xy]_{\infty} (q)_{\infty}^2}{[x, y]_{\infty}}, \quad |x|, |q/x| < 1, \quad (7.4.2)$$

as a function of the complex variable  $y$ . We calculate the partial fraction expansion of  $F(y)$ . Observe that  $F(y)$  has simple poles at  $y = q^n$ ,  $-\infty < n < \infty$ . Set

$$\begin{aligned} L_n &:= \lim_{y \rightarrow q^{-n}} (y - q^{-n}) F(y) \\ &= \lim_{y \rightarrow q^{-n}} -q^{-n} \frac{(xy; q)_{\infty} (q/(xy); q)_{\infty} (q; q)_{\infty}^2}{(x; q)_{\infty} (q/x; q)_{\infty} (1-y) \cdots (1-yq^{n-1})(1-yq^{n+1}) \cdots (q/y; q)_{\infty}} \\ &= -q^{-n} \frac{(x/q^n; q)_{\infty} (q^{n+1}/x; q)_{\infty} (-1)^n q^{n(n+1)/2} (q; q)_{\infty}^2}{(x; q)_{\infty} (q/x; q)_{\infty} (q; q)_{\infty} (q; q)_{\infty} (q^{n+1}; q)_{\infty}} \\ &= -q^{-n} \frac{(-1)^n x^n (q/x; q)_{\infty} (x; q)_{\infty} (-1)^n q^{n(n+1)/2} (q; q)_{\infty}^2}{q^{n(n+1)/2} (x; q)_{\infty} (q/x; q)_{\infty} (q; q)_{\infty}^2} \\ &= -q^{-n} x^n. \end{aligned}$$

Thus, the associated term in the partial fraction decomposition of  $F(y)$  is equal to

$$-\frac{q^{-n} x^n}{y - q^{-n}} = \frac{x^n}{1 - yq^n}, \quad -\infty < n < \infty.$$



Hence, for some entire function  $G(y)$ ,

$$F(y) = \sum_{n=-\infty}^{\infty} \frac{x^n}{1-yq^n} + G(y) =: H(y) + G(y).$$

We want to show that  $G(y) \equiv 0$ . Observe that

$$F(qy) = \frac{[xqy]_{\infty}(q)_{\infty}^2}{[x, qy]_{\infty}} = \frac{(1-1/(xy))(1-y)}{(1-xy)(1-1/y)} F(y) = \frac{F(y)}{x}.$$

It is easy to see that

$$H(qy) = \frac{H(y)}{x}.$$

It follows that

$$F(y/q) = xF(q \cdot y/q) = xF(y) \quad \text{and} \quad H(y/q) = xH(y). \quad (7.4.3)$$

On  $0 \leq |y| \leq 1$ ,  $G(y) = F(y) - H(y)$  is a bounded analytic function. By (7.4.3),  $|G(y/q)| = |xG(y)|$ , and by induction,

$$|G(y/q^n)| = |x^n G(y)|. \quad (7.4.4)$$

Hence, as  $n \rightarrow \infty$ , we see that  $G(y/q^n)$  tends to 0, since  $|x| < 1$ . In conclusion,  $G(y)$  is an entire function that tends to 0 as  $y$  tends to infinity. It is therefore a bounded entire function, and so, by Liouville's theorem,  $G(y)$  is a constant. But since  $G(y) \rightarrow 0$ , as  $y \rightarrow \infty$ , this constant must be 0. Hence, (7.4.1) follows, and the proof is finished.  $\square$

**Lemma 7.4.2.** [*Halphen's Identity*] For arbitrary complex numbers  $a, b, c$ ,

$$\begin{aligned} H(a, b, c, q) &:= \frac{[ab, bc, ca]_{\infty}(q)_{\infty}^2}{[a, b, c, abc]_{\infty}} \\ &= 1 + F(a, q) + F(b, q) + F(c, q) - F(abc, q), \end{aligned} \quad (7.4.5)$$

where

$$F(x, q) := \sum_{k=0}^{\infty} \frac{xq^k}{1-xq^k} - \sum_{k=1}^{\infty} \frac{q^k/x}{1-q^k/x}, \quad (|q| < 1). \quad (7.4.6)$$

In this particular form, Halphen's identity was first established by Andrews, R. Lewis, and Z.-G. Liu [22]. However, equivalent versions of (7.4.5) in terms of Weierstrass elliptic functions appeared in the literature much earlier, in particular, as exercises in the classical text by Whittaker and Watson [101, Examples 19 and 20, p. 458], and also in G. H. Halphen's paper [66, p. 187]. We will follow the lead of Whittaker and Watson and leave Halphen's identity as an exercise.

**Lemma 7.4.3.** For  $|q| < 1$ ,

$$1 + F(a, q) = -F(1/a, q). \quad (7.4.7)$$

**Proof.** Extracting the term with  $n = 0$  in the first sum below and then adding and subtracting the term with  $n = 0$  in the second sum, we find that

$$\begin{aligned} F(a, q) &= \sum_{n=0}^{\infty} \frac{aq^n}{1-aq^n} - \sum_{n=1}^{\infty} \frac{q^n/a}{1-q^n/a} \\ &= \frac{a}{1-a} + \frac{\frac{1}{a}}{1-\frac{1}{a}} + \sum_{n=1}^{\infty} \frac{q^n/(\frac{1}{a})}{1-q^n/(\frac{1}{a})} - \sum_{n=0}^{\infty} \frac{\frac{1}{a}q^n}{1-\frac{1}{a}q^n} \\ &= -1 - F(1/a, q). \end{aligned}$$

□

**Theorem 7.4.4** (Winquist's Identity). *We have*

$$\sum_{m,n=-\infty}^{\infty} (-1)^{m+n} q^{(3m^2+3n^2+3m+n)/2} (a^{-3m}b^{-3n} - a^{-3m}b^{3n+1} - b^{-3m-1}a^{-3n+1} + b^{-3m-1}a^{3n+2}) = (q; q)_{\infty}^2 [a, b, ab, a/b; q]_{\infty} \quad (7.4.8)$$

**Proof.** We first rewrite (7.4.8) in the equivalent form

$$\begin{aligned} &[a, b, ab, a/b; q]_{\infty} (q; q)_{\infty}^2 \\ &= [a^3; q^3]_{\infty} (q^3; q^3)_{\infty}^2 ([b^3q; q^3]_{\infty} - b[b^3q^2; q^3]_{\infty}) \\ &\quad - \frac{a}{b} [b^3; q^3]_{\infty} (q^3; q^3)_{\infty}^2 ([a^3q; q^3]_{\infty} - a[a^3q^2; q^3]_{\infty}). \end{aligned} \quad (7.4.9)$$

To derive this representation of Winquist's identity, we need to identify each of the eight bilateral series on the left side of (7.4.8) in terms of Ramanujan's theta function  $f(\alpha, \beta)$  and then apply the Jacobi triple product identity (1.1.7) to each of the eight series. We leave this task as an exercise. Our goal is thus to prove (7.4.9).

Let  $\omega$  be a primitive cubic root of unity. For brevity, set  $F(a, q) = F(a)$ . In Halphen's identity (7.4.5), we replace  $a$ ,  $b$ , and  $c$  by  $a\omega$ ,  $b\omega^2$ , and  $\omega^2/b$ , respectively, to deduce that

$$H(a\omega, b\omega^2, \omega^2/b, q) = 1 + F(a\omega) + F(b\omega^2) + F(\omega^2/b) - F(a\omega^2). \quad (7.4.10)$$

Simplifying the left side of (7.4.10), we deduce that

$$\begin{aligned} H(a\omega, b\omega^2, \omega^2/b, q) &= \frac{[ab, a/b, \omega; q]_{\infty} (q; q)_{\infty}^2}{[a\omega, b\omega^2, \omega^2/b, a\omega^2; q]_{\infty}} \\ &= -b\omega(1-\omega) \frac{[a, b, ab, a/b; q]_{\infty} (q; q)_{\infty}^2}{[a^3, b^3, q; q^3]_{\infty}}. \end{aligned} \quad (7.4.11)$$

Note that

$$\begin{aligned} &F(a\omega) - F(a\omega^2) \\ &= \sum_{k=0}^{\infty} \frac{a\omega q^k}{1-a\omega q^k} - \sum_{k=1}^{\infty} \frac{\omega^2 q^k/a}{1-\omega^2 q^k/a} - \sum_{k=0}^{\infty} \frac{a\omega^2 q^k}{1-a\omega^2 q^k} + \sum_{k=1}^{\infty} \frac{\omega q^k/a}{1-\omega q^k/a} \end{aligned}$$

$$\begin{aligned}
&= (\omega - \omega^2) \sum_{k=0}^{\infty} \frac{(1 - aq^k)aq^k}{1 - a^3q^{3k}} + (\omega - \omega^2) \sum_{k=1}^{\infty} \frac{(1 - q^k/a)q^k/a}{1 - q^{3k}/a^3} \\
&= (\omega - \omega^2) \sum_{k=0}^{\infty} \frac{(1 - aq^k)aq^k}{1 - a^3q^{3k}} + (\omega - \omega^2) \sum_{k=1}^{\infty} \frac{(1 - aq^{-k})aq^{-k}}{1 - a^3q^{-3k}} \\
&= a(\omega - \omega^2) \sum_{k=-\infty}^{\infty} \frac{q^k}{1 - a^3q^{3k}} - a^2(\omega - \omega^2) \sum_{k=-\infty}^{\infty} \frac{q^{2k}}{1 - a^3q^{3k}} \\
&= a(\omega - \omega^2) \frac{[a^3q; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[a^3, q; q^3]_{\infty}} - a^2(\omega - \omega^2) \frac{[a^3q^2; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[a^3, q^2; q^3]_{\infty}}, \quad (7.4.12)
\end{aligned}$$

where we applied Lemma 7.4.1 twice in the last equality. By invoking (7.4.7), we similarly deduce that

$$\begin{aligned}
&1 + F(b\omega^2) + F(\omega^2/b) \\
&= F(b\omega^2) - F(b\omega) \\
&= b^2(\omega - \omega^2) \frac{[b^3q^2; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[b^3, q^2; q^3]_{\infty}} - b(\omega - \omega^2) \frac{[b^3q; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[b^3, q; q^3]_{\infty}}. \quad (7.4.13)
\end{aligned}$$

Substituting (7.4.11) on the left side of (7.4.10), and (7.4.12) and (7.4.13) on the right side of (7.4.10), and dividing both sides by  $-b\omega(1 - \omega)$ , we arrive at

$$\begin{aligned}
\frac{[a, b, ab, a/b; q]_{\infty} (q; q)_{\infty}^2}{[a^3, b^3, q; q^3]_{\infty}} &= \frac{[b^3q; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[b^3, q; q^3]_{\infty}} - b \frac{[b^3q^2; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[b^3, q^2; q^3]_{\infty}} \\
&\quad - \frac{a}{b} \frac{[a^3q; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[a^3, q; q^3]_{\infty}} + \frac{a^2}{b} \frac{[a^3q^2; q^3]_{\infty} (q^3; q^3)_{\infty}^2}{[a^3, q^2; q^3]_{\infty}}.
\end{aligned}$$

Upon multiplying both sides by  $[a^3, b^3, q; q^3]_{\infty}$ , we obtain Winquist's identity in the form (7.4.9).  $\square$

**Theorem 7.4.5.** *For every nonnegative integer  $n$ ,*

$$p(11n + 6) \equiv 0 \pmod{11}. \quad (7.4.14)$$

**Proof.** In Winquist's identity (7.4.8), replace  $a$  and  $b$  by  $a^2$  and  $b^2$ , respectively, and then multiply both sides by  $a^{-1}b^{-1}$ . Noting that

$$a^{-1}b^{-1}(1 - a^2)(1 - b^2) = (a - a^{-1})(b - b^{-1}),$$

we find that

$$\begin{aligned}
&\sum_{m, n=-\infty}^{\infty} (-1)^{m+n} q^{(3m^2+3n^2+3m+n)/2} (a^{-6m-1}b^{-6n-1} \\
&\quad - a^{-6m-1}b^{6n+1} - b^{-6m-3}a^{-6n+1} + b^{-6m-3}a^{6n+3}) \\
&= (a - a^{-1})(b - b^{-1})(q; q)_{\infty}^2 (a^2q; q)_{\infty} (a^{-2}q; q)_{\infty} (b^2q; q)_{\infty} (b^{-2}q; q)_{\infty} \\
&\quad \times (a^2b^2; q)_{\infty} (qa^{-2}b^{-2}; q)_{\infty} (a^2b^{-2}; q)_{\infty} (qa^{-2}b^2; q)_{\infty}. \quad (7.4.15)
\end{aligned}$$

Differentiate both sides of (7.4.15) with respect to  $b$  and then set  $b = 1$ . Then multiply both sides by  $a^{-2}$ . Because  $b - b^{-1} = 0$  when  $b = 1$ , we see that only one expression on the right-hand side survives, namely, that from differentiating  $b - b^{-1}$ . We note that

$$\left. \frac{d}{db}(b - b^{-1}) \right|_{b=1} = 1 + b^{-2}|_{b=1} = 2.$$

Also,

$$a^{-2}(1 - a^2b^2)(1 - a^2b^{-2})|_{b=1} = (a - a^{-1})^2.$$

With all of these remarks in mind, we find that the differentiation of (7.4.15) with respect to  $b$  and the setting of  $b = 1$  yields

$$\begin{aligned} & \sum_{m,n=-\infty}^{\infty} (-1)^{m+n} q^{(3m^2+3n^2+3m+n)/2} \\ & \times (-2(6n+1)a^{-6m-3} + (6m+3)a^{-6n-1} - (6m+3)a^{6n+1}) \\ & = 2(a - a^{-1})^3 (q; q)_{\infty}^4 (a^2q; q)_{\infty}^3 (a^{-2}q; q)_{\infty}^3. \end{aligned} \quad (7.4.16)$$

We now apply the operator  $a \frac{d}{da}$  three times to (7.4.16). Then we set  $a = 1$ , and lastly divide both sides by 2. On the right-hand side, all of the expressions will equal 0 when we set  $a = 1$ , except for the contribution obtained by three applications of the aforementioned operator to  $(1 - a^{-2})^3$ . To help readers with these calculations, we write  $2(a - a^{-1})^3 = 2a^3(1 - a^{-2})^3$ . The contributions after three applications of the given operator are:

$$\begin{aligned} 6a^3(1 - a^{-2})^2 \cdot 2a^{-3} \cdot a &= 12a(1 - a^{-2})^2, \\ 24a(1 - a^{-2}) \cdot 2a^{-3} \cdot a &= 48a^{-1}(1 - a^{-2}), \\ 48a^{-1} \cdot 2a^{-3} \cdot a &= 96a^{-3}. \end{aligned}$$

With the aid of the calculations above, we find that

$$\begin{aligned} & \sum_{m,n=-\infty}^{\infty} (-1)^{m+n} q^{(3m^2+3n^2+3m+n)/2} \{ (6m+3)^3(6n+1) - (6m+3)(6n+1)^3 \} \\ & = 48(q; q)_{\infty}^{10}. \end{aligned} \quad (7.4.17)$$

Now set  $\alpha = 6m + 3$ ,  $\beta = 6n + 1$ . A straightforward calculation shows that

$$3m^2 + 3n^2 + 3m + n = \frac{1}{12} (\alpha^2 + \beta^2 - 10). \quad (7.4.18)$$

Hence, we may write (7.4.17) in the shape

$$\sum_{\substack{\alpha, \beta = -\infty \\ \alpha \equiv 3 \pmod{6} \\ \beta \equiv 1 \pmod{6}}}^{\infty} (-1)^{(\alpha+\beta-4)/6} (\alpha^3\beta - \alpha\beta^3) q^{(\alpha^2+\beta^2-10)/24} = 48(q; q)_{\infty}^{10}. \quad (7.4.19)$$

If we set

$$(q; q)_{\infty}^{10} =: \sum_{n=0}^{\infty} a(n)q^n, \quad (7.4.20)$$

then, equating coefficients of  $q^n$ ,  $n \geq 0$ , in (7.4.19), we find that

$$a(n) = \frac{1}{48} \sum_{\substack{\alpha, \beta = -\infty \\ \alpha \equiv 3 \pmod{6} \\ \beta \equiv 1 \pmod{6} \\ (\alpha^2 + \beta^2 - 10)/24 = n}}^{\infty} (-1)^{(\alpha + \beta - 4)/6} (\alpha^3 \beta - \alpha \beta^3).$$

Now,

$$\begin{aligned} \frac{1}{24}(\alpha^2 + \beta^2 - 10) &\equiv 6 \pmod{11}, \\ \iff \alpha^2 + \beta^2 - 10 &\equiv 1 \pmod{11}, \\ \iff \alpha^2 + \beta^2 &\equiv 0 \pmod{11}, \\ \iff \alpha, \beta &\equiv 0 \pmod{11}. \end{aligned}$$

Hence, if  $n \equiv 6 \pmod{11}$ , then  $a(n) \equiv 0 \pmod{11^4}$ .

From (7.4.20),

$$\sum_{n=0}^{\infty} p(n)q^n = \frac{1}{(q; q)_{\infty}} = \frac{(q; q)_{\infty}^{10}}{(q; q)_{\infty}^{11}} \equiv \frac{(q; q)_{\infty}^{10}}{(q^{11}; q^{11})_{\infty}} = \frac{1}{(q^{11}; q^{11})_{\infty}} \sum_{n=0}^{\infty} a(n)q^n \pmod{11}.$$

Extracting those terms with powers of  $q$  congruent to 6 modulo 11 and using the subsequent congruence  $a(n) \equiv 0 \pmod{11^4}$ , we see that

$$\sum_{n=0}^{\infty} p(11n + 6)q^{11n+6} \equiv \frac{1}{(q^{11}; q^{11})_{\infty}} \sum_{n=0}^{\infty} a(11n + 6)q^{11n+6} \pmod{11}.$$

Thus,

$$p(11n + 6) \equiv 0 \pmod{11},$$

which is what we wanted to prove. □

Another proof of the identity (7.4.17) for  $(q; q)_{\infty}^{10}$  has been given by the author, S. H. Chan, Z.-G. Liu, and H. Yesilyurt [34]. We remark that (7.4.17) shows that  $(q; q)_{\infty}^{10}$  is lacunary. By the pentagonal number theorem (1.2.23) and Jacobi’s identity (3.1.31), respectively,  $(q; q)_{\infty}$  and  $(q; q)_{\infty}^3$  are also lacunary. It is natural to ask what powers  $(q; q)_{\infty}^n$  are lacunary, and this was answered by J.-P. Serre [97]. Briefly, there are very few powers of the eta-function that are lacunary.

### 7.5. A More General Partition Function

In a letter to Hardy written from Fitzroy House late in 1918 [37, pp. 192–193], Ramanujan writes, “I have considered more or less exhaustively about the congruency of  $p(n)$  and in general that of  $p_r(n)$  where

$$\sum p_r(n)x^n = \frac{1}{(x; x)_{\infty}^r},$$

by four different methods.” This declaration appears to imply that he had established several results about  $p_r(n)$ . However, the only work that remains of Ramanujan on this

more general partition function is page 182 in the volume containing Ramanujan's lost notebook. The page has "5" written in the upper right-hand corner clearly indicating that this page belonged to a much longer manuscript that has evidently been lost. On this page, Ramanujan's general partition function is defined slightly differently, and we adopt this definition here. Define the more general partition function  $p_r(n)$  by

$$\frac{1}{(q; q)_\infty^r} = \sum_{n=0}^{\infty} p_r(n)q^n, \quad |q| < 1. \quad (7.5.1)$$

This definition is actually not provided on page 182, but it is clear that it must have been given somewhere in the missing pages 1–4 of the manuscript. Of course,  $p_1(n) = p(n)$ . The elementary methods developed by Ramanujan to prove congruences for  $p(n)$  modulo 5 and 7 are sufficient to establish all the results on this page. We follow the exposition of the writer, C. Gugg, and S. Kim in their paper [35]. Formerly, a brief account of page 182 was given by K. G. Ramanathan [84].

**Entry 7.5.1** (p. 182). *Let  $\delta$  denote any integer, and let  $n$  denote a nonnegative integer. Suppose that  $\varpi$  is a prime of the form  $6\lambda - 1$ . Then*

$$p_{\delta\varpi-4} \left( n\varpi - \frac{\varpi+1}{6} \right) \equiv 0 \pmod{\varpi}. \quad (7.5.2)$$

**Proof.** Recalling that  $\lambda = (\varpi + 1)/6$ , consider

$$\begin{aligned} \sum_{n=0}^{\infty} p_{\delta\varpi-4}(n)q^{n+\lambda} &= (q; q)_\infty^{-\delta\varpi} (q; q)_\infty^3 (q; q)_\infty q^\lambda \\ &\equiv (q^\varpi; q^\varpi)_\infty^{-\delta} \sum_{\mu=0}^{\infty} \sum_{\nu=-\infty}^{\infty} (-1)^{\mu+\nu} (2\mu+1) q^{\frac{1}{2}\mu(\mu+1) + \frac{1}{2}\nu(3\nu+1) + \lambda} \pmod{\varpi}, \end{aligned} \quad (7.5.3)$$

upon the use of Euler's pentagonal number theorem (1.2.23) and Jacobi's identity (3.1.31). We want to examine those terms for which

$$\frac{1}{2}\mu(\mu+1) + \frac{1}{2}\nu(3\nu+1) + \frac{\varpi+1}{6} \equiv 0 \pmod{\varpi}. \quad (7.5.4)$$

Our goal is to prove that

$$\varpi \mid (2\mu+1). \quad (7.5.5)$$

Multiply (7.5.4) by 24 to obtain the equivalent congruence

$$12\mu(\mu+1) + 12\nu(3\nu+1) + 4\varpi + 4 \equiv 0 \pmod{\varpi},$$

or

$$3(2\mu+1)^2 + (6\nu+1)^2 \equiv 0 \pmod{\varpi}. \quad (7.5.6)$$

Using the fact that, for each prime  $p$ , the Legendre symbol  $\left(\frac{-1}{p}\right) = (-1)^{(p-1)/2}$ , and the law of quadratic reciprocity, we find that

$$\left(\frac{-3}{\varpi}\right) = \left(\frac{\varpi}{3}\right) = \left(\frac{-1}{3}\right) = -1.$$

Thus, the only way that (7.5.6) can hold is for (7.5.5) to happen. But then, from the right-hand side of (7.5.3), we can conclude that

$$p_{\delta\varpi-4} \left( n\varpi - \frac{\varpi+1}{6} \right) \equiv 0 \pmod{\varpi}.$$

Thus, the proof is complete.  $\square$

**Corollary 7.5.2** (p. 182). *For each positive integer  $n$ ,*

$$\begin{aligned} p_6(5n-1) &\equiv 0 \pmod{5}, \\ p_7(11n-2) &\equiv 0 \pmod{11}. \end{aligned}$$

**Proof.** The first congruence arises from the case  $\varpi = 5$  and  $\delta = 2$ , while the second arises from the case  $\varpi = 11$  and  $\delta = 1$  in Entry 7.5.1.  $\square$

Next, Ramanujan gives an elementary proof of the congruence  $p(7n-2) \equiv 0 \pmod{7}$ . He begins with the same first three lines of [87, eq. (13)], [90, p. 212], and then argues in a somewhat more abbreviated fashion than he does in [87] to deduce the congruence

$$p_{-6}(7n-2) \equiv 0 \pmod{49}, \tag{7.5.7}$$

from which it follows that

$$p(7n-2) \equiv 0 \pmod{7}. \tag{7.5.8}$$

It should be remarked that the stronger congruence (7.5.7) is not mentioned by Ramanujan in [87], although it is implicit in his argument.

Unfortunately, the one-page manuscript ends with (7.5.8). It would seem that Ramanujan would have next offered a theorem analogous to Entry 7.5.1, and so we shall state and prove such a theorem here, but, of course, Ramanujan probably would have had lots more to say to us, if his manuscript had survived.

**Theorem 7.5.3.** *For a prime  $\varpi$  with  $4 \mid (\varpi+1)$ , any integer  $\delta$ , and any positive integer  $n$ ,*

$$p_{\delta\varpi-6} \left( n\varpi - \frac{\varpi+1}{4} \right) \equiv 0 \pmod{\varpi}. \tag{7.5.9}$$

In the case  $\delta = 0$  above, we can strengthen (7.5.9).

**Entry 7.5.4** (p. 182). *We have*

$$p_{-6} \left( n\varpi - \frac{\varpi+1}{4} \right) \equiv 0 \pmod{\varpi^2}. \tag{7.5.10}$$

Observe that (7.5.7) is the special case  $\varpi = 7$  of (7.5.10), and so, with slight exaggeration, we affix “p. 182” to the entry above.

**Corollary 7.5.5.** *For each positive integer  $n$ ,*

$$p_{3\delta-6}(3n-1) \equiv 0 \pmod{3}. \tag{7.5.11}$$

**Proof.** Set  $\varpi = 3$  in Theorem 7.5.3.  $\square$

For the case  $\delta = 3$  in (7.5.11), N.D. Baruah and K.K. Ojah [26], using more sophisticated means, obtained the stronger result

$$p_3(3n - 1) \equiv 0 \pmod{3^2}.$$

**Proof of Theorem 7.5.3.** Consider, for  $\lambda = (\varpi + 1)/4$ ,

$$\begin{aligned} \sum_{n=0}^{\infty} p_{\delta\varpi-6}(n)q^{n+\lambda} &= (q; q)_{\infty}^{-\delta\varpi} (q; q)_{\infty}^6 q^{\lambda} \\ &\equiv (q^{\varpi}; q^{\varpi})_{\infty}^{-\delta} \sum_{\mu=0}^{\infty} \sum_{\nu=0}^{\infty} (-1)^{\mu+\nu} (2\mu+1)(2\nu+1) q^{\frac{1}{2}\mu(\mu+1) + \frac{1}{2}\nu(\nu+1) + \lambda} \pmod{\varpi}, \end{aligned} \quad (7.5.12)$$

upon the use of Jacobi's identity (3.1.31). We need to show that if

$$\frac{1}{2}\mu(\mu+1) + \frac{1}{2}\nu(\nu+1) + \frac{\varpi+1}{4} \equiv 0 \pmod{\varpi}, \quad (7.5.13)$$

then

$$\varpi^2 \mid (2\mu+1)(2\nu+1). \quad (7.5.14)$$

The congruence (7.5.9) will then follow from (7.5.14) and (7.5.12). Multiply (7.5.13) by 8 to obtain

$$4\mu(\mu+1) + 4\nu(\nu+1) + 2\varpi + 2 \equiv 0 \pmod{\varpi},$$

or

$$(2\mu+1)^2 + (2\nu+1)^2 \equiv 0 \pmod{\varpi}.$$

Since

$$\left(\frac{-1}{\varpi}\right) = -1,$$

we conclude that

$$\varpi \mid (2\mu+1) \quad \text{and} \quad \varpi \mid (2\nu+1),$$

which completes the proof of (7.5.14).  $\square$

Observe that if  $\delta = 0$ , then the congruence in (7.5.12) can be replaced by an equality. Hence, in (7.5.9), the congruence modulo  $\varpi$  can be replaced by a congruence modulo  $\varpi^2$  in view of (7.5.14). Entry 7.5.4 therefore follows.

Although Entry 7.5.1 and Theorem 7.5.3 are not special cases of the general theorem of Andrews and Roy [23], they would be instances of the general theorem envisioned by the authors in Section 5 of their paper [23].

Recall next that a corollary of Winquist's identity is given by [103]

$$48(q; q)_{\infty}^{10} = \sum_{m, n=-\infty}^{\infty} (-1)^{m+n} ((6m+3)^3(6n+1) - (6m+3)(6n+1)^3) q^{\frac{1}{2}(3m^2+3m+3n^2+n)}. \quad (7.5.15)$$



**Theorem 7.5.6.** For a prime  $\varpi$  with  $12 \mid (\varpi + 1)$ , and any integer  $\delta$ , we have

$$p_{\delta\varpi-10}\left(n\varpi - \frac{5(\varpi+1)}{12}\right) \equiv 0 \pmod{\varpi}.$$

**Proof.** Let  $\lambda = 5(\varpi + 1)/12$ , and from (7.5.15) consider

$$\begin{aligned} \sum_{n=0}^{\infty} p_{\delta\varpi-10}(n)q^{n+\lambda} &= (q; q)_{\infty}^{-\delta\varpi} (q; q)_{\infty}^{10} q^{\lambda} \\ &\equiv (q^{\varpi}; q^{\varpi})_{\infty}^{-\delta} \frac{1}{48} \sum_{m,n=-\infty}^{\infty} (-1)^{m+n} ((6m+3)^3(6n+1) \\ &\quad - (6m+3)(6n+1)^3) q^{\frac{1}{2}(3m^2+3m+3n^2+n)+\lambda} \pmod{\varpi}. \end{aligned} \tag{7.5.16}$$

If

$$\frac{1}{2}(3m^2 + 3m + 3n^2 + n) + \lambda \equiv 0 \pmod{\varpi},$$

then upon multiplying both sides above by 24, we find that

$$12(3m^2 + 3m + 3n^2 + n) + 10(\varpi + 1) \equiv 0 \pmod{\varpi},$$

or

$$(6m+3)^2 + (6n+1)^2 \equiv 0 \pmod{\varpi}.$$

Since

$$\left(\frac{-1}{\varpi}\right) = -1,$$

we see that

$$\varpi \mid (6m+3) \quad \text{and} \quad \varpi \mid (6n+1).$$

Using these observations in (7.5.16), we complete the proof.  $\square$

We observe that in the special case  $\delta = 0$ , our proof yields a stronger result.

**Corollary 7.5.7.** For a prime  $\varpi$  with  $12 \mid (\varpi + 1)$ , we have

$$p_{-10}\left(n\varpi - \frac{5(\varpi+1)}{12}\right) \equiv 0 \pmod{\varpi^4}.$$

## 7.6. Exercises

1. Use the Jacobi triple product identity to show that

$$\varphi(-q) = \frac{(q; q)_{\infty}}{(-q; q)_{\infty}}. \tag{7.6.1}$$

2. Find a proof of Theorem 7.4.5 that is in the spirit of Ramanujan's elementary proofs of the congruences  $p(5n+4) \equiv 0 \pmod{5}$  and  $p(7n+5) \equiv 0 \pmod{7}$ , in particular, that depends on the pentagonal number theorem and Jacobi's identity.

3. Give a proof of Halphen's Identity.

4. Prove (7.4.11).

Define the colored partition function  $p_m(n)$  by

$$(q; q)_\infty^m =: \sum_{n=0}^{\infty} p_m(n) q^n. \quad (7.6.2)$$

5. If  $p_5(n)$ ,  $n \geq 0$ , is defined by (7.6.2), then

$$\sum_{n=0}^{\infty} p_5(5n) q^n = \frac{(q; q)_\infty^6}{(q^5; q^5)_\infty}.$$

Hint: Write

$$\sum_{n=0}^{\infty} p_5(n) q^n = (q; q)_\infty^5 = \frac{(q^5; q^5)_\infty^6}{(q^{25}; q^{25})_\infty} \frac{(q; q)_\infty^5}{(q^{25}; q^{25})_\infty^5} \frac{(q^{25}; q^{25})_\infty^6}{(q^5; q^5)_\infty^6}.$$

6. If  $p_7(n)$ ,  $n \geq 0$ , is defined by (7.6.2), then

$$\sum_{n=0}^{\infty} p_7(7n) q^n = \frac{(q; q)_\infty^8}{(q^7; q^7)_\infty} + 49q(q; q)_\infty^4 (q^7; q^7)_\infty^3.$$

Hint: Write

$$\sum_{n=0}^{\infty} p_7(n) q^n = (q; q)_\infty^7 = \frac{(q^7; q^7)_\infty^8}{(q^{49}; q^{49})_\infty} \frac{(q; q)_\infty^7}{(q^{49}; q^{49})_\infty^7} \frac{(q^{49}; q^{49})_\infty^8}{(q^7; q^7)_\infty^8}.$$

7. Recall that

$$\varphi(q) = \sum_{j=-\infty}^{\infty} q^{j^2} \quad \text{and} \quad \psi(q) = \sum_{j=0}^{\infty} q^{j(j+1)/2}. \quad (7.6.3)$$

In each case, determine for which positive integers  $n$ ,  $\varphi^n(q)$  and  $\psi^n(q)$  are lacunary.

8. Prove (7.3.7), (7.3.9), and (7.3.12).

9. Prove that Winquist's identity (7.4.8) can be put in the form (7.4.9).

---

## Bibliography

- [1] A. K. Agarwal, Padmavathamma, and M. V. Subbarao, *Partition Theory*, Atma Ram & Sons, Chandigarh, 2005.
- [2] S. Ahlgren, *Distribution of parity of the partition function in arithmetic progressions*, Indag. Math., N. S. **10** (1999), 173–181.
- [3] S. Ahlgren and M. Boylan, *Arithmetic properties of the partition function*, Invent. Math. **153** (2003), 487–502.
- [4] H. L. Alder, *Research Problems*, Bull. Amer. Math. Soc. **62** (1956), 76.
- [5] C. Alfes, M. Jameson, and R. Lemke Oliver, *Proof of the Alder–Andrews conjecture*, Proc. Amer. Math. Soc. **139** (2011), 63–78.
- [6] G. E. Andrews, *A generalization of the Göllnitz–Gordon partition theorems*, Proc. Amer. Math. Soc. **18** (1967), 945–952.
- [7] G. E. Andrews, *On a partition problem of H. L. Alder*, Pacific. J. Math. **36** (1971), 279–284.
- [8] G. E. Andrews, *On the general Rogers–Ramanujan theorem*, Memoirs Amer. Math. Soc., No. 152 (1974).
- [9] G. E. Andrews, *On identities implying the Rogers–Ramanujan identities*, Houston Math. J. **2** (1976), 289–298.
- [10] G. E. Andrews, *Connection coefficient problems and partitions*, in *Proc. Sym. Pure Math.*, No. 34, American Mathematical Society, 1979, 1–24
- [11] G. E. Andrews, *The Theory of Partitions*, Addison–Wesley, Reading, MA, 1976; reissued: Cambridge University Press, Cambridge, 1998.
- [12] G. E. Andrews, *Generalized Frobenius Partitions*, Mem. Amer. Math. Soc., No. 301. **49** (1984), American Mathematical Society, Providence, RI, 1984.
- [13] G. E. Andrews, *J. J. Sylvester, Johns Hopkins and partitions*, in *A Century of Mathematics in America*, Part I, P. Duren, ed., American Mathematical Society, Providence, RI, 1988, pp. 21–40.
- [14] G. E. Andrews, *On the proofs of the Rogers–Ramanujan identities*, in *q-Series and Partitions*, D. Stanton, ed., Springer-Verlag, New York, 1989, pp. 1–14.
- [15] G. E. Andrews, *An identity of Sylvester and the Rogers–Ramanujan identities*, in *Number Theory, Madras 1987*, Lecture Notes in Math. 1395, Springer-Verlag, Berlin, 1989, pp. 64–72.

- [16] G. E. Andrews, *A Fine dream*, Internat. J. Number Thy. **3** (2007), 325–334.
- [17] G. E. Andrews and B. C. Berndt, *Ramanujan's Lost Notebook, Part I*, Springer, New York, 2005.
- [18] G. E. Andrews and B. C. Berndt, *Ramanujan's Lost Notebook, Part II*, Springer, New York, 2009.
- [19] G. E. Andrews and B. C. Berndt, *Ramanujan's Lost Notebook, Part III*, Springer, New York, 2013.
- [20] G. E. Andrews and K. Eriksson, *Integer Partitions*, Cambridge University Press, Cambridge, 2004.
- [21] G. E. Andrews and F. G. Garvan, *Dyson's crank of a partition*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), 167–171.
- [22] G. E. Andrews, R. Lewis, and Z.-G. Liu, *An identity relating a theta function to a sum of Lambert series*, Bull. London Math. Soc. **33** (2001), 25–31.
- [23] G. E. Andrews and R. Roy, *Ramanujan's method in  $q$ -series congruences*, Elec. J. Comb. **4**(2): R2 (1997), 7 pp.
- [24] A. O. L. Atkin, *Proof of a conjecture of Ramanujan*, Glasgow Math. J. **8** (1967), 14–32.
- [25] A. O. L. Atkin and H. P. F. Swinnerton-Dyer, *Some properties of partitions*, Proc. London Math. Soc. (3) **4** (1954), 84–106.
- [26] N. D. Baruah and K. K. Ojah, *Some congruences deducible from Ramanujan's cubic continued fraction*, Internat. J. Number Thy. **7** (2011), 1331–1343.
- [27] B. C. Berndt, *Ramanujan's Notebooks, Part III*, Springer-Verlag, New York, 1991.
- [28] B. C. Berndt, *Ramanujan's Notebooks, Part V*, Springer-Verlag, New York, 1998.
- [29] B. C. Berndt, *Number Theory in the Spirit of Ramanujan*, American Mathematical Society, Providence, RI, 2006.
- [30] B. C. Berndt, *What is a  $q$ -series?*, in *Ramanujan Rediscovered: Proceedings of a Conference on Elliptic Functions, Partitions, and  $q$ -Series in memory of K. Venkatchaliengar: Bangalore, 1 – 5 June, 2009*, N. D. Baruah, B. C. Berndt, S. Cooper, T. Huber, and M. J. Schlosser, eds., Ramanujan Mathematical Society, Mysore, 2010, pp. 31–51.
- [31] B. C. Berndt, H. H. Chan, S. H. Chan, and W.-C. Liaw, *Cranks and dissections in Ramanujan's lost notebook*, J. Combin. Thy., Ser. A **109** (2005), 91–120.
- [32] B. C. Berndt, H. H. Chan, S. H. Chan, and W.-C. Liaw, *Ramanujan and cranks*, in *Theory and Applications of Special Functions. A Volume Dedicated to Mizan Rahman*, M. E. H. Ismail and E. Koelink, eds., Springer, New York, 2005, pp. 77–98.
- [33] B. C. Berndt, H. H. Chan, S. H. Chan, and W.-C. Liaw, *Cranks—really the final problem*, Ramanujan J. **23** (2010), 3–15.
- [34] B. C. Berndt, S. H. Chan, Z.-G. Liu, and H. Yesilyurt, *A new identity for  $(q; q)_{\infty}^{10}$  with an application to Ramanujan's partition congruence modulo 11*, Quart. J. Math. (Oxford) **55** (2004), 13–30.
- [35] B. C. Berndt, C. Gugg, and S. Kim, *Ramanujan's elementary method in partition congruences*, in *Partitions,  $q$ -Series and Modular Forms*, K. Alladi and F. Garvan, eds., Developments in Math. **23**, Springer, New York, 2011, pp. 13–22.
- [36] B. C. Berndt and K. Ono, *Ramanujan's unpublished manuscript on the partition and tau functions with proofs and commentary*, Sém. Lotharingien de Combinatoire **42** (1999), 63 pp.; in *The Andrews Festschrift*, D. Foata and G.-N. Han, eds., Springer-Verlag, Berlin, 2001, pp. 39–110.

- [37] B. C. Berndt and R. A. Rankin, *Ramanujan: Letters and Commentary*, American Mathematical Society, Providence, RI, 1995; London Mathematical Society, London, 1995.
- [38] B. C. Berndt, A. J. Yee, and J. Yi, *Theorems on partitions from a page in Ramanujan's lost notebook*, J. Comp. Appl. Math. **160** (2003), 53–68.
- [39] B. C. Berndt, A. J. Yee, and A. Zaharescu, *On the parity of partition functions*, Inter. J. Math. **14** (2003), 437–459.
- [40] B. C. Berndt, A. J. Yee, and A. Zaharescu, *New theorems on the parity of partition functions*, J. Reine Angew. Math. **566** (2004), 91–109.
- [41] L. Carlitz, *Rectangular arrays and plane partitions*, Acta Arith. **13** (1967), 29–67.
- [42] H.-C. Chan, *Ramanujan's cubic continued fraction and an analog of his "most beautiful identity,"* Internat. J. Number Thy. **6** (2010), 673–680.
- [43] H.-C. Chan, *Ramanujan's cubic continued fraction and Ramanujan type congruences for a certain partition function*, Internat. J. Number Thy. **6** (2010), 819–834.
- [44] H.-C. Chan, *Distribution of a certain partition function modulo powers of primes*, Acta Math. Sin. (English Ser.) **27** (2011), 625–634.
- [45] O.-Y. Chan, *Some asymptotics for cranks*, Acta Arith. **120** (2005), 107–143.
- [46] S. H. Chan, *Generalized Lambert series identities*, Proc. London Math. Soc. (3) **91** (2005), 598–622.
- [47] S. D. Chowla, *Congruence properties of partitions*, J. London Math. Soc. **9** (1934), 247.
- [48] W. Chu and L. Di Claudio, *Classical Partition Identities and Basic Hypergeometric Series*, Quaderno 6/2004 del Dipartimento di Matematica "Ennio De Giorgi", Università degli Studi di Lecce, 2004.
- [49] F. J. Dyson, *Some guesses in the theory of partitions*, Eureka (Cambridge) **8** (1944), 10–15.
- [50] D. Eichhorn, *A new lower bound on the number of odd values of the ordinary partition function*, Ann. Comb. **13** (2009), 297–303.
- [51] A. B. Ekin, *Inequalities for the crank*, J. Combin. Thy. Ser. A **83** (1998), 283–289.
- [52] A. B. Ekin, *Some properties of partitions in terms of crank*, Trans. Amer. Math. Soc. **352** (2000), 2145–2156.
- [53] J. Fabrykowski and M. V. Subbarao, *Some new identities involving the partition function  $p(n)$* , in *Number Theory*, R. A. Mollin, ed., Walter de Gruyter, New York, 1990, pp. 125–138.
- [54] N. J. Fine, *Some new results on partitions*, Proc. Nat. Acad. Sci. U.S.A. **34**, 616–618.
- [55] F. G. Garvan, *A simple proof of Watson's partition congruences for powers of 7*, J. Austral. Math. Soc. (Series A) **36** (1984), 316–334.
- [56] F. G. Garvan, *Generalizations of Dyson's Rank*, Ph. D. Thesis, Pennsylvania State University, University Park, PA, 1986.
- [57] F. G. Garvan, *New combinatorial interpretations of Ramanujan's partition congruences mod 5, 7 and 11*, Trans. Amer. Math. Soc. **305** (1988), 47–77.
- [58] F. G. Garvan, *The crank of partitions mod 8, 9 and 10*, Trans. Amer. Math. Soc. **322** (1990), 79–94.
- [59] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd ed., Cambridge University Press, Cambridge, 2004.
- [60] H. Göllnitz, *Einfache Partitionen*, Diplomarbeit W.S., Göttingen, 1960 (unpublished).
- [61] H. Göllnitz, *Partitionen mit Differenzenbedingungen*, J. Reine Angew. Math. **225** (1967), 154–190.

- [62] B. Gordon, *Some Ramanujan-like continued fractions*, Abstracts of Short Communications, International Congress of Mathematicians, Stockholm, 1962, pp. 29–30.
- [63] B. Gordon, *Some continued fractions of the Rogers–Ramanujan type*, Duke Math. J. **32** (1965), 741–748.
- [64] H. Gupta, *A table of partitions*, Proc. London Math. Soc. **39** (1935), 47–53.
- [65] H. Gupta, *Collected Works of Hansraj Gupta*, Vol. 1, R. J. Hans–Gill and M. Raka, eds., Ramanujan Mathematical Society, Mysore, India, 2013.
- [66] G. H. Halphen, *Traité des fonctions elliptiques et de leurs applications vol. 1*, Gauthier-Villars, Paris, 1888.
- [67] G. H. Hardy, *Ramanujan: Twelve Lectures on Subjects Suggested by His Life and Work*, Cambridge University Press, Cambridge, 1940; reprinted by Chelsea, New York, 1960; reprinted by the American Mathematical Society, Providence, RI, 1999.
- [68] E. Heine, *Untersuchungen über die Reihe*  

$$1 + \frac{(1-q^\alpha)(1-q^\beta)}{(1-q)(1-q^\gamma)} \cdot x + \frac{(1-q^\alpha)(1-q^{\alpha+1})(1-q^\beta)(1-q^{\beta+1})}{(1-q)(1-q^2)(1-q^\gamma)(1-q^{\gamma+1})} \cdot x^2 + \dots,$$
 J. Reine Angew. Math. **34** (1847), 285–328.
- [69] M. D. Hirschhorn, *A birthday present for Ramanujan*, Amer. Math. Monthly **97** (1990), 398–400.
- [70] M. D. Hirschhorn, Lecture at Conference, *Number Theory, Partitions, q-Series, and Related Research*, National Institute of Education, Singapore, March 7, 2014.
- [71] M. D. Hirschhorn and D. C. Hunt, *A simple proof of the Ramanujan conjecture for powers of 5*, J. Reine Angew. Math. **336** (1981), 1–17.
- [72] B. Kim, *An analog of crank for a certain kind of partition function arising from the cubic continued fraction*, Acta Arith. **148** (2011), 1–19.
- [73] S. Kim, *A combinatorial proof of a recurrence relation for the partition function due to Euler*, preprint.
- [74] M. I. Knopp, *Modular Functions in Analytic Number Theory*, Chelsea, New York, 1993.
- [75] O. Kolberg, *Note on the parity of the partition function*, Math. Scand. **7** (1959), 377–378.
- [76] L. Lorentzen and H. Waadeland, *Continued Fractions with Applications*, North Holland, Amsterdam, 1992.
- [77] P. A. MacMahon, *Combinatory Analysis* (2 volumes), Cambridge University Press, London, 1916; reprinted by Chelsea, New York, 1960.
- [78] L. Mirsky, *The distribution of values of the partition function in residue classes*, J. Math. Anal. Appl. **93** (1983), 593–598.
- [79] M. Newman, *Congruences for the coefficients of modular forms and some new congruences for the partition function*, Canad. J. Math. **9** (1957), 549–552.
- [80] J.–L. Nicolas and A. Sárközy, *On the parity of partition functions*, Illinois J. Math. **39** (1995), 586–597.
- [81] J.–L. Nicolas, I. Z. Ruzsa and A. Sárközy, *On the parity of additive representation functions. With an appendix by J.–P. Serre*, J. Number Thy. **73** (1998), 292–317.
- [82] J.–L. Nicolas, *Valeurs impaires de la fonction de partition  $p(n)$* , Internat. J. Number Thy. **2** (2004), 469–487.
- [83] T. R. Parkin and D. Shanks, *On the distribution of parity in the partition function*, Math. Comp. **21** (1967), 466–480.

- [84] K. G. Ramanathan, *Ramanujan and the congruence properties of partitions*, Proc. Indian Acad. Sci. (Math. Sci.) **89** (1980), 133–157.
- [85] S. Ramanujan, *Highly composite numbers*, Proc. London Math. Soc. **14** (1915), 347–409.
- [86] S. Ramanujan, *Proof of certain identities in combinatory analysis*, Proc. Cambridge Philos. Soc. **19** (1919), 214–216.
- [87] S. Ramanujan, *Some properties of  $p(n)$ , the number of partitions of  $n$* , Proc. Cambridge Philos. Soc. **19** (1919), 210–213.
- [88] S. Ramanujan, *Congruence properties of partitions*, Proc. London Math. Soc. **18** (1920), records for 13 March 1919, xix.
- [89] S. Ramanujan, *Congruence properties of partitions*, Math. Z. **9** (1921), 147–153.
- [90] S. Ramanujan, *Collected Papers*, Cambridge University Press, Cambridge, 1927; reprinted by Chelsea, New York, 1962; reprinted by the American Mathematical Society, Providence, RI, 2000.
- [91] S. Ramanujan, *Notebooks* (2 volumes), Tata Institute of Fundamental Research, Bombay, 1957.
- [92] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Narosa, New Delhi, 1988.
- [93] Z. Reti, *Five Problems in Combinatorial Number Theory*, Ph.D. Thesis, University of Florida, Gainesville, FL, 1994.
- [94] L. J. Rogers, *Second memoir on the expansion of certain infinite products*, Proc. London Math. Soc. **25** (1894), 318–343.
- [95] I. Schur, *Zur additiven Zahlentheorie*, Sitz. Preuss. Akad. Wiss. Phys.–Math. Kl. **1926**, 488–495.
- [96] I. Schur, *Ein Beitrag zur additiven Zahlentheorie und zur Theorie der Kettenbrüche*, S.–B. Preuss. Akad. Wiss. Phys.–Math. Kl. (1917), 302–321.
- [97] J.–P. Serre, *Sur la lacunarité des puissances de  $\eta$* , Glasgow Math. J. **27** (1985), 203–221.
- [98] L. J. Slater, *Further identities of the Rogers–Ramanujan type*, Proc. London Math. Soc. **54** (1952), 147–167.
- [99] J. J. Sylvester, *A constructive theory of partitions, arranged in three acts, an interact, and an exodion*, in *The Collected Papers of J. J. Sylvester*, Vol. 3, Cambridge University Press, London, pp. 1–83; reprinted by Chelsea, New York, 1973.
- [100] G. N. Watson, *Ramanujans Vermutung über Zerfallungsanzahlen*, J. Reine Angew. Math. **179** (1938), 97–128.
- [101] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, 4th ed., Cambridge University Press, Cambridge, 1966.
- [102] S. Wigert, *Sur l’ordre de grandeur du nombre des diviseurs d’un entier*, Ark. Mat. Astron. Fys. **3** (1906–1907), 1–9.
- [103] L. Winquist, *An elementary proof of  $p(11n + 6) \equiv 0 \pmod{11}$* , J. Comb. Thy. **6** (1969), 56–59.
- [104] E. M. Wright, *An enumerative proof of an identity of Jacobi*, J. London Math. Soc. **40** (1965), 55–57.
- [105] A. J. Yee, *Partitions with difference conditions and Alder’s conjecture*, Proc. Nat. Acad. Sci. **101** (2004), 16417–16418.
- [106] A. J. Yee, *Alder’s conjecture*, J. Reine Angew. Math. **616** (2008), 67–88.
- [107] A. Yong, Critique of Hirsch’s citation index: A combinatorial Fermi problem, preprint.