

UNNORMALIZED DIFFERENCES BETWEEN ZEROS OF L -FUNCTIONS

KEVIN FORD, ALEXANDRU ZAHARESCU

ABSTRACT. We study a subtle inequity in the distribution of unnormalized differences between imaginary parts of zeros of the Riemann zeta function, which was observed by a number of authors. We establish a precise measure which explains the phenomenon, that the location of each Riemann zero is encoded in the distribution of large Riemann zeros. We also extend these results to zeros of more general L -functions. In particular, we show how the rank of an elliptic curve over \mathbb{Q} is encoded in the sequences of zeros of *other* L -functions, not only the one associated to the curve.

1 Introduction

The study of local spacing distribution of sequences was initiated by physicists (see Wigner [38] and Dyson [9]) in order to understand the spectra of high energies. These notions have received a great deal of attention in many areas of mathematical physics, analysis, probability theory and number theory. After the pioneering work of Montgomery [25] on the pair correlation of the imaginary parts of zeros of the Riemann zeta function, followed by Hejhal's investigation [14] of the triple correlation, higher level correlations for general L -functions have been studied by Rudnick and Sarnak [30], and Katz and Sarnak [19], [20]. In particular, in the case of the Riemann zeta function, the above mentioned works establish the existence for all m of the limiting m -level correlations, for certain large classes of test functions, and confirm that the limiting correlations are the same as those from the Gaussian unitary ensemble (GUE) model. One important feature of the GUE model is that the density function vanishes at the origin. This means strong repulsion between consecutive zeros of the Riemann zeta function (see also supporting numerical computations by Odlyzko [26]).

More recently, the distribution of *unnormalized* differences between zeros has been considered by a number of researchers. Motivated in part by attempts to understand the limitations of the predictions of random matrix theory, Bogomolny and Keating [3] were the first to make a precise conjecture for the discrepancy of such differences; see also the survey articles of Berry-Keating [1] and Snaith [36]. In particular, as pointed out in [1], there seems to be a striking resurgence phenomenon: in the pair correlation of high Riemann zeros, the low Riemann zeros appear as resonances; see also [36, Fig. 3] showing the gaps between the first 100,000 Riemann zeros.

In 2011, Pérez-Marco [23] performed extensive numerical studies of the distribution of differences of zeros of the Riemann zeta function and of other L -functions. He also highlighted an interesting discrepancy phenomenon for zeros of $\zeta(s)$, namely that the differences tend to avoid imaginary parts of small zeros, e.g. there is a deficiency of differences near $\gamma_1 = 14.1347\dots$, the imaginary part of the smallest nontrivial zero of ζ . One may interpret this as saying that there is some repulsion between the imaginary parts of zeros of $\zeta(s)$, and the translates of these imaginary parts by γ_1 . The reader is referred to [23] for extensive data supporting this phenomenon, for $\zeta(s)$ as well as for more general L -functions.

In the present paper, our goal is to prove, unconditionally, a precise measure of the discrepancy of gaps in the distribution of unnormalized differences between zeros. To proceed, let $\hat{h}(\xi) = \int h(x)e^{-2\pi i x \xi} dx$ denote the Fourier transform of h . We will consider $h \in \mathcal{H}$, the class of functions h satisfying

Date: October 3, 2014.

Key words and phrases. Riemann zeta function, L -functions, distribution of zeros.

The first author was supported in part by National Science Foundation grant DMS-1201442.

2010 Mathematics Subject Classification: Primary 11M26; Secondary 11K38.

- (1) $\int h = 0$,
- (2) $\int |xh(x)| dx < \infty$
- (3) $|\hat{h}'(\xi)| \ll |\xi|^{-5}$ for $|\xi| \geq 1$.

Let \mathcal{H}_0 be the set of functions $h \in \mathcal{H}$ such that in addition, 0 is not in the closure of the support of h .

Theorem 1. *Suppose $h \in \mathcal{H}_0$. For $T \geq 2$,*

$$\sum_{0 < \gamma, \gamma' \leq T} h(\gamma - \gamma') = \frac{T}{2\pi} \int_{-\infty}^{\infty} h(t)(K(1+it) + K(1-it)) dt + O\left(\frac{T}{(\log T)^{1/3}}\right),$$

where $K(s)$ is defined as

$$K(s) = \sum_{n=1}^{\infty} \frac{\Lambda^2(n)}{n^s} \quad (\Re s > 1),$$

and by analytic continuation to $\{s : \Re s \geq 1, s \neq 1\}$. Here γ, γ' are imaginary parts of nontrivial zeros of $\zeta(s)$, each zero occurring in the sum the number of times of its multiplicity.

We mention here that the conclusion of Theorem 1, with suitable test functions h , has been obtained by Conrey and Snaith [6] under the assumption of the L -functions ratios conjecture of Conrey, Farmer and Zirnbauer [5]. More recently, the conclusion of Theorem 1 was deduced, for test functions h with compactly supported Fourier transforms and under the assumption of the Riemann Hypothesis, by Rodgers [29, Theorem 1.4].

Remarks. (i) The constant implied by the O -symbol depends on h . By standard counts of zeros (see (2.1) and (2.2) below), if h has compact support then there are $\asymp T \log^2 T$ nonzero summands $h(\gamma - \gamma')$. Thus, Theorem 1 implies that there is a discrepancy in the distribution of $\gamma - \gamma'$ of relative order $1/\log^2 T$. The discrepancy has a “density function” $K(1+it) + K(1-it)$, which is graphed in Figure 1.

(ii) $K(s)$ is close to the “nice” (from an analytic point of view) function

$$\left(\frac{\zeta'}{\zeta}\right)'(s) = \sum_{n=1}^{\infty} \frac{\Lambda(n) \log n}{n^s} \quad (\Re s > 1).$$

Let $b_m = \mu(\ker(m))\phi(\ker(m))$, where $\ker(m) = \prod_{p|m} p$. Using the identity $1 = \sum_{m|k} (k/m)b_m$, we obtain the meromorphic continuation

$$K(s) = \sum_{m=1}^{\infty} b_m \left(\frac{\zeta'}{\zeta}\right)'(ms) \quad (\Re s > 0),$$

provided that $ms \neq 1$ and $\zeta(ms) \neq 0$ for all $m \in \mathbb{N}$, the sum on m converges absolutely, since $|b_m| \leq m$ and $(\zeta'/\zeta)'(\sigma+it) = O(2^{-\sigma})$ for $\sigma \geq 2$.

Invoking the explicit formula for $(\zeta'/\zeta)(s)$ [7, §12, (8)] one gets, for any fixed $M \geq 4$ and $\sigma = \Re s > \frac{1}{2}$,

$$(1.1) \quad K(s) = \sum_{m=1}^M \frac{b_m}{m^2} \left(\frac{1}{(s-1/m)^2} - \sum_{\rho}^* \frac{1}{(s-\rho/m)^2} \right) + O(M2^{-\sigma M})$$

where \sum_{ρ}^* denotes a sum over *all* zeros of ζ , including the trivial zeros.

The terms in (1.1) corresponding to $m=1$ are the most significant. Here $b_1 = 1$, and we see that if $s = 1+it$ and $t \approx \Im \rho$ for some ρ , the term in (1.1) corresponding to ρ will be

$$-\frac{b_m}{m^2} \frac{1}{(s-\rho/m)^2} \approx -\frac{1}{(1-1/2)^2} = -4.$$

This accounts for the noticeable dips of about -8 in magnitude on the graph of $K(1+it) + K(1-it)$ (Figure 1) corresponding to low-lying zeros of $\zeta(s)$ (imaginary parts $\gamma_1 = 14.134 \dots, \gamma_2 = 21.022 \dots, \dots$). Likewise, for small $m > 1$, if $t \approx \Im \rho/m$ for some ρ there will be a large positive or negative (depending on the sign of b_m) term in (1.1) corresponding to ρ . In particular, if $t \approx \Im \rho/2$, the term in (1.1) corresponding to




FIGURE 1. Plot of $K(1 + it) + K(1 - it)$ together with small zeros of $\zeta(s)$

ρ is about $-\frac{b_2}{2^2} \frac{1}{(1-1/4)^2} = \frac{4}{9}$, and if $t \approx \Im \rho/3$, the term corresponding to ρ is about $\frac{8}{25}$. We have highlighted a few values of t in Figure 1 where terms corresponding to $m = 2$ and $m = 3$ produce noticeable peaks in the graph.

It is not difficult to generalize Theorem 1 to the study of differences of imaginary parts of zeros of a large class of L -functions. Generally speaking, an L -function is a Dirichlet series $F(s) = \sum_{n=1}^{\infty} a_F(n)n^{-s}$ whose coefficients possess both *multiplicative structure* (that is, F has an Euler product) and *additive structure* (encoded in a functional equation equation for F). One framework in which to operate is the Selberg Class \mathcal{S} , the set of F satisfying the following axioms:

- (i) there exists an integer $m \geq 0$ such that $(s - 1)^m F(s)$ is an entire function of finite order;
- (ii) F satisfies a functional equation

$$(1.2) \quad \Phi(s) = w\overline{\Phi}(1-s), \quad \Phi(s) = Q^s \prod_{j=1}^r \Gamma(\lambda_j s + \mu_j) F(s)$$

with $Q > 0$, $\lambda_j > 0$, $\mu_j \in \mathbb{C}$, $\Re(\mu_j) \geq 0$ and $|w| = 1$. (Here, $\overline{f}(s) = \overline{f(\overline{s})}$);

- (iii) $F(s)$ has an Euler product ($a_F(n)$ is multiplicative), which we write as

$$-\frac{F'}{F}(s) = \sum_{n=1}^{\infty} \Lambda_F(n)n^{-s},$$

where $\Lambda_F(n)$ is supported on powers of primes;

- (iv) $\Lambda_F(n) \ll n^{\theta_F}$ for some $\theta_F < \frac{1}{2}$; and
- (v) for every $\varepsilon > 0$, $a_F(n) \ll_{\varepsilon} n^{\varepsilon}$.

The functional equation is not uniquely determined in light of the duplication formula for Γ -function, however the real sum $d_F = 2 \sum_{j=1}^r \lambda_j$ is well-defined and is known as the degree of F .

For an introduction to results and conjectures concerning the Selberg class, the reader may consult the survey papers of Kaczorowski and Perelli [18], Kaczorowski [17] and Perelli [27, 28]. In particular, \mathcal{S} includes the Riemann zeta function, Dirichlet L -functions, as well as L -functions attached to number fields, elliptic curves, and holomorphic cusp forms. Subject to the truth of the open Ramanujan-Petersson conjecture (condition (v) above), the Selberg class also includes primitive Maass cusp forms, and more generally,

all automorphic L -functions (in the sense of Godement-Jacquet). Chapter 5 of the book of Iwaniec and Kowalski [15] is another excellent reference for the basic theory of L -functions.

By convention, all sums, products and counts of zeros of a function $F \in \mathcal{S}$ count zeros with their multiplicity. A generic zero will be written as $\rho = \beta + i\gamma$ (the variable with or without subscripts), with $\beta, \gamma \in \mathbb{R}$. We denote $Z(F)$ the multi-set of nontrivial zeros (those with $0 < \beta < 1$). All sums over zeros will include only nontrivial zeros unless otherwise noted.

To generalize Theorem 1, we will require two additional hypotheses on F , a mild Mertens-type estimate for $|\Lambda_F(n)^2|$ and a zero-density estimate for F near $\Re s = \frac{1}{2}$. It is expected that all zeros in $Z(F)$ have real part $\frac{1}{2}$ (the analog of Riemann's Hypothesis for $\zeta(s)$).

(vi) We have

$$\sum_{n \leq x} \frac{|\Lambda_F(n)|^2}{n} \ll \log^2 x;$$

(vii) There exist constants $A(F) > 0, B(F) > 0$ such that

$$N_F(\sigma, T) := |\{\rho = \beta + i\gamma \in Z(F) : \beta \geq \sigma, 0 < \gamma \leq T\}| \ll T^{1-A(F)(\sigma-1/2)} (\log T)^{B(F)},$$

uniformly for $\sigma \geq 1/2$ and $T \geq 2$.

Let \mathcal{S}^* denote the set of $F \in \mathcal{S}$ satisfying (vi) and (vii) above. Condition (vi) is very mild and holds in practice, in particular for all Dirichlet L -functions (trivially from the bound $|\Lambda(n)| \leq \log n$) and also for all automorphic L -functions (see Theorem 3 below). Condition (vii) is known for the Riemann zeta function and Dirichlet L -functions (Selberg [31], [32] with $B(F) = 1$), and degree 2 L -functions attached to newforms on the full modular group (Luo [22]; also with $B(F) = 1$). As we will see later in Section 7, the method of Luo also can be used to show (vii) for newforms attached to congruence subgroups.

Theorem 2. *Let $F, G \in \mathcal{S}^*$ and suppose further that the Dirichlet series*

$$K_{F,G}(s) = \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n^s}$$

can be analytically continued to the region $\{s : \Re s \geq 1, s \neq 1\}$. For any $h \in \mathcal{H}_0$ with compact support,

$$(1.3) \quad \sum_{\substack{0 < \gamma \leq T \\ \rho \in Z(F)}} \sum_{\substack{0 < \gamma' \leq T \\ \rho' \in Z(G)}} h(\gamma - \gamma') = \frac{T}{\pi} \Re \int_{-\infty}^{\infty} h(t) K_{F,G}(1+it) dt + O\left(\frac{T}{(\log T)^{1/3}}\right).$$

Furthermore, if $K_{F,G}$ is analytic at $s = 1$, then (1.3) holds also for all $h \in \mathcal{H}$ with compact support.

Remarks. The hypothesis that h has compact support may be omitted, provided that the function $K_{F,G}$ satisfies $|K_{F,G}(\sigma + it)| \ll 1 + |t|$ uniformly for $\sigma \geq 1$, (excluding $|t| \ll 1$ if $K_{F,G}$ has a pole at $s = 1$). See the proof of (4.11) at the end of Section 4. This is the case, e.g. if F and G are both Dirichlet L -functions. In general, in order to ensure that the integral (1.3) converges absolutely, one needs a hypothesis on the decay of $h(t)$ as $|t| \rightarrow \infty$ and/or a hypothesis on the growth of $K_{F,G}(\sigma + it)$ for $\sigma + it$ as $|t| \rightarrow \infty$.

In general, the function $K_{F,G}(s)$ should satisfy the hypotheses of Theorem 2, although it is not known to be true for every $F, G \in \mathcal{S}$. The function $K_{F,G}$ is closely related to the Rankin-Selberg convolution of the L -functions F and G . In the special case where χ and ψ are Dirichet characters (principal characters allowed), $F(s) = L(s, \chi)$, and $G(s) = L(s, \psi)$, we have $\Lambda_F(n) \overline{\Lambda_G(n)} = \chi(n) \overline{\psi(n)} \Lambda^2(n)$ and hence

$$K_{F,G}(s) = \frac{d^2}{ds^2} \log L(s, \chi \overline{\psi}) + \tilde{K}(s),$$

where $\tilde{K}(s)$ is a Dirichlet series, absolutely convergent and analytic for $\Re s > 1/2$. By classical arguments [7, §14, main theorem], $L(s, \chi \overline{\psi}) \neq 0$ for $\Re s \geq 1$ and thus $K_{F,G}(s)$ is analytic on $\{s : \Re s \geq 1, s \neq 1\}$. An elaboration of this argument gives the following, which we will prove later in Section 5. The proof is based on properties of Rankin-Selberg convolution L -functions.

Theorem 3. (a) Condition (vi) holds for any automorphic L -function F ; (b) Suppose F and G are automorphic L -functions, of arbitrary degree. Then $K_{F,G}$ is analytic on $\{s : \Re s \geq 1, s \neq 1\}$. Furthermore, $K_{F,G}$ is analytic at $s = 1$ unless $F = G$.

It is natural to ask about more general linear combinations of zeros in (1.3). Let α be an arbitrary positive real number. Our method of proof of Theorem 2 indeed yields the more general result that

$$\sum_{\substack{0 < \gamma \leq T \\ \rho \in Z(F)}} \sum_{\substack{0 < \gamma' \leq T \\ \rho' \in Z(G)}} h(\gamma - \alpha\gamma') = \frac{T}{\pi} \Re \int_{\mathbb{R}} h(t) K_{F,G}^{(\alpha)}(1 + it) dt + O\left(\frac{T}{(\log T)^{1/3}}\right),$$

where

$$K_{F,G}^{(\alpha)}(s) = \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n^\alpha)}}{n^s} \quad (\Re s > 1).$$

Here $\Lambda_G(x) = 0$ unless $x \in \mathbb{N}$. We observe that the sum on n is empty unless $\alpha = \frac{\log a}{\log b}$ for some positive integers a, b , and that $K_{F,G}^{(\alpha)}$ is trivially analytic for $\Re s = 1$ if α is irrational.

We emphasize that the local minima of $K(1 + it) + K(1 - it)$ (see Figure 1) are not necessarily attained exactly at $t = \gamma_j$. However, statistics on gaps between zeros are capable of identifying the exact location of every zero. This comes from working with higher derivatives of $K(s)$, in which the influence of each zero is sharpened. This is confirmed by our next theorem, which shows that the Riemann Hypothesis for the Riemann zeta function is encoded in the (distribution of) zeros of every single Dirichlet L -function.

Theorem 4. Fix an integer $q > 1$ and a primitive Dirichlet character $\chi \pmod{q}$. Fix a compactly supported function g in $C^\infty(\mathbb{R})$ such that $g \geq 0$ and $\int g = 1$. Fix a real number $\beta > \frac{1}{2}$. Then, for any positive integer k , the function $f_{\chi,g,\beta,k} : (\frac{1}{2}, \infty) \rightarrow \mathbb{R}$ given by

$$f_{\chi,g,\beta,k}(\alpha) := \frac{(-1)^{k+1}}{2^{k+1}(k+1)!} \lim_{m \rightarrow \infty} \lim_{T \rightarrow \infty} \frac{2\pi}{T} \sum_{\substack{0 < \gamma, \gamma' \leq T \\ L(\rho, \chi) = L(\rho', \chi) = 0}} h_m^{(k)}(\gamma - \gamma'),$$

is well defined, where

$$h_m(x) := m \left(g(m(x - \alpha)) - g(m(x - \beta)) \right).$$

Moreover, the following are equivalent:

- (a) The sequence of functions $(f_{\chi,g,\beta,k})_{k \geq 1}$ converges (pointwise) on $(\frac{1}{2}, \infty)$,
- (b) The Riemann Hypothesis holds true for $\zeta(s)$.

More precisely, assuming the Riemann Hypothesis for $\zeta(s)$, for any $\beta > \alpha > 1/2$,

$$\lim_{k \rightarrow \infty} f_{\chi,g,\beta,k}(\alpha) = m_\zeta \left(\frac{1}{2} + i\alpha \right) - m_\zeta \left(\frac{1}{2} + i\beta \right),$$

where $m_\zeta(\rho)$ denotes the multiplicity of the zero of the Riemann zeta function at ρ .

Theorem 4 is an instance where a property involving the distribution of zeros of an L -function is proved to be equivalent to the Riemann Hypothesis for another L -function. One can use the methods employed to extend Theorem 1 to Theorem 2 in order to obtain a generalization of Theorem 4. We discuss only one such application here, to illustrate the ideas further.

For two general L -functions $F, G \in \mathcal{S}$ we define

$$f_{F,G,g,\beta,k}(\alpha) := \frac{(-1)^{k+1}}{2^{k+1}(k+1)!} \lim_{m \rightarrow \infty} \lim_{T \rightarrow \infty} \frac{2\pi}{T} \sum_{\substack{0 < \gamma, \gamma' \leq T \\ F(\rho) = G(\rho') = 0}} h_m^{(k)}(\gamma - \gamma'),$$

where g and h_m are the same as in Theorem 4.

Let E be an elliptic curve over the rational field \mathbb{Q} with discriminant Δ and conductor N . For squarefree d , let E_d be the quadratic twist of E ; e.g., if E is given by the equation

$$E : y^2 = x^3 + ax + b,$$

then E_d is given by

$$E_d : dy^2 = x^3 + ax + b.$$

Let $L(s, E)$ be the (normalized) L -function attached to E . By Hasse's theorem, we have

$$(1.4) \quad L(s, E) = \sum_{n=1}^{\infty} a(n)n^{-s} = \prod_p (1 - \alpha_p p^{-s})^{-1} (1 - \beta_p p^{-s})^{-1} \quad (\Re s > 1),$$

where $|\alpha_p| = |\beta_p| = 1$ and $\alpha_p = \overline{\beta_p}$ for $p \nmid \Delta$, and for $p|\Delta$, $0 \leq \alpha_p \leq 1$ and $\beta_p = 0$. That $L(s, E)$ lies in the Selberg class (modularity) is a celebrated result of Wiles et al [39], [37], [4]. In fact, $L(s, E)$ coincides with the L -function of a weight-2 newform on the congruence subgroup $\Gamma_0(N)$. The same holds for all quadratic twists $L(s, E_d)$. The Dirichlet coefficients of $L(s, E_d) = \sum_{n=1}^{\infty} a_d(n)n^{-s}$ are given by $a_d(n) = \chi_d(n)a(n)$, where χ_d is the primitive quadratic character modulo d (the Jacobi symbol modulo d). Now let

$$(1.5) \quad F(s) = L(s, E_d), \quad G(s) = L(s, \chi_d), \quad H(s) = L(s, E), \quad W(s) = \left(\frac{H'}{H} \right)'(s).$$

We can then apply Theorem 2 and obtain that the distribution of differences $\gamma - \gamma'$, with $F(\rho) = G(\rho') = 0$, is related to the behavior of $K_{F,G}(1 + it)$. The latter function is closely associated with the zeros of H . Of particular interest is the multiplicity $m_E(1/2)$ of the zero at the point $s = 1/2$, which carries important arithmetic information. By Mordell's theorem the group $E(\mathbb{Q})$ of rational points on E is finitely generated,

$$E(\mathbb{Q}) \cong \mathbb{Z}^r + E(\mathbb{Q})^{\text{tors}},$$

where $E(\mathbb{Q})^{\text{tors}}$ is a finite abelian group. According to the Birch and Swinnerton-Dyer Conjecture [2] (see also [40]), the rank r of E equals the multiplicity $m_E(1/2)$.

Theorem 5. *Assume that d is squarefree, and that all zeros of $H(s)$ with imaginary part $\leq \max(|\alpha|, |\beta|) + 1$ have real part equal to $\frac{1}{2}$. Then*

$$\lim_{k \rightarrow \infty} f_{F,G,g,\beta,k}(\alpha) = m_E \left(\frac{1}{2} - i\alpha \right) - m_E \left(\frac{1}{2} - i\beta \right),$$

where $m_E(\rho)$ denotes the multiplicity of the zero of $H(s)$ at $s = \rho$.

Taking $\alpha = 0$ and β an ordinate where $H(1/2 + i\beta) \neq 0$, the right side above equals the ‘‘analytic rank’’ $m_E(\frac{1}{2})$, which conjecturally equals the rank of $E(\mathbb{Q})$. The left side, on the other hand, is an expression involving only the distribution of zeros of $L(s, E_d)$ and zeros of $L(s, \chi_d)$. Conceptually what this means is that some arithmetic information on the given elliptic curve E is encoded in the sequences of zeros of *other* L -functions, not only the one associated to E .

2 Classical sums over zeros

Assume that $F \in \mathcal{S}^*$. We use frequently the estimate (cf. [33], (1.6))

$$(2.1) \quad N_F(T) = |\{\rho = \beta + i\gamma \in Z(F) : 0 < \beta < 1, 0 < \gamma \leq T\}| = \frac{d_F}{2\pi} T \log T + c_1 T + O(\log T)$$

for some constant $c_1 = c_1(F)$. An easy consequence is

$$(2.2) \quad N_F(T + A) - N_F(T) = O(A \log T) \quad (1 \leq A \leq T).$$

Another easy corollary is, for $F \in \mathcal{S}^*$ and $G \in \mathcal{S}^*$,

$$(2.3) \quad \sum_{\substack{0 < \gamma, \gamma' \leq T \\ \rho \in Z(F), \rho' \in Z(G)}} \min \left(K, \frac{1}{|\gamma - \gamma'|} \right) \ll T(K + \log T) \log^2 T \quad (K \geq 1).$$

Finally, we need a uniform version of Landau's theorem [10, Lemma 1]. The following is given in [11].

Lemma 2.1. *Let $F \in \mathcal{S}$, $x > 1$, $T \geq 2$, and let n_x be a nearest integer to x . Then, for any $\varepsilon > 0$,*

$$\sum_{0 < \gamma \leq T} x^\rho = -\frac{\Lambda_F(n_x)}{2\pi} \frac{e^{iT \log(x/n_x)} - 1}{i \log(x/n_x)} + O_\varepsilon \left(x^{1+\theta_F} \log(2x) + x^{1+\varepsilon} \log T + \frac{\log T}{\log x} \right).$$

Remark. If $1 < x < 3/2$, then $n_x = 1$, $\Lambda_F(n_x) = 0$ and the error term in Lemma 2.1 is $O(1 + \frac{\log T}{\log x})$. For $x = 1 + o(1/T)$, this is worse than the trivial bound $O(T \log T)$ coming from (2.1). Thus, we may replace the term $\frac{\log T}{\log x}$ in the statement of Lemma 2.1 by $\min(T \log T, \frac{\log T}{\log x})$.

3 The contribution of zeros off the critical line

For $F \in \mathcal{S}^*$, define

$$(3.1) \quad \begin{aligned} D_F(\xi) &= D_F(\xi; T) = \sum_{\substack{\rho \in Z(F) \\ 0 < \gamma \leq T}} e^{2\pi i \gamma \xi} \left(1 - e^{2\pi \xi (\beta - 1/2)} \right) \\ &= \frac{1}{2} \sum_{0 < \gamma \leq T} e^{2\pi i \gamma \xi} \left(2 - e^{2\pi \xi (\beta - 1/2)} - e^{2\pi \xi (1/2 - \beta)} \right). \end{aligned}$$

The second equality follows from the fact that if $\beta + i\gamma$ is a nontrivial zero of F then so is $1 - \beta + i\gamma$ (this follows from the functional equation (ii)). Throughout the remainder of this section, $\rho_j = \beta_j + i\gamma_j \in Z(F)$ ($j = 1, 2$).

Lemma 3.1. *Suppose that $F \in \mathcal{S}^*$. Uniformly for $T \geq 3$ and $1 \leq U \leq \frac{A(F)}{8\pi} \log T$,*

$$\int_0^U |D_F(\xi; T)|^2 d\xi \ll \frac{U^4 T}{\log T} (\log \log T)^4 + T^{1-A(F)/U} (\log T)^{B(F)+2}.$$

Proof. We have

$$(3.2) \quad \int_0^U |D_F(\xi; T)|^2 d\xi = \frac{1}{4} \sum_{0 < \gamma_1, \gamma_2 \leq T} \int_0^U e^{2\pi i \xi (\gamma_1 - \gamma_2)} \prod_{k=1}^2 (2 - e^{2\pi \xi (\beta_k - 1/2)} - e^{2\pi \xi (1/2 - \beta_k)}) d\xi.$$

Let $I(\rho_1, \rho_2, \xi)$ be the integrand on the right side of (3.2), and put

$$\Delta = \gamma_1 - \gamma_2, \quad \alpha = \max(|\beta_1 - 1/2|, |\beta_2 - 1/2|).$$

We consider four conditions on the pair (ρ_1, ρ_2) .

Case 1: $|\Delta| \leq 1$ and $\alpha \leq \frac{1}{U}$. Since $|2 - e^x - e^{-x}| \ll x^2$ for $0 \leq x \leq 2\pi$, we have

$$I(\rho_1, \rho_2, \xi) \ll \alpha^4 \xi^4 \leq \alpha^4 U^4.$$

Let

$$N_F^*(\sigma, T) = \#\{(\rho_1, \rho_2) : 0 < \gamma_1, \gamma_2 \leq T : \max(\beta_1, \beta_2) \geq \sigma, |\gamma_1 - \gamma_2| \leq 1\}.$$

From (2.2), we have

$$N_F^*(\sigma, T) \ll (\log T) N_F(\sigma, T).$$

The sum of $\int_0^U I(\rho_1, \rho_2, \xi) d\xi$ over all such pairs (ρ_1, ρ_2) is, using (2.1) and the zero-density estimate (vi),

$$\begin{aligned} &\ll U^5 \int_0^{1/U} \alpha^4 (-dN_F^*(\tfrac{1}{2} + \alpha, T)) \ll 3U^5 \int_0^{1/U} \alpha^3 N_F^*(\tfrac{1}{2} + \alpha, T) d\alpha \\ &\ll U^5 T \int_0^\infty \alpha^3 \min\left(T^{-\alpha A(F)} (\log T)^{B(F)+1}, \log^2 T\right) d\alpha \ll \frac{U^5 T}{\log^2 T} (\log \log T)^4. \end{aligned}$$

Case 2: $|\Delta| \leq 1$ and $\alpha > \frac{1}{U}$. Then $I(\rho_1, \rho_2, \xi) \ll e^{4\pi\alpha\xi}$ and thus

$$\int_0^U I(\rho_1, \rho_2, \xi) d\xi \ll \alpha^{-1} e^{4\pi\alpha U} \leq U e^{4\pi\alpha U}.$$

The contribution to the right side of (3.2) from all such pairs (ρ_1, ρ_2) is therefore

$$\begin{aligned} &\ll U \int_{1/U}^{1/2} e^{4\pi\alpha U} (-dN_F^*(\tfrac{1}{2} + \alpha, T)) \\ &= U \left[e^{4\pi N_F^*(\tfrac{1}{2} + \frac{1}{U}, T)} + 4\pi U \int_{1/U}^{1/2} e^{4\pi\alpha U} N_F^*(\tfrac{1}{2} + \alpha, T) d\alpha \right] \\ &\ll UT (\log T)^{B(F)+1} \left[T^{-A(F)/U} + U \int_{1/U}^\infty e^{(4\pi U - (\log T)A(F))\alpha} d\alpha \right] \\ &\ll T^{1-A(F)/U} (\log T)^{B(F)+2}. \end{aligned}$$

We used repeatedly the upper bound $U \leq \frac{A(F)}{8\pi} \log T$.

Case 3: $|\Delta| > 1$ and $\alpha \leq \frac{1}{U}$. We have

$$(3.3) \quad \int_0^U I(\rho_1, \rho_2, \xi) d\xi = \frac{1}{2\pi} \left[2g(0) - 2g(\beta_1 - \tfrac{1}{2}) - 2g(\beta_2 - \tfrac{1}{2}) + g(\beta_1 + \beta_2 - 1) + g(\beta_1 - \beta_2) \right],$$

where

$$g(z) = \frac{e^{2\pi i U \Delta + 2\pi U z} - 1}{i\Delta + z} + \frac{e^{2\pi i U \Delta - 2\pi U z} - 1}{i\Delta - z}.$$

Since we apply this with $z \leq \frac{1}{U}$, by Taylor's theorem we get

$$\begin{aligned} g(z) &= e^{2\pi i U \Delta} \frac{2i\Delta + (4\pi^2 i U^2 \Delta - 4\pi U)z^2 + O(|\Delta|z^4 U^4)}{-\Delta^2 - z^2} + \frac{2i\Delta}{\Delta^2 + z^2} \\ &= e^{2\pi i U \Delta} \frac{2i\Delta + (4\pi^2 i U^2 \Delta - 4\pi U - 2i/\Delta)z^2 + O(|\Delta|z^4 U^4)}{-\Delta^2} + \frac{2i}{\Delta} \left(1 - \frac{z^2}{i\Delta^2} + O\left(\frac{z^4}{\Delta^4}\right) \right). \end{aligned}$$

For any even quadratic polynomial $f : \mathbb{C} \rightarrow \mathbb{C}$, $2f(0) - 2f(a) - 2f(b) + f(a+b) + f(a-b) = 0$. Hence, by (3.3),

$$\int_0^U I(\rho_1, \rho_2, \xi) d\xi \ll \frac{\alpha^4 U^4}{|\Delta|}.$$

For non-negative integer j , define

$$N_j(\sigma, T) = \#\{(\rho_1, \rho_2) : 2^j < |\gamma_1 - \gamma_2| \leq 2^{j+1}, 0 < \gamma_1, \gamma_2 \leq T, \beta_1 \geq \sigma\}$$

and observe that by (2.1),

$$(3.4) \quad N_j(\sigma, T) \ll (2^j \log T) N_F(\sigma, T).$$

Using condition (vii) in the definition of \mathcal{S}^* , the contribution to the right side of (3.2) from all pairs (ρ_1, ρ_2) counted in case 3 is

$$\begin{aligned} &\ll \sum_{1 \leq 2^j \leq T} \frac{U^4}{2^j} \int_0^{1/U} \alpha^4 (-dN_j(\tfrac{1}{2} + \alpha, T)) \\ &\ll \sum_{1 \leq 2^j \leq T} \frac{U^4}{2^j} \int_0^{1/U} \alpha^3 N_j(\tfrac{1}{2} + \alpha, T) d\alpha \\ &\ll \sum_{1 \leq 2^j \leq T} U^4 T \int_0^\infty \alpha^3 \min\left(T^{-\alpha A(F)} (\log T)^{B(F)+1}, \log^2 T\right) d\alpha \ll \frac{U^4 T}{\log T} (\log \log T)^4. \end{aligned}$$

Case 4: $|\Delta| > 1$, $\alpha > \frac{1}{U}$. Using formula (3.3), the crude bound $|g(z)| \ll \frac{1}{|\Delta|} (e^{2\pi U|z|} + 1)$, and estimate (3.4), the contribution to the right side of (3.2) from all such pairs (ρ_1, ρ_2) is

$$\begin{aligned} &\ll \sum_{1 \leq 2^j \leq T} 2^{-j} \int_{1/U}^{1/2} e^{4\pi U \alpha} (-dN_j(\tfrac{1}{2} + \alpha, T)) \\ &= \sum_{1 \leq 2^j \leq T} 2^{-j} \left[e^{4\pi} N_j(\tfrac{1}{2} + \tfrac{1}{U}, T) + 4\pi U \int_{1/U}^{1/2} e^{4\pi U \alpha} N_j(\tfrac{1}{2} + \alpha, T) d\alpha \right] \\ &\ll T (\log T)^{B(F)+1} \sum_{1 \leq 2^j \leq T} \left[T^{-A(F)/U} + \frac{U}{\log T} T^{-A(F)/U} \right] \ll T^{1-A(F)/U} (\log T)^{B(F)+2}. \end{aligned}$$

Combining the estimates in the four cases completes the proof of the lemma. \square

4 Proof of Theorems 1 and 2

Throughout this section, $\rho_1 = \beta_1 + i\gamma_1 \in Z(F)$ and $\rho_2 = \beta_2 + i\gamma_2 \in Z(G)$.

Writing

$$Q_F(\xi; T) = \sum_{\substack{\rho \in Z(F) \\ 0 < \gamma \leq T}} e^{2\pi i \xi \gamma},$$

we then have

$$(4.1) \quad S := \sum_{0 < \gamma_1, \gamma_2 \leq T} h(\gamma_1 - \gamma_2) = \int_{\mathbb{R}} \hat{h}(\xi) Q_F(\xi; T) \overline{Q_G(\xi; T)} d\xi = 2\Re \int_0^\infty \hat{h}(\xi) Q_F(\xi; T) \overline{Q_G(\xi; T)} d\xi.$$

Decompose the integral as

$$(4.2) \quad S = 2\Re \int_0^U \hat{h}(\xi) Q_F(\xi; T) \overline{Q_G(\xi; T)} d\xi + \int_{|\xi| > U} \hat{h}(\xi) Q_F(\xi; T) \overline{Q_G(\xi; T)} d\xi =: 2\Re S_1 + S_2,$$

where

$$U = \frac{c \log T}{\log \log T}, \quad c = \frac{\min(1, A(F), A(G))}{100\pi}.$$

We can quickly dispense with S_2 due to the rapid decay of $\hat{h}'(\xi)$ from condition (3) in the definition of \mathcal{H} . Since

$$\int_W^V Q_F(\xi; T) \overline{Q_G(\xi; T)} d\xi \ll \sum_{0 < \gamma_1, \gamma_2 \leq T} \min\left(V - W, \frac{1}{|\gamma_1 - \gamma_2|}\right),$$

we obtain from (2.3) and integration by parts the bound

$$(4.3) \quad \left| \int_U^\infty \hat{h}(\xi) Q_F(\xi; T) \overline{Q_G(\xi; T)} d\xi \right| \ll \int_U^\infty |\hat{h}'(\xi)| \sum_{0 < \gamma_1, \gamma_2 \leq T} \min\left(\xi, \frac{1}{|\gamma_1 - \gamma_2|}\right) d\xi \\ \ll \frac{T \log^3 T}{U^4} + \frac{T \log^2 T}{U^3} \ll \frac{T(\log \log T)^4}{\log T}.$$

For S_1 , we decompose $Q_F(\xi; T)$ as follows:

$$Q_F(\xi; T) = \sum_{\substack{\rho \in Z(F) \\ 0 < \gamma \leq T}} \left[e^{2\pi\xi(\rho-1/2)} + e^{2\pi i\gamma\xi} \left(1 - e^{2\pi\xi(\beta-1/2)}\right) \right] = M_F(\xi) + E_F(\xi) + D_F(\xi),$$

where $D_F(\xi)$ is given by (3.1), and writing $x = e^{2\pi\xi}$,

$$M_F(\xi) = -\frac{\Lambda_F(n_x)}{2\pi\sqrt{n_x}} \frac{e^{iT \log(x/n_x)} - 1}{i \log(x/n_x)}.$$

By Lemma 2.1 (cf. the remark following the lemma) and condition (iv) defining the Selberg class ($|\Lambda_F(n)| \ll n^{\theta_F}$),

$$E_F(\xi) = -\frac{\Lambda_F(n_x)}{2\pi} \frac{e^{iT \log(x/n_x)} - 1}{i \log(x/n_x)} \left(\frac{1}{\sqrt{x}} - \frac{1}{\sqrt{n_x}} \right) + O\left(x^{1/2+\theta_F+\varepsilon} \log T + \min\left(\frac{\log T}{\log x}, T \log T\right)\right) \\ \ll |\Lambda_F(n_x)| \frac{|x - n_x|}{n_x^{3/2} |\log(x/n_x)|} + e^{2\pi\xi} \log T + \min\left(\frac{\log T}{\xi}, T \log T\right) \\ \ll e^{2\pi\xi} \log T + \min\left(\frac{\log T}{\xi}, T \log T\right).$$

Think of $M_F(\xi)$ as the main term and $D_F(\xi)$ and $E_F(\xi)$ as error terms. Making a similar decomposition $Q_G(\xi; T) = M_G(\xi) + D_G(\xi) + E_G(\xi)$, we have

$$(4.4) \quad \left| Q_F(\xi; T) \overline{Q_G(\xi; T)} - M_F(\xi) \overline{M_G(\xi)} \right| = \left| M_F(\xi) (\overline{E_G(\xi)} + \overline{D_G(\xi)}) + \overline{M_G(\xi)} (E_F(\xi) + D_F(\xi)) \right. \\ \left. + (E_F(\xi) + D_F(\xi)) (\overline{E_G(\xi)} + \overline{D_G(\xi)}) \right|.$$

Applying the Cauchy-Schwartz inequality, we obtain

$$(4.5) \quad \int_0^U |\hat{h}(\xi)| \cdot \left| Q_F(\xi; T) \overline{Q_G(\xi; T)} - M_F(\xi) \overline{M_G(\xi)} \right| d\xi \ll \mathcal{E} + \mathcal{D} + \mathcal{M}^{1/2} (\mathcal{E}^{1/2} + \mathcal{D}^{1/2}),$$

where

$$\mathcal{D} = \int_0^U |\hat{h}(\xi)| (|D_F(\xi)|^2 + |D_G(\xi)|^2) d\xi, \\ \mathcal{E} = \int_0^U |\hat{h}(\xi)| (|E_F(\xi)|^2 + |E_G(\xi)|^2) d\xi, \\ \mathcal{M} = \int_0^U |\hat{h}(\xi)| (|M_F(\xi)|^2 + |M_G(\xi)|^2) d\xi.$$

Since $\hat{h}(0) = 0$, we have $|\hat{h}(\xi)| \ll \xi$ and hence

$$(4.6) \quad \mathcal{E} \ll \int_0^{1/T} \xi (T \log T)^2 d\xi + \int_{1/T}^U \xi \left[\left(\frac{\log T}{\xi} \right)^2 + e^{4\pi\xi} \log^2 T \right] d\xi \\ \ll \log^2 T + \log^3 T + U e^{4\pi U} \log^2 T \ll T^{1/2}.$$

Observe that the hypothesized decay $|\hat{h}'(\xi)| \ll |\xi|^{-5}$ for $|\xi| \geq 1$ implies that

$$(4.7) \quad |\hat{h}(\xi)| \ll |\xi|^{-4} \quad (|\xi| \geq 1).$$

By (4.7), Lemma 3.1 and integration by parts,

$$(4.8) \quad \begin{aligned} \mathcal{D} &\ll \int_0^U (1+\xi)^{-5} \int_0^\xi |D_F(v)|^2 + |D_G(v)|^2 dv d\xi \\ &\ll \frac{T(\log \log T)^4}{\log T} \int_0^U (1+\xi)^{-1} d\xi \\ &\ll \frac{T(\log \log T)^5}{\log T}. \end{aligned}$$

Next, using condition (vi) in the definition of \mathcal{S}^* ,

$$(4.9) \quad \begin{aligned} \mathcal{M} &\ll 1 + \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{|\Lambda_F(n)|^2 + |\Lambda_G(n)|^2}{n} \int_{\frac{\log(n-1/2)}{2\pi}}^{\frac{\log(n+1/2)}{2\pi}} |\hat{h}(\xi)| \min\left(T, \frac{1}{|2\pi\xi - \log n|}\right)^2 d\xi \\ &\ll 1 + \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{T(|\Lambda_F(n)|^2 + |\Lambda_G(n)|^2)}{n(\log n)^4} \ll T. \end{aligned}$$

Thus, by (4.1), (4.2), (4.3), (4.5), (4.6), (4.8) and (4.9),

$$(4.10) \quad \begin{aligned} S &= 2\Re \int_0^U \hat{h}(\xi) M_F(\xi) \overline{M_G(\xi)} d\xi + O\left(\frac{T}{(\log T)^{1/3}}\right) \\ &= \frac{1}{2\pi^2} \Re \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \int_{\log(1-\frac{1}{2n})}^{\log(1+\frac{1}{2n})} \hat{h}\left(\frac{\log n + u}{2\pi}\right) \left|\frac{e^{iTu} - 1}{iu}\right|^2 du + O\left(\frac{T}{(\log T)^{1/3}}\right). \end{aligned}$$

Because \hat{h}' is bounded (a consequence of condition (3) in the definition of \mathcal{H}), replacing $\hat{h}(\frac{\log n + u}{2\pi})$ by $\hat{h}(\frac{\log n}{2\pi})$ produces a total error \mathcal{R} which satisfies

$$\begin{aligned} \mathcal{R} &\ll \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{|\Lambda_F(n)|^2 + |\Lambda_G(n)|^2}{n} \int_{\log(1-\frac{1}{2n})}^{\log(1+\frac{1}{2n})} |u| \min\left(T, \frac{1}{|u|}\right)^2 du \\ &\ll (\log T) \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{|\Lambda_F(n)|^2 + |\Lambda_G(n)|^2}{n} \ll \log^3 T. \end{aligned}$$

This implies that

$$S = \frac{1}{2\pi^2} \Re \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \hat{h}\left(\frac{\log n}{2\pi}\right) \int_{\log(1-\frac{1}{2n})}^{\log(1+\frac{1}{2n})} \left|\frac{e^{iTu} - 1}{iu}\right|^2 du + O\left(\frac{T}{(\log T)^{1/3}}\right).$$

Replacing the limits of integration with $(-\infty, \infty)$ produces an error of

$$\sum_{n \leq e^{2\pi U+1}} |\Lambda_F(n)|^2 + |\Lambda_G(n)|^2 \ll T^{1/2},$$

while the ‘‘complete’’ integral equals $2\pi T$. We obtain

$$S = \frac{T}{\pi} \Re \sum_{2 \leq n \leq e^{2\pi U+1}} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \hat{h}\left(\frac{\log n}{2\pi}\right) + O\left(\frac{T}{(\log T)^{1/3}}\right).$$

Using (vi) once again, we extend the sum on n to all positive integers n . The error \mathcal{R}' induced satisfies

$$\mathcal{R}' \ll T \sum_{n > e^{2\pi U}} \frac{|\Lambda_F(n)|^2 + |\Lambda_G(n)|^2}{n \log^4 n} \ll \frac{T}{U^2},$$

hence

$$S = \frac{T}{\pi} \Re \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \hat{h} \left(\frac{\log n}{2\pi} \right) + O \left(\frac{T}{(\log T)^{1/3}} \right).$$

Finally, writing \hat{h} as an integral and interchanging the sum and integral, we will show that

$$(4.11) \quad \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \hat{h} \left(\frac{\log n}{2\pi} \right) = \int_{\mathbb{R}} h(t) K_{F,G}(1+it) dt.$$

To show (4.11), we first observe that (vi) and the decay of $\hat{h}(\xi)$ imply that the left side of (4.11) is absolutely convergent, and thus

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \hat{h} \left(\frac{\log n}{2\pi} \right) &= \lim_{\delta \rightarrow 0^+} \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n^{1+\delta}} \hat{h} \left(\frac{\log n}{2\pi} \right) \\ &= \lim_{\delta \rightarrow 0^+} \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n^{1+\delta}} \int_{-\infty}^{\infty} h(t) n^{-it} dt. \end{aligned}$$

For each fixed $\delta > 0$, the sum-integral on the right side above is also absolutely convergent by (vi) and the assumption that $h \in L^1(\mathbb{R})$. Hence, by Fubini's theorem, we may interchange the sum and integral, obtaining

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n} \hat{h} \left(\frac{\log n}{2\pi} \right) &= \lim_{\delta \rightarrow 0^+} \int_{-\infty}^{\infty} h(t) \sum_{n=1}^{\infty} \frac{\Lambda_F(n) \overline{\Lambda_G(n)}}{n^{1+\delta+it}} dt \\ &= \lim_{\delta \rightarrow 0^+} \int_{-\infty}^{\infty} h(t) K_{F,G}(1+\delta+it) dt. \end{aligned}$$

Here we used the fact that the Dirichlet series for $K_{F,G}(s)$ converges to $K_{F,G}(s)$ for $\Re s > 1$. Finally, we must take the limit back inside the integral. We observe that by assumption, $K_{F,G}(s)$ is analytic in $\{s : \Re s \geq 1, s \neq 1\}$. In the case of $F(s) = G(s) = \zeta(s)$ in Theorem 1, the explicit formula (6.3) plus standard density bounds for the zeros immediately give

$$K_{F,G}(1+\delta+it) \ll \frac{1}{|\delta+it|^2} + \log^3(2+|t|)$$

uniformly for $\delta \geq 0$. Since $h \in \mathcal{H}_0$,

$$\int_{-\infty}^{\infty} h(t) |K_{F,G}(1+\delta+it)| dt$$

converges uniformly for $\delta > 0$, and this proves (4.11). For general F and G , we do not necessarily have a good enough zero-free region to deduce strong uniform bounds on $K_{F,G}(1+\delta+it)$ as $\delta \rightarrow 0^+$. But here we assume that h has compact support, so that the analyticity of $K_{F,G}$ still allows us to conclude (4.11). This completes the proof of Theorem 2.

5 Analyticity of $K_{F,G}$. Proof of Theorem 3

Let F and G be automorphic L -functions on $GL(n)$ and $GL(m)$, respectively, where n and m are positive integers. For basic analytic properties of automorphic L -functions, the reader may consult Sections 5.11 and

5.12 of [15]. In particular, the Euler products for F and G have the shape

$$F(s) = \prod_p \prod_{i=1}^n (1 - \alpha_i(p)p^{-s})^{-1}, \quad G(s) = \prod_p \prod_{j=1}^m (1 - \beta_j(p)p^{-s})^{-1}$$

for complex constants $\alpha_i(p), \beta_j(p)$. Both Euler products are absolutely convergent for $\Re s > 1$, and both functions have analytic continuation to $\mathbb{C} \setminus \{1\}$ and satisfy functional equations of type (1.2) (see e.g. [15, §5.12] and the references therein). Although the Ramanujan-Petersson conjecture (which states that $|\alpha_i(p)| \leq 1, |\beta_j(p)| \leq 1$ for all p, i and j) is not known for L -functions of degree exceeding 2, we do have the Luo-Rudnick-Sarnak bounds [15, (5.95)]

$$(5.1) \quad |\alpha_i(p)| \leq p^{\frac{1}{2} - \frac{1}{n^2+1}}, \quad |\beta_j(p)| \leq p^{\frac{1}{2} - \frac{1}{m^2+1}}.$$

The Rankin-Selberg convolution L -function of F and \overline{G} , which we denote by $H(s)$, has Euler product of the form [15, (5.9),(5.10)]

$$(5.2) \quad H(s) = \prod_{p \nmid Q} \prod_{i=1}^n \prod_{j=1}^m (1 - \alpha_i(p)\overline{\beta_j(p)}p^{-s})^{-1} \prod_{p \mid Q} \prod_{j=1}^{mn} (1 - \gamma_j(p)p^{-s})^{-1},$$

for some positive integer Q and numbers γ_j satisfying $|\gamma_j| < p$. The function $H(s)$ also has absolutely convergent Euler product for $\Re s > 1$ (and hence is nonzero in this open half-plane) [16]. Also, Shahidi proved that $H(1+it) \neq 0$ for all real t ([35]; with an announcement in [34]). Moreover, $H(s)$ is analytic at $s = 1$ unless $F = G$, in which case H has a simple pole at $s = 1$ [24].

By standard arguments, this is enough to establish a Mertens-type bound for sums of $\Lambda_H(n)$. From (5.2) we have that

$$\Lambda_H(p^k) = \frac{\Lambda_F(p^k)\overline{\Lambda_G(p^k)}}{\log p} \quad (p \nmid Q).$$

Hence, taking $F = G$ and using (5.1), we get the crude upper bound

$$\begin{aligned} \sum_{b \leq x} \frac{|\Lambda_F(b)|^2}{b} &\leq \sum_{(b,Q) > 1} \frac{|\Lambda_F(b)|^2}{b} + (e \log x) \sum_{(b,Q)=1} \frac{|\Lambda_F(b)|^2}{b^\sigma \log b} \quad \left(\text{take } \sigma = 1 + \frac{1}{\log x} \right) \\ &\ll 1 + (\log x) \left(-\frac{H'}{H}(\sigma) + \sum_{(b,Q) > 1} |\Lambda_H(b)| b^{-\sigma} \right). \end{aligned}$$

We have $-\frac{H'}{H}(\sigma) = \frac{1}{\sigma-1} + O(1) = \log x + O(1)$ and

$$\sum_{(b,Q) > 1} |\Lambda_H(b)| b^{-\sigma} \leq mn \sum_{p \mid Q} \log p \sum_{k=1}^{\infty} \frac{1}{p^{(\sigma-1)k}} \ll \log x.$$

This proves part (a) of Theorem 3; that is, condition (vi) holds.

To prove part (b), we observe that from the absolute convergence of the Euler product in the case $F = G$ we obtain the bound

$$(5.3) \quad \sum_{i=1}^n \sum_p \frac{|\alpha_i(p)|^2}{p^\sigma} < \infty \quad (\sigma > 1).$$

For $\Re s > 1$,

$$\log H(s) = \sum_{p \nmid Q} \frac{a_F(p)\overline{a_G(p)}}{p^s} + J(s) - \sum_{p \mid Q} \sum_{j=1}^{mn} \log(1 - \gamma_j(p)p^{-s}),$$

where

$$J(s) = \sum_{k=2}^{\infty} \frac{1}{k} \sum_{i=1}^n \sum_{j=1}^m \frac{(\alpha_i(p)\overline{\beta_j(p)})^k}{p^{ks}}.$$

Let $\delta = \min(\frac{1}{n^2+1}, \frac{1}{m^2+1})$. We claim that $J(s)$ is an absolutely convergent Dirichlet series (hence analytic) for $\Re s \geq 1 - \delta/2$. Indeed, by (5.1) and (5.3), for such s ,

$$\sum_{k=2}^{\infty} \frac{1}{k} \sum_{i,j} \left| \frac{(\alpha_i(p)\beta_j(p))^k}{p^{ks}} \right| \leq \sum_{k=2}^{\infty} \frac{1}{k} \sum_{i,j} \frac{p^{(1-2\delta)(k-1)}}{p^{(1-\delta/2)k}} |\alpha_i(p)|^2 < \infty.$$

Similarly,

$$K_{F,G}(s) = \sum_{p \nmid Q} \frac{a_F(p) \overline{a_G(p)} \log^2 p}{p^s} + \tilde{K}(s),$$

where $\tilde{K}(s)$ is analytic for $\Re s \geq 1$. Therefore, initially for $\Re s > 1$ and for $\Re s \geq 1$ (but $s \neq 1$ if $F = G$) by analytic continuation,

$$K_{F,G}(s) = \left(-\frac{H'}{H} \right)'(s) + E(s),$$

where $E(s)$ is analytic for $\Re s \geq 1$. This proves part (b) of Theorem 3.

6 Proof of Theorem 4

Fix an integer $q > 1$ and a primitive Dirichlet character χ mod q . We also fix a compactly supported function g in $C^\infty(\mathbb{R})$ such that $g \geq 0$ and $\int g = 1$. Denoting $g_m(x) = mg(mx)$, g_m is more concentrated at the origin (as $m \rightarrow \infty$, g_m approaches a Dirac delta function), and also $\int g_m = 1$. Fix real numbers α, β in $(\frac{1}{2}, \infty)$, and let

$$h_m(x) = g_m(x - \alpha) - g_m(x - \beta).$$

Here the key idea is that integrating by parts k times gives

$$\int h_m^{(k)}(t) H(t) dt = \int h_m(t) H^{(k)}(t) dt.$$

By Theorem 2 with $F = L(s, \chi)$, as $T \rightarrow \infty$,

$$S_{m,k}(T) := \frac{2\pi}{T} \sum_{\substack{0 < \gamma_1, \gamma_2 \leq T \\ \rho_1, \rho_2 \in Z(L(s, \chi))}} h_m^{(k)}(\gamma_1 - \gamma_2) \rightarrow \int h_m^{(k)}(t) H_\chi(t) dt = \int h_m(t) H_\chi^{(k)}(t) dt,$$

where

$$H_\chi(t) = K_\chi(1 + it) + K_\chi(1 - it),$$

$$K_\chi(s) = \sum_{n=1}^{\infty} \frac{|\Lambda_F(n)|^2}{n^s} \quad (\Re s > 1),$$

and $K_\chi(s)$ is given by analytic continuation for $\Re s = 1$. Since $\Lambda_F(n) = \Lambda(n)\chi(n)$,

$$K_\chi(s) = \sum_{(n,q)=1} \frac{\Lambda^2(n)}{n^s} = K(s) - \sum_{p|q} \frac{\log^2 p}{p^s - 1}.$$

Since $H_\chi(t) \in C^\infty(0, \infty)$ and h_m approaches the difference of two Dirac deltas as $m \rightarrow \infty$,

$$(6.1) \quad \lim_{m \rightarrow \infty} \lim_{T \rightarrow \infty} S_{m,k}(T) = \lim_{m \rightarrow \infty} \int h_m(t) H_\chi^{(k)}(t) dt = H_\chi^{(k)}(\alpha) - H_\chi^{(k)}(\beta),$$

for every fixed k .

Lemma 6.1. *Uniformly for $\Re s = 1$, $s \neq 1$,*

$$\left| K_\chi^{(k)}(s) - K^{(k)}(s) \right| \ll k! \log q.$$

Proof.

$$\begin{aligned}
K^{(k)}(s) - K_{\chi}^{(k)}(s) &= \sum_{p|q} \log^2 p [(p^s - 1)^{-1}]^{(k)} = \sum_{p|q} (-\log p)^{k+2} \sum_{j=1}^{\infty} \frac{j^k}{p^{js}} \\
&\ll \sum_{p|q} (\log p)^{k+2} \int_0^{\infty} x^k p^{-x} dx \\
&= \sum_{p|q} (\log p) \cdot k! \leq k! \log q.
\end{aligned}$$

□

Corollary 1. *Uniformly for $t \neq 0$,*

$$H_{\chi}^{(k)}(t) - H^{(k)}(t) \ll k! \log q.$$

Lemma 6.2. *Let*

$$W(s) = \left(\frac{\zeta'}{\zeta} \right)'(s),$$

and let S be a compact set of real numbers not containing zero. Then, uniformly for $s = 1 + it$, with $t \in S$, we have

$$\lim_{k \rightarrow \infty} \left| \frac{K^{(k)}(s) - W^{(k)}(s)}{2^k (k+1)!} \right| = 0.$$

Proof. First, we have

$$\begin{aligned}
(6.2) \quad K^{(k)}(s) - W^{(k)}(s) &= (-1)^k \sum_{n=1}^{\infty} \frac{\Lambda^2(n) \log^k n - \Lambda(n) \log^{k+1} n}{n^s} \\
&= (-1)^k \sum_{l=2}^{\infty} \sum_p \frac{(l^k - l^{k+1})(\log p)^{k+2}}{p^{ls}} \\
&=: (-1)^k \sum_{l=2}^{\infty} (l^k - l^{k+1}) R_{k,l}.
\end{aligned}$$

If $l \geq 3$, then

$$\begin{aligned}
|R_{k,l}| &\leq \int_{2^-}^{\infty} \frac{(\log u)^{k+1}}{u^l} d\theta(u) = \int_2^{\infty} \theta(u) \frac{(\log u)^{k+1}}{u^l} \left(\frac{l}{u} - \frac{k+1}{u \log u} \right) du \\
&\ll l \int_2^{\infty} \frac{(\log u)^{k+1}}{u^l} du = \frac{l}{(l-1)^{k+2}} \int_{(l-1) \log 2}^{\infty} w^{k+1} e^{-w} dw \\
&\ll \begin{cases} \frac{l}{(l-1)^{k+2}} \cdot (k+1)! & (\text{all } l), \\ \frac{l}{(l-1)^{k+2}} \cdot 2^{-(l-1)/2} \int_0^{\infty} w^{k+1} e^{-w/2} dw \ll \frac{l}{(l-1)^{k+2}} \cdot 2^{k-l/2} (k+1)! & (l > 2k). \end{cases}
\end{aligned}$$

Therefore

$$\begin{aligned}
\left| \sum_{l \geq 3} (l^k - l^{k+1}) R_{k,l} \right| &\ll (k+1)! \left[\sum_{l=3}^{2k} \left(\frac{l}{l-1} \right)^{k+2} + 2^k \sum_{l=2k+1}^{\infty} \left(\frac{l}{l-1} \right)^{k+2} 2^{-l/2} \right] \\
&\ll (k+1)! \left\{ \left(\frac{3}{2} \right)^k + k \left(\frac{4}{3} \right)^k + 1 \right\} \ll \left(\frac{3}{2} \right)^k (k+1)!.
\end{aligned}$$

When $l = 2$, we must be more precise, using the Prime Number Theorem in the form $\theta(u) \sim u$ as $u \rightarrow \infty$. Write $\theta(u) = u + E(u)$. In what follows, $o(1)$ stands for a function of k that tends to zero as $k \rightarrow \infty$. By partial summation we get

$$R_{k,2} = \int_{2^-}^{\infty} \frac{(\log u)^{k+1}}{u^{2+2it}} du + \int_2^{\infty} \frac{(\log u)^{k+1}}{u^{3+2it}} E(u) \left(2 + 2it - \frac{k+1}{\log u}\right) du.$$

Making the change of variables $w = \log u$ and using that $E(u) = o(u)$ as $u \rightarrow \infty$, we obtain

$$\begin{aligned} |R_{k,2}| &\ll 1 + \left| \int_0^{\infty} \frac{w^{k+1}}{e^{(1+2it)w}} dw \right| + \int_{\log 2}^{k/10} w^{k+1} \left(1 + |t| + \frac{k+1}{w}\right) dw + o(1) \int_{k/10}^{\infty} \frac{w^{k+1}}{e^w} (1 + |t|) dw \\ &= 1 + \frac{(k+1)!}{|1+2it|^{k+2}} + O((1+|t|)(k/10)^{k+2}) + o(1)(1+|t|)(k+1)! \\ &= o((k+1)!) \end{aligned}$$

uniformly for $t \in S$. It follows from (6.2) that $K^{(k)}(1+it) - W^{(k)}(1+it) = o(2^k(k+1)!)$ uniformly for $t \in S$. \square

In the same way, we can prove an analogous estimate needed for Theorem 5.

Lemma 6.3. *Let F, G, H, W be given by (1.5), and let S be a compact set of real numbers not containing zero. Then, uniformly for $s = 1 + it$, with $t \in S$, we have*

$$\lim_{k \rightarrow \infty} \left| \frac{K_{F,G}^{(k)}(s) - W^{(k)}(s)}{2^k(k+1)!} \right| = 0.$$

Proof. We compute that $\Lambda_H(p^l) = (\alpha_p^l + \beta_p^l) \log p$ and

$$\Lambda_F(p^l) \Lambda_G(p^l) = \begin{cases} \Lambda_H(p^l) \log p & p \nmid Nd \\ 0 & p | Nd. \end{cases}$$

Hence,

$$(-1)^k \left(W^{(k)}(s) - K_{F,G}^{(k)}(s) \right) = \sum_{\substack{p, l \geq 2 \\ p \nmid Nd}} \frac{(l^{k+1} - l^k) \Lambda_H(p^l) (\log p)^{k+1}}{p^{ls}} + \sum_{p|Nd} \sum_{l=1}^{\infty} \frac{(l \cdot \log p)^{k+1} \Lambda_H(p^l)}{p^{ls}}.$$

For $\Re s = 1$, the second double sum above is, in absolute value, at most

$$\begin{aligned} 2 \sum_{p|Nd} (\log p)^{k+2} \sum_{l=1}^{\infty} \frac{l^{k+1}}{p^l} &\ll \sum_{p|Nd} (\log p)^{k+2} \int_0^{\infty} x^{k+1} p^{-x} dx \\ &= (k+1)! \sum_{p|Nd} 1 \ll (k+1)!. \end{aligned}$$

Since $|\Lambda_H(p^l)| \leq 2 \log p$, the argument used in the proof of Lemma 6.2 implies that the terms in the sum over $p \nmid Nd$ and $l \geq 3$ total $O((3/2)^k(k+1)!)$. The terms with $l = 2$ and $p \nmid Nd$ total

$$2^k \sum_{p \nmid Nd} \frac{(a(p)^2 - 2)(\log p)^{k+2}}{p^{2s}}.$$

From the standard theory of Rankin-Selberg convolutions [15, Theorems 5.13, 5.44], we have $\sum_{p \leq x} a(p)^2 \log p \sim x$ (the analog of the Prime Number Theorem for $a(p)^2$) and the argument in the proof of Lemma 6.2 implies that the above sum is $o(2^k(k+1)!)$, uniformly for the values of s under consideration. \square

Proof of Theorem 4. Denote by

$$H_k(t) = \frac{(-1)^{k+1}}{2^{k+1}(k+1)!} \left(W^{(k)}(1+it) + W^{(k)}(1-it) \right).$$

By (6.1), Corollary 1, and Lemma 6.2,

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{(-1)^{k+1}}{2^{k+1}(k+1)!} \lim_{m \rightarrow \infty} \lim_{T \rightarrow \infty} S_{m,k}(T) &= \lim_{k \rightarrow \infty} \frac{(-1)^{k+1}}{2^{k+1}(k+1)!} \left(H^{(k)}(\alpha) - H^{(k)}(\beta) \right) \\ &= \lim_{k \rightarrow \infty} (H_k(\alpha) - H_k(\beta)). \end{aligned}$$

By the explicit formula for $\zeta'(s)/\zeta(s)$, e.g. [7, §12, (8)],

$$W(s) = \frac{1}{(s-1)^2} - \sum_{\rho} \frac{1}{(s-\rho)^2} - \frac{1}{2} \left(\frac{\Gamma'}{\Gamma} \left(\frac{s}{2} + 1 \right) \right)'$$

Using a well-know series expansion for $\Gamma'(s)/\Gamma(s)$, we get for $k \geq 0$ that

$$(6.3) \quad W^{(k)}(s) = (-1)^k (k+1)! \left(\frac{1}{(s-1)^{k+2}} - \sum_{\rho}^* \frac{1}{(s-\rho)^{k+2}} \right),$$

where the sum on ρ is over *all* zeros of ζ , including the trivial zeros at the points $\rho = -2, -4, \dots$

Assume RH: For any $t > \frac{1}{2}$,

$$\begin{aligned} \lim_{k \rightarrow \infty} H_k(t) &= \lim_{k \rightarrow \infty} \frac{(-1)^{k+k+1}}{2^{k+1}} \left(\frac{1}{(it)^{k+2}} + \frac{1}{(-it)^{k+2}} - \sum_{\rho}^* \frac{1}{(1+it-\rho)^{k+2}} + \frac{1}{(1-it-\rho)^{k+2}} \right) \\ &= \frac{1}{2} \left(m_{\zeta} \left(\frac{1}{2} + it \right) + m_{\zeta} \left(\frac{1}{2} - it \right) \right) = m_{\zeta} \left(\frac{1}{2} + it \right), \end{aligned}$$

where $m_{\zeta}(\rho)$ denotes the multiplicity of the zero of the Riemann zeta function at ρ . We note that the above limit calculation remains valid under the weaker assumption that there are no zeros off the critical line with imaginary part in $[t - \frac{1}{2}, t + \frac{1}{2}]$; that is, the full strength of RH is not necessary.

This proves the implication (b) \implies (a) in Theorem 4, and it also shows that

$$\lim_{k \rightarrow \infty} f_{\chi,g,\beta,k}(\alpha) = m_{\zeta} \left(\frac{1}{2} + i\alpha \right) - m_{\zeta} \left(\frac{1}{2} + i\beta \right).$$

Assume RH fails: Then let ρ_0 be a zero of the Riemann zeta function with $\Re \rho_0 > \frac{1}{2}$. We distinguish two cases, according as to whether $\zeta(s)$ has any zeros inside the open disk of radius $\frac{1}{2}$ centered at $1 + i\beta$.

Assume first that there are no zeros of $\zeta(s)$ inside this disk. We then choose $\alpha = \Im \rho_0$, and consider the zeros of $\zeta(s)$ closest to $1 + i\alpha$. Denote the minimum distance to $1 + i\alpha$ by δ_0 , and let ρ_1, \dots, ρ_r denote the zeros of $\zeta(s)$ at distance δ_0 from $1 + i\alpha$, with multiplicities m_1, \dots, m_r . We write the differences $1 + i\alpha - \rho_j$ in the form

$$1 + i\alpha - \rho_j = \delta_0 e^{2\pi i \theta_j}, \quad j = 1, \dots, r.$$

Then the contribution of ρ_1, \dots, ρ_r in $f_{\chi,g,\beta,k}(\alpha)$ is of the order of magnitude of $\left(\frac{1}{2\delta_0} \right)^k \sum_{j=1}^r m_j \cos 2\pi k \theta_j$. We now let k tend to infinity along a subsequence for which all the fractional parts $\{k\theta_j\}$ tend to zero. Then along this subsequence, $\sum_{j=1}^r m_j \cos 2\pi k \theta_j \rightarrow \sum_{j=1}^r m_j$. By comparison, the contribution of all the other zeros of $\zeta(s)$ in $f_{\chi,g,\beta,k}(\alpha)$ is exponentially small. Since $\delta_0 < \frac{1}{2}$, it follows that $f_{\chi,g,\beta,k}(\alpha)$ tends to infinity along this subsequence.

Assume now that $\zeta(s)$ has at least one zero inside the open disk of radius $\frac{1}{2}$ centered at $1 + i\beta$. In this case we choose any $\alpha \in (\frac{1}{2}, \infty)$ for which all the zeros of $\zeta(s)$ are at distance larger than $\frac{1}{2}$ from $1 + i\alpha$. We then repeat the reasoning from the previous case, where the role of ρ_1, \dots, ρ_r is now played by the zeros of $\zeta(s)$ closest to $1 + i\beta$. As above, it follows that $f_{\chi,g,\beta,k}(\alpha)$ tends to infinity along a suitable subsequence.

This completes the proof of the implication (a) \implies (b) in Theorem 4. \square

Proof of Theorem 5. This follows in the same way, using Lemma 6.3 instead of Lemma 6.2. Condition (vii) from the definition of the class \mathcal{S}^* holds by the analysis in Section 7. Condition (vi) holds by Theorem 3, and the other conditions hold by the discussion preceding the statement of Theorem 5. Unlike the situation in Theorem 4, F , G and H are all entire functions and thus α and β are not excluded from the interval $[-\frac{1}{2}, \frac{1}{2}]$. This comes from the analog of the explicit formula (6.3), where there will be no term corresponding to the pole at $s = 1$ as with $\zeta(s)$. \square

7 Notes on shifted convolution sums of Fourier coefficients of $\Gamma_0(D)$ cusp forms

7.1 Zero density result

Luo's zero density result follows from understanding a mollified second moment. Since $L(1/2 + it, f)$ has conductor $|t|^2$, this requires understanding the shifted convolution problem for the Fourier coefficients $\lambda_f(n)$ of f (see Lemma 2.1 of [22]). In the case of the full modular group, results were proven by Hafner [12] and subsequently by Luo [22]. Unfortunately, there does not appear to be such a result for general congruence subgroups in the literature and we provide a proof below for the sake of completeness.

7.2 Shifted convolution sum

Fix $\mu, \nu > 0$ and $l \in \mathbb{Z}$, with $l \neq 0$. Let $\omega(x)$ be a smooth function supported on $[1/2, 5/2]$ and $N \geq 1$ (with $N \gg (\mu\nu)^\epsilon$ for convenience). Let

$$(7.1) \quad \mathcal{S} = \mathcal{S}(\mu, \nu, l) := \sum_{\mu m - \nu n = l} \lambda_f(n) \lambda_f(m) \omega\left(\frac{n}{N}\right).$$

We have normalized the coefficients λ_f so that $\lambda_f(n) \ll n^\epsilon$ is known by the work of Deligne. We study \mathcal{S} using the delta method, as used by Duke, Friedlander and Iwaniec in [8]. We state the version developed by Heath-Brown [13]. As usual, let $\delta(0) = 1$ and $\delta(n) = 0$ for $n \neq 0$. Then there exists a smooth function $h : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\delta(n) = \frac{1}{Q^2} \sum_{q \geq 1} \sum_{\substack{a \pmod{q} \\ (a, q) = 1}} e\left(\frac{an}{q}\right) h\left(\frac{q}{Q}, \frac{n}{Q^2}\right) + O_A\left(\frac{1}{Q^A}\right).$$

Applying this to (7.1) with $Q = \sqrt{\nu N}$, we see that

$$(7.2) \quad \mathcal{S} = \frac{1}{Q^2} \sum_{q \geq 1} \sum_{\substack{a \pmod{q} \\ (a, q) = 1}} e\left(\frac{-al}{q}\right) \mathcal{S}(a) + O_A\left(\frac{1}{N^A}\right),$$

where

$$(7.3) \quad \mathcal{S}(a) = \sum_{m, n} \lambda_f(n) \lambda_f(m) e\left(\frac{a(\mu m - \nu n)}{q}\right) g(m, n),$$

where $g(x, y)$ is a smooth function compactly supported on $[\frac{\nu N}{2\mu} + \frac{l}{\mu}, \frac{5\nu N}{2\mu} + \frac{l}{\mu}] \times [N/2, 5N/2]$, and moreover

$$\frac{\partial^{j+k}}{\partial x^j \partial y^k} g(x, y) \ll_{j, k} \max\left(N^{-j}, \left(\frac{\mu}{Q^2}\right)^j\right) \max\left(\left(\frac{\nu}{Q^2}\right)^k, N^{-k}\right) \ll \frac{1}{N^{l+j}},$$

for all $j, l \geq 0$.

Let $\mu' = \mu/(\mu, q)$, $q_\mu = q/(\mu, q)$, $\nu' = \nu/(\nu, q)$, and $q_\nu = q/(\nu, q)$. Further, let $D_{1, \mu} = (q_\mu, D)$ and let $D_{2, \mu} = D/D_{1, \mu}$ and similarly define $D_{1, \nu}$ and $D_{2, \nu}$. By Chinese Remainder Theorem, we have that $\chi = \chi_{D_{1, \mu}} \chi_{D_{2, \mu}}$ for unique $\chi_{D_{i, \mu}}$ characters modulo $D_{i, \mu}$.

Now we apply the Voronoi summation formula for these coefficients due to Kowalski, Michel and VanderKam (see Appendix A [21] for the proof and exact notation) to get that

$$(7.4) \quad \mathcal{S}(a) = \chi_{D_{1,\mu}}(\bar{a})\chi_{D_{1,\nu}}(\bar{a})\chi_{D_{2,\mu}}(-q_\mu)\chi_{D_{2,\nu}}(-q_\nu) \frac{\eta_f(D_{2,\mu})\eta_f(D_{2,\nu})}{\sqrt{D_{2,\mu}D_{2,\nu}}} \\ \sum_{m,n} \lambda_{fD_{2,\mu}}(n)\lambda_{fD_{2,\nu}}(m) e\left(\frac{\overline{a\mu'}m}{q_\mu}\right) e\left(\frac{-\overline{a\nu'}n}{q_\nu}\right) G(m,n),$$

where

$$G(m,n) = \frac{4\pi^2}{q_\nu q_\mu} \int_0^\infty \int_0^\infty g(x,y) J_{k-1}\left(\frac{4\pi\sqrt{mx}}{q_\mu\sqrt{D_{2,\mu}}}\right) J_{k-1}\left(\frac{4\pi\sqrt{ny}}{q_\nu\sqrt{D_{2,\nu}}}\right) dx dy.$$

The rest of the proof is similar to pg. 156 of [22]. To be more precise, using integration by parts and the fact that

$$J_{k-1}(2\pi x) = \frac{1}{\pi\sqrt{x}} \Re(W(2\pi x)e(x - k/4 + 1/8)),$$

for an essentially flat function W satisfying $x^l W^l(x) \ll 1$, we see that the sum over m and n is essentially restricted to be of length $\frac{q_\mu^2 D_{2,\mu} \mu}{\nu N^{1-\epsilon}}$ and $\frac{q_\nu^2 D_{2,\nu}}{N^{1-\epsilon}}$, respectively. Moreover, this tells us that trivially

$$G(m,n) \ll \frac{1}{\sqrt{q_\nu q_\mu} (mn)^{1/4}} \left(N \frac{\nu}{\mu} N\right)^{3/4}.$$

Also, the sum over a creates a Kloosterman sum $S_\chi(l, *; q)$ for which the Weil bound may be applied. This gives the bound

$$\mathcal{S} \ll_f \sqrt{l\nu}^{3/4} N^{3/4+\epsilon}.$$

Note that the above bound is symmetric in N and $\frac{\nu}{\mu} N$ (in the sense that it is equal to $(\frac{\nu}{\mu} N)^{3/4} \mu^{3/4}$).

One may then derive analytic continuation and bounds for the Dirichlet series

$$D_{\mu,\nu}(s,l) = \sum_{n \geq 1} \frac{a(n)a((\nu n + l)/\mu)}{(\nu n + l/2)^s}$$

by using a smooth partition of unity. This suffices for the purposes of proving a zero density result as in Theorem 1.1 of [22].

Acknowledgements. The authors are grateful to the referees for their helpful comments and suggestions. The authors thank Xiannan Li for the material in Section 7, for helpful discussions about Rankin-Selberg L -functions, and for posing the question of general linear combinations of zeros $\gamma - \alpha\gamma'$. The authors are also thankful to Christopher Hughes, Jon Keating, Jeffrey Lagarias, Brad Rodgers, and Michael Rubinstein for informing them about previous work on this topic.

References

- [1] M. V. Berry and J. P. Keating, *The Riemann zeros and eigenvalue asymptotics*, SIAM Rev. **41** (1999), 236–266.
- [2] B. J. Birch and H. P. F. Swinnerton-Dyer, *Notes on elliptic curves. II*, J. Reine Angew. Math. **218** 1965 79–108.
- [3] E. B. Bogomolny and J. P. Keating, *Gutzwiller's trace formula and spectral statistics: beyond the diagonal approximation*, Phys. Rev. Lett. **77** (1996), 1472–1475.
- [4] C. Breuil, B. Conrad, F. Diamond, and R. Taylor, *On the modularity of elliptic curves over \mathbb{Q} : wild 3–adic exercises*, J. Amer. Math. Soc. **14** (2001), 843–939.
- [5] J. B. Conrey, D. W. Farmer, and M. R. Zirnbauer, *Autocorrelation of ratios of L -functions*, Commun. Number Theory Phys. **2** (2008), no. 3, 593–636.
- [6] J. B. Conrey and N. C. Snaith, *Applications of the L -functions ratios conjectures*, Proc. Lond. Math. Soc. **94** (2007), 594–646.
- [7] H. Davenport, *Multiplicative Number Theory*, 3rd ed., Springer-Verlag, 2000.
- [8] W. Duke, J.B. Friedlander and H. Iwaniec, *Bounds for automorphic L -functions*. Invent. Math. **112** (1993), 1–8.
- [9] F. J. Dyson, *Statistical theory of the energy levels of complex systems, I, II, III*, J. Math. Phys. **3** (1962), 14–156, 15–165, 16–175.

- [10] K. Ford and A. Zaharescu, *On the distribution of imaginary parts of zeros of the Riemann zeta function*, J. reine angew. Math. **579** (2005), 145–158.
- [11] K. Ford, K. Soundararajan and A. Zaharescu, *On the distribution of imaginary parts of zeros of the Riemann zeta function, II*, Math. Ann. **343** (2009), 487–505.
- [12] J.L. Hafner, Explicit estimates in the arithmetic theory of cusp forms and Poincaré series. Math. Ann. 264 (1983), no. 1, 9–20.
- [13] D.R. Heath-Brown, A new form of the circle method, and its application to quadratic forms. J. Reine Angew. Math. 481 (1996), 149–206.
- [14] D. Hejhal, *On the triple correlation of the zeros of the zeta function*, Int. Math. Res. Not. (1994), 293–302.
- [15] H. Iwaniec and E. Kowalski, Analytic Number Theory, Amer. Math. Soc., 2004.
- [16] H. Jacquet and J. A. Shalika, *On Euler products and the classification of automorphic representations. I.*, Amer. J. Math. **103** (1981), no. 3, 499–558; *On Euler products and the classification of automorphic forms. II.* Amer. J. Math. **103** (1981), no. 4, 777–815.
- [17] J. Kaczorowski, *Axiomatic Theory of L-Functions: the Selberg Class*, Analytic number theory, 133–209, Lecture Notes in Math., vol. 1891, Springer, Berlin, 2006.
- [18] J. Kaczorowski and A. Perelli, *The Selberg class: a survey*, Number Theory in Progress, vol. II, de Gruyter, Berlin (1999), 953–992.
- [19] N. Katz, P. Sarnak, *Zeros of zeta functions and symmetry*, Bull. Amer. Math. Soc. (N.S.) **36** (1999), no. 1, 1–26.
- [20] N. Katz, P. Sarnak, *Random matrices*, Frobenius eigenvalues, and monodromy, Amer. Math. Soc. Colloqu. Publ. **45** (1999).
- [21] E. Kowalski, P. Michel, and J. VanderKam, Rankin-Selberg L-functions in the level aspect. Duke Math. J. 114 (2002), no. 1, 123–191.
- [22] W. Luo, *Zeros of Hecke L-functions associated with cusp forms.*, Acta Arith. **71** (1995), no. 2, 139–158.
- [23] R. Pérez-Marco, *Statistics on Riemann zeros*, preprint, ArXiv:1112.0346
- [24] C. Mœglin and J.-L. Waldspurger, Le spectre résiduel de $GL(n)$ (**French**) [The residual spectrum of $GL(n)$], Ann. Sci. École Norm. Sup. (4) **22** (1989), 605–674.
- [25] H. L. Montgomery, The pair correlation of zeros of the zeta function, Analytic number theory, Proc. Sympos. Pure Math. XXIV, St. Louis Univ., St. Louis, Mo. (1972), 181–193.
- [26] A. M. Odlyzko, *On the distribution of spacings between zeros of the zeta function*, Math. Comp. **48** (1987), 273–308.
- [27] A. Perelli, *A Survey of the Selberg Class of L-Functions, Part I*, Milan J. Math. **73** (2005), 19–52.
- [28] A. Perelli, *A Survey of the Selberg Class of L-functions. Part II*, Riv. Mat. Univ. Parma (7) **3** (2004), 83–118.
- [29] B. Rodgers, *Macroscopic pair correlation of the Riemann zeroes for smooth test functions*, Quart. J. Oxford, to appear.
- [30] Z. Rudnick, P. Sarnak, *Zeros of principal L-functions and random matrix theory*, A celebration of John F. Nash, Jr., Duke Math. J. **81** (1996), no. 2, 269–322.
- [31] A. Selberg, *Contributions to the theory of the Riemann zeta-function*, Arch. Math. Naturvid. **48** (1946), 89–155; Collected papers, vol. I, 214–280, Springer, Berlin 1989.
- [32] A. Selberg, *Contributions to the theory of Dirichlet’s L-functions*, Skr. Norske Vid. Akad. Oslo. I. **1946**, (1946), no. 3, 62 pp.; Collected papers, vol. I, 281–340, Springer, Berlin 1989.
- [33] A. Selberg, *Old and new conjectures and results about a class of Dirichlet series*, Proceedings of the Amalfi Conference on Analytic Number Theory (Maiori, 1989), Univ. Salerno, (1992), 367–385; Collected papers, vol. II, 47–63, Springer, Berlin 1989.
- [34] F. Shahidi, *On nonvanishing of L-functions*, Bull. Amer. Math. Soc. (N.S.) **2** (1980), no. 3, 462–464.
- [35] F. Shahidi, *On certain L-functions*, Amer. J. Math. **103** (1981), no. 2, 297–355.
- [36] N. C. Snaith, *Riemann zeros and random matrix theory*, Milan Math. J. **78** (2010), 135–152.
- [37] R. Taylor, A. Wiles, *Ring-theoretic properties of certain Hecke algebras*, Ann. Math. **141** (1995), 553–572.
- [38] E. Wigner, *Random matrices in physics*, SIAM Rev. **9** (1967), 1–23.
- [39] A. Wiles, *Modular elliptic curves and Fermat’s last theorem*, Ann. Math. **142** (1995), 443–551.
- [40] A. Wiles, *The Birch and Swinnerton-Dyer conjecture*, The millennium prize problems, 31–41, Clay Math. Inst., Cambridge, MA, 2006.

DEPARTMENT OF MATHEMATICS, 1409 WEST GREEN STREET, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801, USA

E-mail address: ford@math.uiuc.edu

E-mail address: zaharesc@math.uiuc.edu