

On Avoider-Enforcer games

József Balogh* Ryan Martin†

August 16, 2007

Abstract

In the Avoider-Enforcer game on the complete graph K_n , the players (Avoider and Enforcer) each take an edge in turn. Enforcer wins the game when he can require Avoider's graph to have a given property \mathcal{P} . The important parameter is $\tau_E(\mathcal{P})$, the most number of rounds required for Enforcer to win if Avoider plays with an optimal strategy ($\tau_E(\mathcal{P}) = \infty$ if Avoider can finish the game without creating a graph with property \mathcal{P}).

In this paper, let \mathcal{F} be an arbitrary family of graphs and \mathcal{P} be the property that a member of \mathcal{F} is a subgraph or is an induced subgraph. We determine the asymptotic value of $\tau_E(\mathcal{P})$ when \mathcal{F} contains no bipartite graph and establish that $\tau_E(\mathcal{P}) = o(n^2)$ if \mathcal{F} contains a bipartite graph.

The proof uses the game of JumbleG and the Szemerédi Regularity Lemma.

1 Introduction

An **unbiased positional game** is one in which two players alternately select a vertex from a hypergraph. In the most common formulation of the game, one player is Maker and the other Breaker. Maker attempts to occupy all vertices in some hyperedge and Breaker attempts to occupy at least one vertex in every hyperedge. In the graph context, players select edges of a complete graph K_n or possibly some other graph, such as a random graph [8]. Here Maker attempts to create a graph with a given monotone property and Breaker attempts to prevent Maker from achieving this.

Hefetz, Krivelevich and Szabó [6] recently investigated the so-called **Avoider-Enforcer game**. The players are, unsurprisingly, Avoider and Enforcer. In the context of graph games, Avoider attempts to prevent having her edges induce a graph with a property \mathcal{P} for as many rounds as possible. Enforcer selects his edges in such a way as to force **Avoider's graph** to have property \mathcal{P} as early as possible.

*Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL, 61801, jobal@math.uiuc.edu. This author's research supported in part by NSF grants DMS-0302804, DMS-0600303 and DMS-0603769, UIUC Campus Research Board #07048 and by OTKA grants T034475 and T049398.

†Department of Mathematics, Iowa State University, Ames, IA 50011, rymartin@iastate.edu. This author's research supported in part by NSA grant H98230-05-1-0257.

At the end of r rounds, Avoider and Enforcer each have chosen exactly r edges. Let $\tau_E(\mathcal{P})$ be the smallest integer t such that Enforcer can win the \mathcal{P} -property game in t rounds. Let \mathcal{C}_n^k denote the property that an n -vertex graph is k -colorable and let \mathcal{NC}_n^k denote the property that an n -vertex graph is not k -colorable.

In [5], Hefetz, Krivelevich, Stojaković and Szabó establish that

$$\frac{n^2}{8} + \frac{n-2}{12} \leq \tau_E(\mathcal{NC}_n^2) \leq \frac{n^2}{8} + \frac{n}{2} + 1,$$

but for $k \geq 3$ show only that

$$(1 - o(1)) \frac{(k-1)n^2}{4k} \leq \tau_E(\mathcal{NC}_n^k) < \frac{1}{2} \binom{n}{2}.$$

They state that it “seems reasonable” that $\tau_E(\mathcal{NC}_n^k) \leq (1 + o(1)) \frac{(k-1)n^2}{4k}$.

In this paper, we prove that this is the case, as a consequence of a stronger result.

1.1 Main Results

Let $t(n, k)$ denote the Turán number that is the maximum number of edges in a graph with no K_{k+1} . In particular, $\binom{k-1}{k} \frac{n^2}{2} - \frac{k}{8} \leq t(n, k) \leq \binom{k-1}{k} \frac{n^2}{2}$.

Definition 1 Let \mathcal{F} be a family of graphs such that

$$k = \min\{\chi(F) : F \in \mathcal{F}\}.$$

We use the notation \mathcal{F}_n to be the property that a graph on n vertices has a copy of some member of \mathcal{F} as a subgraph and let $\mathcal{F}_n^{\text{ind}}$ be the property that a graph on n vertices has a copy of some member of \mathcal{F} as an *induced* subgraph.

Theorem 2 In the Avoider-Enforcer game for properties \mathcal{F}_n and $\mathcal{F}_n^{\text{ind}}$, if n is large enough, then

$$\left(\frac{k-2}{k-1}\right) \frac{n^2}{4} - O(k) = \left\lfloor \frac{1}{2} t(n, k-1) \right\rfloor \leq \tau_E(\mathcal{F}_n) \leq \tau_E(\mathcal{F}_n^{\text{ind}}) \leq (1 + o(1)) \left(\frac{k-2}{k-1}\right) \frac{n^2}{4}.$$

The last inequality in Theorem 2 is our main result. Enforcer’s strategy is simple, he tries to achieve to maintain that the board “looks like a random graph” after each round. We simply show that in this case Avoider’s graph when it has $(1 + o(1)) \frac{k-2}{k-1} \left(\frac{n^2}{4}\right)$ edges, will satisfy property $\mathcal{F}_n^{\text{ind}}$.

Let $\mathcal{F} = \{K_{k+1}\}$. The upper bound for Theorem 3 is an immediate consequence of Theorem 2. The lower bound comes from the same trivial strategy that establishes the lower bound in Theorem 2.

Theorem 3 *Let $k \geq 2$ be an integer and let \mathcal{NC}_n^k be the property that a graph is not k -colorable. In the Avoider-Enforcer game,*

$$\left(\frac{k-1}{k}\right) \frac{n^2}{4} - O(k) = \left\lfloor \frac{1}{2}t(n, k) \right\rfloor \leq \tau_E(\mathcal{NC}_n^k) \leq (1 + o(1)) \left(\frac{k-1}{k}\right) \frac{n^2}{4}.$$

This result is weaker than that in [5] for $k = 2$, but gives the correct asymptotic approximation for all $k \geq 2$.

This paper is organized as follows: In Section 2, we describe the game of JumbleG [4], along with important consequences thereof. In Section 3, we state the version of the Szemerédi Regularity Lemma that is useful for our purposes. Section 4 contains the proofs of the theorems. Section 5 contains some concluding remarks.

2 The game of JumbleG

The game of JumbleG is a traditional Maker-Breaker game. The goal of Maker is to create a pseudo-random graph. Frieze, Krivelevich, Pikhurko and Szabó [4] use two different conditions for pseudo-randomness. We will use their first definition.

Definition 4 *If G is a graph and S, T are disjoint vertex sets in $V(G)$, then denote $e_G(S, T)$ to be the number of edges which have one endpoint in S and the other in T .*

A pair of disjoint vertex sets (S, T) is ϵ -unbiased if

$$\left| \frac{e_G(S, T)}{|S||T|} - \frac{1}{2} \right| \leq \epsilon.$$

A graph G on n vertices, with minimum degree $\delta(G)$, is ϵ -regular if both:

P1 $\delta(G) \geq (1/2 - \epsilon)n$.

P2 *Any pair S, T of disjoint subsets of $V(G)$ with $|S|, |T| > \epsilon n$ is ϵ -unbiased.*

Theorem 5 ([4]) *Maker has a winning strategy in JumbleG, provided $\epsilon \geq 2(\log n/n)^{1/3}$ and n is sufficiently large.*

In our results, Enforcer will simply play as Maker in the game of JumbleG. As a result, between any pair of disjoint sets, both of them large enough, Avoider cannot occupy too many of the edges. We formalize this as follows.

Definition 6 *After round r in the game let $e_B(S, T; r)$ denote the number of edges that Breaker occupies in the pair (S, T) and $e_M(S, T; r)$ denote the number of edges that Maker occupies.*

Lemma 7 *Let $S, T \subset V(G)$ be disjoint sets and $|S|, |T| \geq \epsilon n$. If Maker plays the strategy to win JumbleG, then for every r*

$$e_B(S, T; r) - e_M(S, T; r) \leq 2\epsilon|S||T| + 1.$$

Proof. Fix any integer $r > 0$. Suppose $e_B(S, T; r) - e_M(S, T; r) > 2\epsilon|S||T| + 1$. Assume, at this stage, Breaker can change his strategy to occupy as many edges between S and T as possible. When the game concludes, the number of edges that Breaker occupies between S and T is at least

$$\begin{aligned} & e_B(S, T; r) + \left\lfloor \frac{1}{2} (|S||T| - e_B(S, T; r) - e_M(S, T; r)) \right\rfloor \\ & \geq e_B(S, T; r) + \frac{1}{2} (|S||T| - e_B(S, T; r) - e_M(S, T; r)) - \frac{1}{2} \\ & > \frac{1}{2}|S||T| + \frac{1}{2} (2\epsilon|S||T| + 1) - \frac{1}{2} \\ & \geq \left(\frac{1}{2} + \epsilon \right) |S||T|. \end{aligned}$$

This contradicts the assumption that Maker played a winning strategy for JumbleG. ■

3 The Regularity Lemma

Definition 8 *Let A and B be disjoint vertex sets. The number of edges with one endpoint in A and the other in B is denoted by $e(A, B)$. The **density** of (A, B) is denoted*

$$d(A, B) = \frac{e(A, B)}{|A||B|}.$$

A pair (A, B) is α -regular if, for every $X \subseteq A$ and $Y \subseteq B$ with $|X| > \alpha|A|$ and $|Y| > \alpha|B|$,

$$|d(A, B) - d(X, Y)| < \alpha.$$

*The partition V_1, \dots, V_ℓ is an **equipartition** of V if for every i, j we have $||V_i| - |V_j|| \leq 1$.*

We use a form of the Szemerédi Regularity Lemma, introduced by Alon, Fischer, Krivelevich and M. Szegedy [1]. Lemma 9 is a simplified version.

It is possible to prove our results using the original version [9] but we use this version, as its statement is convenient for our purposes.

Lemma 9 (Regularity Lemma [1]) *For every integer m and constant $\mathcal{E} > 0$, there is an $S = S(m, \mathcal{E})$ which satisfies the following: For any graph G on $n \geq S$ vertices, there exists an equipartition $\mathcal{A} = \{V_i : 1 \leq i \leq \ell\}$ of $V(G)$ and an induced subgraph U of G , with an equipartition $\mathcal{B} = \{U_i : 1 \leq i \leq \ell\}$ of the vertices of U , that satisfy:*

- $m \leq \ell \leq S$.
- $U_i \subseteq V_i$ for all $i \geq 1$, and $|U_i| = L \geq \lceil n/S \rceil$.
- In the equipartition \mathcal{B} , **all** pairs are \mathcal{E} -regular.
- All but at most $\mathcal{E} \binom{\ell}{2}$ of the pairs $1 \leq i < j \leq \ell$ are such that $|d(V_i, V_j) - d(U_i, U_j)| < \mathcal{E}$.

Since we want to establish that a large enough density in G will enable us to apply Turán's theorem, we need to bound $e(U)$ in terms of $e(G)$.

Corollary 10 *Let m be an integer, and \mathcal{E} be a constant such that $m \leq \mathcal{E}^{-1}$ and let $S = S(m, \mathcal{E})$ be the integer provided by Lemma 9. Let $n \geq 2S/\mathcal{E}$, let G be a graph on n vertices. Let U be the induced subgraph provided by Lemma 9. If the equipartition of $V(U)$ is $\{U_i\}_{i=1}^{\ell}$ with $|U_1| = \dots = |U_{\ell}| = L$, then let \tilde{U} be the graph formed by deleting all edges which have both endpoints in the same set U_i , $i = 1, \dots, \ell$. In this case,*

$$e(U) \geq e(\tilde{U}) \geq e(G) \frac{\ell^2 L^2}{n^2} - 3\mathcal{E} \ell^2 L^2.$$

Proof. First, we bound $e(G)$, using the conditions in Lemma 9.

$$\begin{aligned} e(G) &\leq \sum_i \binom{|V_i|}{2} + \sum_{1 \leq i < j \leq \ell} e(V_i, V_j) \\ &\leq \ell \frac{\lceil n/\ell \rceil^2}{2} + \mathcal{E} \binom{\ell}{2} \left[\left\lceil \frac{n}{\ell} \right\rceil^2 + \left\lceil \frac{n}{\ell} \right\rceil^2 \sum_{1 \leq i < j \leq \ell} [d(U_i, U_j) + \mathcal{E}] \right] \\ &\leq \frac{\ell}{2} \left\lceil \frac{n}{\ell} \right\rceil^2 + 2\mathcal{E} \binom{\ell}{2} \left[\left\lceil \frac{n}{\ell} \right\rceil^2 + \left\lceil \frac{n}{\ell} \right\rceil^2 \sum_{1 \leq i < j \leq \ell} d(U_i, U_j) \right] \\ &\leq \frac{\ell}{2} \left\lceil \frac{n}{\ell} \right\rceil^2 + \mathcal{E} \ell^2 \left\lceil \frac{n}{\ell} \right\rceil^2 + \frac{\lceil n/\ell \rceil^2}{L^2} e(\tilde{U}). \end{aligned} \tag{1}$$

Recall that $L = |U_1| = \dots = |U_\ell|$. The calculations below, use the fact that $\ell^{-1} \leq m^{-1} \leq \mathcal{E}$ and $\ell/n \leq S/n \leq \mathcal{E}/2$. By rearranging the terms in inequality (1), we get a lower bound for $e(\tilde{U})$.

$$\begin{aligned}
e(\tilde{U}) &\geq e(G) \frac{L^2}{\lceil n/\ell \rceil^2} - \frac{\ell L^2}{2} - \mathcal{E} \ell^2 L^2 \\
&\geq e(G) \frac{\ell^2 L^2}{n^2} - e(G) \left(\frac{\ell^2 L^2}{n^2} - \frac{L^2}{\lceil n/\ell \rceil^2} \right) - \frac{1}{2\ell} \ell^2 L^2 - \mathcal{E} \ell^2 L^2 \\
&\geq e(G) \frac{\ell^2 L^2}{n^2} - \frac{n^2}{2} \left(\frac{\ell^2 L^2}{n^2} - \frac{\ell^2 L^2}{(n+\ell)^2} \right) - \frac{\mathcal{E}}{2} \ell^2 L^2 - \mathcal{E} \ell^2 L^2 \\
&\geq e(G) \frac{\ell^2 L^2}{n^2} - \left(1 - \frac{n^2}{(n+\ell)^2} \right) \frac{\ell^2 L^2}{2} - \frac{\mathcal{E}}{2} \ell^2 L^2 - \mathcal{E} \ell^2 L^2 \\
&\geq e(G) \frac{\ell^2 L^2}{n^2} - 3\mathcal{E} \ell^2 L^2.
\end{aligned}$$

Trivially, $e(U) \geq e(\tilde{U})$ and this concludes the proof. \blacksquare

Lemma 11 establishes that a regular f -tuple will induce a graph on f vertices, given necessary density conditions. Versions of it have appeared, for example, in Komlós and Simonovits [7], Bollobás and Thomason [3] and Alon and Shapira [2].

Lemma 11 *For every real η , $0 < \eta < 1$, and integer $f \geq 1$, there exists a $\gamma = \gamma(\eta, f)$ with the following property: Suppose that H is a graph on f vertices v_1, \dots, v_f , and that U_1, \dots, U_f is an f -tuple of disjoint nonempty vertex sets of a graph G such that for every $1 \leq i < j \leq f$, the pair (U_i, U_j) is γ -regular. Moreover, whenever $(v_i, v_j) \in E(H)$ we have $d(U_i, U_j) \geq \eta$ and whenever $(v_i, v_j) \notin E(H)$ we have $d(U_i, U_j) \leq 1 - \eta$. Then, some f -tuple $u_1 \in U_1, \dots, u_f \in U_f$ spans an **induced** copy of H , where each u_i plays the role of v_i .*

The Slicing Lemma (Fact 1.5 in [7]) is a basic fact of regular pairs, common in proofs involving the Regularity Lemma.

Lemma 12 (Slicing Lemma) *Let (U_i, U_j) be an α -regular pair with density d and $|U_i| = |U_j| = L_0$. If $X \subseteq U_i$ and $Y \subseteq U_j$ with $|X| = L_i$ and $|Y| = L_j$, then (X, Y) is α' -regular, where*

$$\alpha' = \max \left\{ 2\alpha, \frac{L_0}{L_i} \alpha, \frac{L_0}{L_j} \alpha \right\},$$

with density in $(d - \alpha, d + \alpha)$.

4 Proof of Theorem 2

As in many theorems involving the Regularity Lemma, there is a sequence of constants. Fix an $F \in \mathcal{F}$ with chromatic number k and order f . With $a \ll b$ meaning that a is small enough relative to b , the

constants are

$$\epsilon \ll \mathcal{E}_0 \ll \mathcal{E}_1 \ll \eta \ll \delta \ll f^{-1}.$$

We will determine the precise relationships later.

In order to establish the lower bound for $\tau_E(\mathcal{F}_n)$, Avoider equipartitions the n vertices into $k-1$ clusters, and choose edges only between different clusters. By this strategy, Avoider's graph will always be $(k-1)$ -colorable, thus, have no member of \mathcal{F} as a subgraph. Avoider can make at least $\lfloor (1/2)t(n, k-1) \rfloor$ moves. Careful calculations establish that

$$t(n, k) = \left(\frac{k-1}{k} \right) \frac{n^2}{2} - \frac{k}{2} \left(\left\lceil \frac{n}{k} \right\rceil - \frac{n}{k} \right) \left(\frac{n}{k} - \left\lfloor \frac{n}{k} \right\rfloor \right).$$

Since the existence of an induced copy of some $F \in \mathcal{F}$ implies the existence of a copy of F as a subgraph, it is trivial that $\tau_E(\mathcal{F}_n) \leq \tau_E(\mathcal{F}_n^{\text{ind}})$.

Finally, we shall prove the upper bound. Among all $F \in \mathcal{F}$ choose an F with $\chi(F) = k$ and let $f = |V(F)|$. Enforcer will play the game JumbleG, and we shall prove that Avoider's graph, after $\left(\frac{k-2}{4(k-1)} + o(1) \right) n^2$ rounds will contain F as an induced subgraph. Assume, for the sake of contradiction, that for (a small) $\delta > 0$, Avoider managed to build a graph G of order n (where n is sufficiently large) and $\left(\frac{k-2}{4(k-1)} + 2\delta \right) n^2$ edges. Apply the Regularity Lemma (Lemma 9) to Avoider's graph with parameters \mathcal{E}_0 and $m := \max\{k, \lceil \mathcal{E}_0^{-1} \rceil\}$, (where \mathcal{E}_0 will be determined later) which yields the subsets U_1, \dots, U_{ℓ_0} and a constant $S_0 = S(m, \mathcal{E}_0)$ so that $|U_1| = \dots = |U_{\ell_0}| = L_0 > n/S_0$. Construct an auxiliary graph H on the vertex set $\{1, \dots, \ell_0\}$ where $i \sim j$ if and only if $d(U_i, U_j) \geq \delta + \epsilon$. Let \tilde{U} be the graph induced by $U_1 \cup \dots \cup U_{\ell_0}$, with the edges inside of each cluster deleted. We use $e(\tilde{U})$ in order to compute $e(H)$. Observe that Lemma 7 gives that $d(U_i, U_j) \leq (1/2 + \epsilon)$ as long as $L_0 \geq n/S_0 > \epsilon n$.

$$\begin{aligned} e(\tilde{U}) &\leq e(H) \left(\frac{1}{2} + \epsilon \right) L_0^2 + \left[\binom{\ell_0}{2} - e(H) \right] (\delta + \epsilon) L_0^2 \\ &\leq e(H) \left(\frac{1}{2} - \delta \right) L_0^2 + \frac{\delta + \epsilon}{2} \ell_0^2 L_0^2. \end{aligned}$$

We use Corollary 10 to bound $e(\tilde{U})$ by $e(G)$.

$$\begin{aligned} e(H) &\geq \frac{e(\tilde{U}) - \frac{\delta + \epsilon}{2} \ell_0^2 L_0^2}{\left(\frac{1}{2} - \delta \right) L_0^2} \\ &\geq \frac{e(G) \frac{\ell_0^2}{n^2} - 3\mathcal{E}_0 \ell_0^2 - \frac{\delta + \epsilon}{2} \ell_0^2}{1/2 - \delta}. \end{aligned} \tag{2}$$

If

$$2\mathcal{E}_0 + 3\epsilon/2 < \delta, \tag{3}$$

then

$$e(G) \geq \left(\frac{k-2}{4(k-1)} + 2\delta \right) n^2 > \left(\frac{k-2}{4(k-1)} + \delta + 2\mathcal{E}_0 + \frac{\epsilon}{2} \right) n^2.$$

Plugging this into (2), we obtain that $e(H) > \left(\frac{k-2}{k-1} \right) \frac{\ell_0^2}{2}$. Therefore, H contains a K_k by Turán's theorem. Without loss of generality, this copy of K_k is spanned by (U_1, \dots, U_k) . Each pair (U_i, U_j) is \mathcal{E}_0 -regular with density in the interval $(\delta + \epsilon, 1/2 + \epsilon)$. \blacksquare

Note: At this stage, we can use, say, the Blow-up lemma (see [7]) to show that if n is large enough, a not necessary induced copy of F occurs as a subgraph in (U_1, \dots, U_k) , hence in G . This provides an upper bound for $\tau_E(\mathcal{F}_n)$. The rest of the proof is required to obtain the bound for $\tau_E(\mathcal{F}_n^{\text{ind}})$.

We apply the Regularity Lemma (Lemma 9) to the portion of Avoider's graph in U_i with parameters \mathcal{E}_1 and $m = f$, for each $i = 1, \dots, k$ (where \mathcal{E}_1 will be determined later), which yields the constants ℓ_i , the subclasses $U_{i,1}, \dots, U_{i,\ell_i}$ and a constant $S_1 = S(m, \mathcal{E}_1)$ so that $|U_{i,1}| = \dots = |U_{i,\ell_i}| = L_i \geq L_0/S_1 \geq n/(S_0 S_1)$. If

$$\epsilon \leq S_0^{-1} S_1^{-1} \tag{4}$$

then each pair $(U_{i,x}, U_{j,y})$ has density at most $1/2 + \epsilon$, by Lemma 7.

Each pair $(U_{i,x}, U_{i,y})$ is \mathcal{E}_1 -regular. Now consider a pair $(U_{i,x}, U_{j,y})$ where $i \neq j$. The pair (U_i, U_j) is \mathcal{E}_0 -regular with density at least $\delta + \epsilon$. Using the Slicing Lemma (Lemma 12), the pair $(U_{i,x}, U_{j,y})$ is $\mathcal{E}_0 L_0 \cdot \max\{L_i^{-1}, L_j^{-1}\}$ -regular. Since $L_i, L_j \geq L_0/S_1$, the pair $(U_{i,x}, U_{j,y})$ is $\mathcal{E}_0 S_1$ -regular with density at most $1/2 + \epsilon$ and at least $\delta/2$, as long as

$$\mathcal{E}_0 \leq \delta/2 + \epsilon. \tag{5}$$

Finally, we apply Lemma 11 with $\eta = \delta/2$ to F and the tuple $\{U_{i,x}\}_{1 \leq i \leq k, 1 \leq x \leq \ell_i}$. This implies the existence of the constant $\gamma = \gamma(f, \delta/2)$. As F is k -colorable graph of order f , and $f \leq \ell_i$ for every i , all pair of classes are $\mathcal{E}_0 S_1$ -regular. In order to apply Lemma 11, it is sufficient to have

$$\mathcal{E}_0 S_1 \leq \gamma(f, \delta/2). \tag{6}$$

This way F can be embedded into G , Avoider's graph.

The order of choosing the constants. First a forbidden graph F is fixed with chromatic number k , and of order f , and an arbitrary $\delta > 0$ is chosen, assuming that Avoider built a graph with $\left(\frac{k-2}{4(k-1)} + 2\delta \right) n^2$ edges. Then from Lemma 11, with $\eta = \delta/2$ we obtain the existence of a $\gamma < m^{-1}$. The Regularity Lemma (Lemma 9) is applied to the classes U_i with $\mathcal{E}_1 = \gamma$ and $m = f$. This also implies the existence of an $S_1 = S_1(\mathcal{E}_1, m)$. We need that the pairs $(U_{i,x}, U_{j,y})$ are γ -regular, hence $\mathcal{E}_0 S_1 \leq \gamma$ needed, so a positive constant $\mathcal{E}_0 < \gamma/S_1$ is chosen. The Regularity Lemma (Lemma 9) is

applied to G with \mathcal{E}_0 and $m = f$. This also implies the existence of an $S_0 = S_0(\mathcal{E}_0, m)$. For JumbleG we use $\epsilon = S_0^{-1}S_1^{-1}$. In order to apply all these Lemmas, we need to assume that n is sufficiently large. This specific choice of constants satisfies the inequalities (3), (4), (5) and (6).

To summarize, we proved that for every $\delta > 0$, if the Avoider-Enforcer induced- \mathcal{F} -avoidance game is played for $\left(\frac{k-2}{4(k-1)} + 2\delta\right)n^2$ rounds, then the Avoider's graph will contain an induced F , and the Enforcer will win. Thus, $\tau_E(\mathcal{F}_n^{\text{ind}}) \leq (1 + o(1))\left(\frac{k-2}{k-1}\right)\frac{n^2}{4}$.

5 Concluding remarks

The game \mathcal{NC}_n^1 is trivial, but the \mathcal{F}_n question is not trivial for $k = 2$. That is, the case in which \mathcal{F} contains a bipartite graph. Theorem 2 gives only that, if $k = 2$ and $\mathcal{F} = \{F\}$, where F is a bipartite graph, then

$$\left\lfloor \frac{1}{2}t(n, F) \right\rfloor \leq \tau_E(\mathcal{F}_n) \leq \tau_E(\mathcal{F}_n^{\text{ind}}) \leq o(n^2),$$

which is not very helpful. And certainly in the noninduced case, we have trivially

$$\tau_E(\mathcal{F}_n) \leq t(n, F).$$

The authors believe the approach needed to prove Theorem 2 may solve other problems in the area of positional games.

We also remark that one can prove the non-induced case of $k = 3$ without the use of the Szemerédi Regularity Lemma.

Acknowledgements: We would like to thank Mathematisches Forschungsinstitut Oberwolfach for hosting a miniworkshop on positional games and Danny Hefetz, Michael Krivelevich, Miloš Stojaković, and Tibor Szabó for their helpful remarks.

References

- [1] N. Alon, E. Fischer, M. Krivelevich and M. Szegedy, Efficient testing of large graphs, *Combinatorica* **20** (2000), 451–476.
- [2] N. Alon and A. Shapira, A characterization of the (natural) graph properties testable with one-sided error, Proc. of the 46 IEEE FOCS, IEEE (2005), 429–438.
- [3] B. Bollobás and A. Thomason, Hereditary and monotone properties of graphs, in “The Mathematics of Paul Erdős II” (R.L. Graham and J. Nešetřil, eds.) Algorithms and Combinatorics **14** Springer-Verlag (1997), 70–78.

- [4] A. Frieze, M. Krivelevich, O. Pikhurko and T. Szabó, The game of JumbleG, *Combin. Probab. Comput.* **12** (2005), R51 (electronic).
- [5] D. Hefetz, M. Krivelevich, M. Stojaković and T. Szabó, Fast winning strategies in positional games, preprint.
- [6] D. Hefetz, M. Krivelevich and T. Szabó, Avoider-Enforcer games, *J. Combin. Th. Ser. A*, **114** (2007), 840–853.
- [7] J. Komlós and M. Simonovits, Szemerédi’s Regularity Lemma and its applications in graph theory. In: *Combinatorics, Paul Erdős is Eighty*, Vol. II (D. Miklós, V.T. Sós, T. Szőnyi eds.), János Bolyai Math. Soc., Budapest (1996), 295–352.
- [8] M. Stojaković and T. Szabó, Positional games on random graphs, *Random Structures Algorithms*, **26** (2005) 204–223.
- [9] E. Szemerédi, Regular partitions of graphs, In: *Proc. Colloque Inter. CNRS* (J.C. Bermond, J.C. Fournier, M. Las Vergnas and D. Sotteau, eds.), 1978, 399–401.