

# On the number of $K_4$ -saturating edges

József Balogh <sup>\*</sup>      Hong Liu <sup>†</sup>

May 15, 2013

## Abstract

Let  $G$  be a  $K_4$ -free graph, an edge in its complement is a  $K_4$ -saturating edge if the addition of this edge to  $G$  creates a copy of  $K_4$ . Erdős and Tuza conjectured that for any  $n$ -vertex  $K_4$ -free graph  $G$  with  $\lfloor n^2/4 \rfloor + 1$  edges, one can find at least  $(1 + o(1))\frac{n^2}{16}$   $K_4$ -saturating edges. We construct a graph with only  $\frac{2n^2}{33}$   $K_4$ -saturating edges. Furthermore, we prove that it is best possible, i.e., one can always find at least  $(1 + o(1))\frac{2n^2}{33}$   $K_4$ -saturating edges in an  $n$ -vertex  $K_4$ -free graph with  $\lfloor n^2/4 \rfloor + 1$  edges.

## 1 Introduction

The notation in this paper is standard. For a graph  $G$ , denote by  $\overline{G}$  its complement. For vertex subsets  $U, W \subseteq V(G)$ , denote  $N(U) := \bigcap_{v \in U} N(v)$  and  $E(U, W)$  the set of cross edges between  $U$  and  $W$ .

Mantel [14] showed that the maximum number of edges in an  $n$ -vertex triangle-free graph is  $\lfloor n^2/4 \rfloor$ . Rademacher (unpublished) extended this result by showing that any  $n$ -vertex graph with  $\lfloor n^2/4 \rfloor + t$  edges contains at least  $t\lfloor n/2 \rfloor$  triangles, for  $t = 1$ . Lovász and Simonovits [13], improving Erdős [6], proved this for every  $t \leq n/2$ . Erdős [7] showed analogue results for cliques and Mubayi [15, 16] for color-critical graphs and for some hypergraphs.

In general, we call Erdős-Rademacher-type problem the following: for any extremal question, what is the number of forbidden configurations appearing in a graph somewhat denser than the extremal graph? This type of problems have been studied in various contexts: A *book* of size  $q$  consists of  $q$  triangles sharing a common edge. Khadžiivanov and Nikiforov [11], answering a question of Erdős, showed that any  $n$ -vertex graph with  $\lfloor n^2/4 \rfloor + 1$  edges contains a book of size at least  $n/6$ . In the context of Sperner's Theorem, Kleitman [12], answering a question of Erdős and Katona, determined the minimum number of 2-chains a poset must

---

<sup>\*</sup>Department of Mathematical Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA [jobal@math.uiuc.edu](mailto:jobal@math.uiuc.edu). Research is partially supported by Simons Fellowship, NSF CAREER Grant DMS-0745185 and Arnold O. Beckman Research Award (UIUC Campus Research Board 13039).

<sup>†</sup>Department of Mathematical Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA [hliu36@illinois.edu](mailto:hliu36@illinois.edu).

contain if its size is larger than its largest anti-chain. Recently, this theorem was extended to  $k$ -chains by Das, Gan and Sudakov [5].

Let  $G$  be an  $n$ -vertex  $K_4$ -free graph, an edge in  $\overline{G}$  is a  $K_4$ -saturating edge if the addition of this edge to  $G$  creates a copy of  $K_4$ . Denote by  $f(G)$  the number of  $K_4$ -saturating edges in  $\overline{G}$  and by  $f(n, e)$  the maximum integer  $\ell$  such that every  $n$ -vertex  $K_4$ -free graph with  $e$  edges must have at least  $\ell$   $K_4$ -saturating edges. The first extremal result related to clique-saturating edges was by Bollobás [4] who proved that if every edge in  $\overline{G}$  is a  $K_r$ -saturating edge, then  $e(G) \geq \binom{n}{2} - \binom{n-r+2}{2}$  and this bound is best possible. Later it was extended by Alon [1], Frankl [9] and Kalai [10] using linear algebraic method. Recently, saturation problems were phrased in the language of ‘graph bootstrap percolation’, see [2] and [3].

In the case of  $K_4$ , Bollobás’ example is the following: let  $F$  be an  $n$ -vertex  $K_4$ -free graph with two vertices adjacent to all other vertices which form an independent set. This graph has only linear many edges,  $e(F) = 2n - 3$ , and yet all edges in  $\overline{F}$  are  $K_4$ -saturating edges. To the other extreme,  $K_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}$  shows that a graph could have up to  $\lfloor n^2/4 \rfloor$  edges with no  $K_4$ -saturating edge, i.e.  $f(n, \lfloor n^2/4 \rfloor) = 0$ . Erdős and Tuza [8] conjectured that if a  $K_4$ -free graph  $G$  has  $\lfloor n^2/4 \rfloor + 1$  edges, then suddenly there are quadratic many  $K_4$ -saturating edges. This conjecture can be considered, as formulated before, an Erdős-Rademacher-type problem concerning the number of  $K_4$ -saturating edges.

**Conjecture 1.1** (Erdős-Tuza [8]).

$$f(n, \lfloor n^2/4 \rfloor + 1) = (1 + o(1)) \frac{n^2}{16}.$$

We disprove this conjecture. We give a counterexample with only  $\frac{2n^2}{33}$   $K_4$ -saturating edges. Furthermore, we prove that  $(1 + o(1)) \frac{2n^2}{33}$  is best possible, that is, one can always find at least  $(1 + o(1)) \frac{2n^2}{33}$   $K_4$ -saturating edges in an  $n$ -vertex  $K_4$ -free graph with  $\lfloor n^2/4 \rfloor + 1$  edges.

**Theorem 1.2.** For  $n \geq 73$ ,

$$f(n, \lfloor n^2/4 \rfloor + 1) = \frac{2n^2}{33} + O(n).$$

We shall prove the following theorem, which implies the lower bound in Theorem 1.2.

**Theorem 1.3.** Let  $G$  be an  $n$ -vertex  $K_4$ -free graph with  $\lfloor n^2/4 \rfloor$  edges, for  $n \geq 73$ . If  $G$  contains a triangle, then

$$f(G) \geq \frac{2n^2}{33} - \frac{3n}{11}.$$

This is best possible when  $n$  is divisible by 66.

*Proof of Theorem 1.2.* The upper bound is by the construction described in Section 2. For the lower bound, let  $G$  be a  $K_4$ -free graph with  $\lfloor n^2/4 \rfloor + 1$  edges. By Mantel’s theorem, it contains a triangle. Let  $G'$  be a subgraph obtained from  $G$  by removing an edge such that  $G'$  contains a triangle. By Theorem 1.3,  $f(G') \geq \frac{2n^2}{33} - \frac{3n}{11}$ . The relation  $f(G) \geq f(G')$  completes the proof.  $\square$

**Remark:** (i) Slight modification of our proof gives the following stability result: Given any  $K_4$ -free graph  $G$  with  $(1 - o(1))n^2/4$  edges, if  $G$  contains a triangle, then  $f(G) \geq (1 - o(1))2n^2/33$ .

(ii) Unlike the case about the number of triangles in [6] and [13], where every additional edge, up to  $n/2$ , guarantees  $\lfloor n/2 \rfloor$  additional triangles, in our problem, even with linear many extra edges, the number of  $K_4$ -saturating edges is still at most  $(1 + o(1))2n^2/33$ . In particular,  $f\left(n, \lfloor \frac{n^2}{4} \rfloor + t\right) = \frac{2n^2}{33} + O(n)$  for  $1 \leq t \leq \frac{n}{66}$ .

(iii) One might define a  $K_r$ -saturating edge of a graph  $G$ , for  $r \geq 5$ , as we did for  $K_4$ . Denote by  $\text{ex}(n, K_{r-1})$  the maximum size of an  $n$ -vertex  $K_{r-1}$ -free graph. We think that a similar phenomenon holds: if  $G$  is  $K_r$ -free and  $e(G) = \text{ex}(n, K_{r-1}) + 1$ , then the number of  $K_r$ -saturating edges is at least  $\left(\frac{2(r-3)^2}{(r-1)(4r^2-19r+23)} + o(1)\right)n^2$ . A straightforward generalization of our construction shows that if the conjecture is true, then it is best possible. Some of the ideas of our proof works for  $r \geq 5$  as well, but some does not.

The paper is organized as follows: We give a construction for the upper bound in Theorem 1.2 and an extremal example for Theorem 1.3 in Section 2. The proof for Theorem 1.3 is given in Section 3. We will omit floors and ceilings when it is not critical and we make no effort optimizing some of the constants.

## 2 Upper bound constructions

Fix an integer  $n$  divisible by 66. We present an  $n$ -vertex  $K_4$ -free graph  $H$  with  $n^2/4 + n/66$  edges and  $f(H) = 2n^2/33 - 7n/33$ . Note that from this graph one can easily remove  $n/66 - 1$  edges without changing the number of  $K_4$ -saturating edges. We also give an extremal example showing the bound in Theorem 1.3 is best possible.

**Construction for Theorem 1.2:** To construct  $H$ , start with a  $C_5$  on  $\{v_1, v_2, v_3, v_4, v_5\}$  with a chord  $v_1v_3$ . Blow up each  $v_i$  to an independent set  $V_i$  of the following size:  $|V_1| = |V_3| = 16n/66$ ,  $|V_2| = 4n/66 + 1$ ,  $|V_4| = 15n/66$  and  $|V_5| = 15n/66 - 1$ , see Figure 1. Then  $H$  is  $K_4$ -free with  $n^2/4 + n/66$  edges. The only  $K_4$ -saturating edges are those in  $V_1, V_2, V_3$ , which gives  $f(H) = 2n^2/33 - 7n/33$ .

**Construction for Theorem 1.3:** Define  $H'$  the same way as  $H$ , except that  $|V_2'| = 4n/66$  and  $|V_4'| = 15n/66$ . This graph is  $K_4$ -free with  $n^2/4$  edges and  $f(H') = \frac{2n^2}{33} - \frac{3n}{11}$ .

## 3 Proof of Theorem 1.3

Let  $G$  be a  $K_4$ -free graph with  $n^2/4$  edges and containing a triangle. Fix, in  $G$ , a maximum family of vertex-disjoint triangles, say  $\mathcal{T} = \{T_1, T_2, \dots, T_t\}$ , where  $0 < t \leq 1/3$ . We write  $V(\mathcal{T})$  for  $\bigcup_{i=1}^t V(T_i)$ ,  $E(\mathcal{T})$  for  $E(G[V(\mathcal{T})])$  and  $e(\mathcal{T}) := |E(\mathcal{T})|$ . Let  $G' = G - V(\mathcal{T})$ , since  $\mathcal{T}$  is of maximum size,  $G'$  is a  $K_3$ -free graph with  $e(G') \leq \frac{(1-3t)^2 n^2}{4}$ . Denote by  $r_1 n^2$  the number of  $K_4$ -saturating edges incident to  $V(\mathcal{T})$ , and by  $r_2 n^2$  the number of  $K_4$ -saturating edges in  $V(G')$ . Hence  $f(G) = (r_1 + r_2)n^2$ . First we give a lower bound on  $r_1$ .




Figure 1: A  $K_4$ -free graph  $H$  with  $e(H) = \frac{n^2}{4} + \frac{n}{66}$  and  $f(H) = \frac{2n^2}{33} - \frac{7n}{33}$ .

**Lemma 3.1.**

$$r_1 n^2 \geq \left( \frac{1}{4} - t + \frac{3t^2}{2} \right) n^2 - e(G') - \frac{3}{2}tn \geq \left( \frac{t}{2} - \frac{3t^2}{4} \right) n^2 - \frac{3}{2}tn.$$

*Proof.* Let  $t_i = e(T_i, G \setminus \bigcup_{j=1}^i T_j)$ , clearly  $\sum_{i=1}^{tn} t_i = e(G) - e(G') - 3tn$ . Since  $G$  is  $K_4$ -free, every vertex can have at most two neighbors in each triangle. Thus  $t_i - (n - 3i)$  is a lower bound on the number of vertices in  $G \setminus \bigcup_{j=1}^i T_j$  having degree 2 in  $T_i$ , each of which gives a  $K_4$ -saturating edge. Indeed, say  $V(T_1) = \{x, y, z\}$ , and  $w \in N(x) \cap N(y)$ , then  $wz$  is a  $K_4$ -saturating edge. Thus,

$$\begin{aligned} r_1 n^2 &\geq \sum_{i=1}^{tn} (t_i - (n - 3i)) = (e(G) - e(G') - 3tn) - \left( tn^2 - 3 \frac{tn(tn+1)}{2} \right) \\ &\geq \left( \frac{1}{4} - t + \frac{3t^2}{2} \right) n^2 - e(G') - \frac{3}{2}tn \geq \left( \frac{t}{2} - \frac{3t^2}{4} \right) n^2 - \frac{3}{2}tn, \end{aligned}$$

where the last inequality follows from  $e(G') \leq \frac{(1-3t)^2 n^2}{4}$ .  $\square$

Let  $T_i \in \mathcal{T}$  be a triangle in  $\mathcal{T}$ . Denote by  $N_j(T_i) \subseteq V(G')$ , for  $0 \leq j \leq 3$ , the set of vertices in  $G'$  that has exactly  $j$  neighbors in  $T_i$ . Since  $G$  is  $K_4$ -free,  $N_3(T_i) = \emptyset$ , for every  $T_i$ 's. Further define  $p_0(T_i) = \frac{|N_0(T_i)|}{n}$ ,  $p_1(T_i) = \frac{|N_1(T_i)|}{n}$  and  $p_2(T_i) = \frac{|N_2(T_i)|}{n}$ . Thus by definition,  $p_0(T_i) + p_1(T_i) + p_2(T_i) = 1 - 3t$ .

The next lemma shows that there is a triangle  $T \in \mathcal{T}$  with large  $|N_2(T)|$ .

**Lemma 3.2.** *There exists a triangle  $T \in \mathcal{T}$ , such that*

- (i)  $e(T, G') \geq \left( \frac{3}{2} - \frac{21t}{4} \right) n$ , and
- (ii)  $p_2(T) \geq \frac{1}{2} - \frac{9t}{4} + p_0(T)$ .

*Proof.* (i) The edge set of  $G$  can be partitioned into  $E(G')$ ,  $E(\mathcal{T}, G')$  and  $E(\mathcal{T})$ . Notice that since  $G$  is  $K_4$ -free, there are at most 6 edges between any pair of triangles in  $\mathcal{T}$ . Hence  $e(\mathcal{T}) \leq 3tn + 6\binom{tn}{2} = 3t^2n^2$ .

Thus we have  $e(\mathcal{T}, G') = e(G) - e(G') - e(\mathcal{T}) \geq \frac{n^2}{4} - \frac{(1-3t)^2n^2}{4} - 3t^2n^2 \geq \left(\frac{3t}{2} - \frac{21t^2}{4}\right)n^2$ .

Therefore, there exists a triangle  $T \in \mathcal{T}$  with  $e(T, G') \geq e(\mathcal{T}, G')/(tn) \geq \left(\frac{3}{2} - \frac{21t}{4}\right)n$ .

(ii) Let  $T \in \mathcal{T}$  be a triangle satisfying (i). Note that  $2p_2(T) + p_1(T) = \frac{e(T, G')}{n}$ . Using  $p_0(T) + p_1(T) + p_2(T) = 1 - 3t$ , we have  $p_2(T) - p_0(T) \geq \frac{3}{2} - \frac{21t}{4} - (1 - 3t) = \frac{1}{2} - \frac{9t}{4}$ .  $\square$

From now on, we let  $T = \{x, y, z\}$  be a triangle in  $\mathcal{T}$  sending the most edges to  $G'$ , hence it has the two properties of Lemma 3.2. For brevity we write  $p_j = p_j(T)$  and  $N_i = N_i(T)$  for  $0 \leq j \leq 2$ . Furthermore, define  $A = N_{G'}(xy)$ ,  $B = N_{G'}(yz)$  and  $C = N_{G'}(xz)$ . Note that  $A, B, C$  are pairwise disjoint independent sets, otherwise  $T \cup A \cup B \cup C$  contains a copy of  $K_4$ . Define  $N_x := N_{G'}(x)$ ,  $N_y := N_{G'}(y)$  and  $N_z := N_{G'}(z)$ . Let  $a = \frac{|A|}{|N_2|}$ ,  $b = \frac{|B|}{|N_2|}$  and  $c = \frac{|C|}{|N_2|}$ , thus  $a + b + c = 1$ . For  $1 \leq k \leq 3$ , we say that  $T$  spans a  $k$ -joint-book, if among  $A, B, C$ , exactly  $3 - k$  of them are empty sets.

**Lemma 3.3.** *If  $T$  spans a 3-joint-book, then we have*

$$r_2n^2 \geq \frac{1}{6} \left[ \frac{3}{2} - \frac{21t}{4} \right]^2 n^2 - e(\overline{G'}) - (1 - 3t)n.$$

*Proof.* First notice that  $N_x, N_y$  and  $N_z$  are all independent sets. Indeed, suppose  $N_x$  contains an edge, then  $T \cup N_x \cup B$  contains two vertex-disjoint triangles, contradicting the maximality of  $\mathcal{T}$ .

Note that  $\binom{|N_x|}{2} + \binom{|N_y|}{2} + \binom{|N_z|}{2} \leq r_2n^2 + e(\overline{G'})$ . Indeed, every pair of vertices in  $N_x, N_y$  or  $N_z$  gives a non-edge in  $G'$  and those  $K_4$ -saturating edges in  $A, B, C$  are counted twice. Additionally,  $|N_x| + |N_y| + |N_z| = e(T, G') \geq \left(\frac{3}{2} - \frac{21t}{4}\right)n$ , and  $e(T, G') \leq 2(1 - 3t)n$ . Thus,

$$\begin{aligned} r_2n^2 + e(\overline{G'}) &\geq \binom{|N_x|}{2} + \binom{|N_y|}{2} + \binom{|N_z|}{2} \geq 3 \binom{e(T, G')/3}{2} \\ &= \frac{1}{6}(e(T, G'))^2 - \frac{1}{2}e(T, G') \geq \frac{n^2}{6} \left[ \frac{3}{2} - \frac{21t}{4} \right]^2 - (1 - 3t)n. \end{aligned}$$

$\square$

We first show that if  $T$  spans a 3-joint-book, then  $f(G) \geq 2n^2/33 - 3n/11$ .

**Lemma 3.4.** *For  $n \geq 73$ , if  $T$  spans a 3-joint-book, then  $f(G) \geq \frac{2n^2}{33} - \frac{3n}{11}$ .*

*Proof.* Note that  $e(G') + e(\overline{G'}) = \frac{(1-3t)^2n^2}{2} - \frac{(1-3t)n}{2}$ . By Lemmas 3.1 and 3.3, we have

$$\begin{aligned} f(G) &= (r_1 + r_2)n^2 \geq \left(\frac{1}{4} - t + \frac{3t^2}{2}\right)n^2 - e(G') - \frac{3}{2}tn \\ &\quad + \frac{1}{6} \left[ \frac{3}{2} - \frac{21t}{4} \right]^2 n^2 - e(\overline{G'}) - (1 - 3t)n \\ &\geq \left(\frac{51t^2}{32} - \frac{5t}{8} + \frac{1}{8}\right)n^2 - \frac{n}{2} \geq \frac{13n^2}{204} - \frac{n}{2} \geq \frac{2n^2}{33} - \frac{3n}{11}, \end{aligned}$$

since  $\frac{51t^2}{32} - \frac{5t}{8} + \frac{1}{8} \geq \frac{13}{204}$  when  $0 < t \leq 1/3$ , and the last inequality holds for  $n \geq 73$ .  $\square$

*Proof of Theorem 1.3.* By Lemma 3.4, we may assume that  $T$  spans a  $k$ -joint-book with  $k \leq 2$ . Without loss of generality assume that  $B = \emptyset$ , i.e.  $b = 0$ . Then  $a + c = 1$  and  $|A| + |C| = p_2 n$ . Notice that each pair of vertices in  $A$  and  $C$  is a  $K_4$ -saturating edge, hence

$$r_2 n^2 \geq \binom{|A|}{2} + \binom{|C|}{2} \geq 2 \binom{p_2 n / 2}{2} = \frac{p_2^2}{4} n^2 - \frac{p_2 n}{2}. \quad (1)$$

If  $t \geq \frac{1}{5}$ , then Lemma 3.1 implies  $f(G) \geq r_1 n^2 \geq \left(\frac{t}{2} - \frac{3t^2}{4}\right) n^2 - \frac{n}{2} \geq \frac{2n^2}{33}$  for  $n \geq 54$ . Thus we may assume that  $t < \frac{1}{5}$ . The right hand side in (1) is minimized when  $p_2$  is at its lower bound provided by Lemma 3.2, as  $\frac{1}{2} - \frac{9t}{4} > \frac{1}{n}$  for  $n \geq 20$ . Hence

$$r_2 n^2 \geq \frac{1}{4} \left(\frac{1}{2} - \frac{9t}{4}\right)^2 n^2 - \frac{1}{2} \left(\frac{1}{2} - \frac{9t}{4}\right) n.$$

Therefore using Lemma 3.1, we have

$$\begin{aligned} f(G) &= (r_1 + r_2) n^2 \geq \left( \left(\frac{t}{2} - \frac{3t^2}{4}\right) + \frac{1}{4} \left(\frac{1}{2} - \frac{9t}{4}\right)^2 \right) n^2 - \frac{1}{2} \left(3t + \frac{1}{2} - \frac{9t}{4}\right) n \\ &= \left(\frac{33t^2}{64} - \frac{t}{16} + \frac{1}{16}\right) n^2 - \frac{1}{2} \left(\frac{3t}{4} + \frac{1}{2}\right) n \geq \frac{2n^2}{33} - \frac{3n}{11} - \frac{3}{44}, \end{aligned}$$

where the function on the right hand side is minimized at  $t = \frac{2}{33} + \frac{4}{11n}$ . Since both  $tn$  and  $f(G)$  are integers, checking all  $n$  modulo 33, we have

$$f(G) \geq \frac{2n^2}{33} - \frac{3n}{11}.$$

$\square$

## References

- [1] N. Alon, An extremal problem for sets with applications to graph theory. *J. Combin. Theory, Ser. A*, 40, (1985), 82–89.
- [2] J. Balogh, B. Bollobás, R. Morris. Graph bootstrap percolation. *Random Structures & Algorithms*, 41, (2012), 413–440.
- [3] J. Balogh, B. Bollobás, R. Morris, O. Riordan. Linear algebra and bootstrap percolation. *J. Combin. Theory, Ser. A*, 119, (2012), 1328–1335.
- [4] B. Bollobás, Weakly  $k$ -saturated graphs. *Beiträge zur Graphentheorie*, (H. Sachs, H.J. Voss, and H. Walther, Eds.), pp. 25–31, Leipzig, Barth, 1968.

- [5] S. Das, W. Gan, B. Sudakov, Sperner's Theorem and a problem of Erdős-Katona-Kleitman. <http://arxiv.org/abs/1302.5210>.
- [6] P. Erdős, On a theorem of Rademacher-Turán. *Illinois Journal of Math*, 6, (1962), 122–127.
- [7] P. Erdős, On the number of complete subgraphs contained in certain graphs. *Magy. Tud. Acad. Mat. Kut. Int. Közl.*, 7, (1962), 459–474.
- [8] P. Erdős, Some of my old and new combinatorial problems. *Paths, flows, and VLSI-layout (Bonn, 1988)*, *Algorithms Combin.*, 9, 35–45, Springer, Berlin, 1990.
- [9] P. Frankl, An extremal problem for two families of sets. *Europ. J. Math.*, 3, (1982), 125–127.
- [10] G. Kalai, Weakly saturated graphs are rigid, in Convexity and Graph Theory, M. Rosenfeld and J. Zaks (eds). *Ann. Discrete Math.*, 20, (1984), 189–190.
- [11] N. Khadžiivanov, V. Nikiforov, Solution of a problem of P. Erdős about the maximum number of triangles with a common edge in a graph (Russian). *C. R. Acad. Bulgare Sci.*, 32, (1979), 1315–1318.
- [12] D. Kleitman, A conjecture of Erdős-Katona on commensurable pairs among subsets of an  $n$ -set. *Theory of Graphs, Proc. Colloq., Tihany*, (1966), 215–218.
- [13] L. Lovász, M. Simonovits, On the number of complete subgraphs of a graph, II. *Studies in pure mathematics*, 459–495, Birkhuser, Basel, 1983.
- [14] W. Mantel, Problem 28. *Wiskundige Opgaven*, 10, (1907), 60–61.
- [15] D. Mubayi, Counting substructures I: color critical graphs. *Advances in Mathematics*, 225, (2010), 2731–2740.
- [16] D. Mubayi, Counting substructures II: hypergraphs. *Combinatorica*, to appear.