

Name _____

(circle your TA discussion section)

- | | |
|---|---|
| ▷ AD1 , TR 9:00-10:50, Darlayne Addabbo | ▷ ADH , TR 3:00-3:50, Paulina Koutsaki |
| ▷ AD2 , TR 1:00-2:50, Ben Fulan | ▷ ADJ , TR 9:00-9:50, Jed Chou |
| ▷ ADA , TR 8:00-8:50, Chris Bailey | ▷ ADK , TR 10:00-10:50, Jed Chou |
| ▷ ADB , TR 9:00-9:50, Chris Bailey | ▷ ADL , TR 11:00-11:50, Andrew McConvey |
| ▷ ADC , TR 10:00-10:50, Andrew McConvey | ▷ ADM , TR 12:00-12:50, Benjamin Wright |
| ▷ ADD , TR 11:00-11:50, Diaa Taha | ▷ ADN , TR 1:00-1:50, Benjamin Wright |
| ▷ ADE , TR 12:00-12:50, Paul Spiegelhalter | ▷ ADO , TR 2:00-2:50, Paul Spiegelhalter |
| ▷ ADF , TR 1:00-1:50, Diaa Taha | ▷ ADP , TR 3:00-3:50, Wan-Yu Wu |
| ▷ ADG , TR 2:00-2:50, Paulina Koutsaki | ▷ ADQ , TR 4:00-4:50, Wan-Yu Wu |

- You may work with other MATH 220 students. However each student should write up solutions separately and independently – nobody should copy someone else’s work.
- You may use your notes, the textbook, or information found on my course home page.
- You may use a calculator only for basic arithmetic. In particular you should not use its graphing features.
- You are not allowed to search the Internet, use Wolfram Alpha, or use technology for anything beyond what is stated above.
- The quiz should be submitted to your TA at the beginning of your official discussion period on Tuesday, April 8th.
- There is a higher expectation for the quality of your work on a take-home quiz. Everything should be written logically and legibly with sufficient work to justify each answer. Blank copies of the quiz are available on the course home page.
- Be sure that the pages are nicely stapled – do not just fold the corners.
- **Note to TAs and Tutors – you should not help students with these specific problems or go over solutions until after 5pm Tuesday.**

1. (2 points) Find a formula for $f(x)$ given that $f''(x) = 2 \cos x + 5 \sin x$, $f(0) = 10$ and $f(\pi/2) = \pi + 7$.

2. (2 points) Suppose that $p(x)$ is continuous at all real numbers and satisfies the following equations.

- $\int_2^5 p(x) dx = 4$
- $\int_2^{10} p(x) dx = 13$
- $\int_{10}^{25} p(x) dx = 61$
- $\int_{20}^{25} p(x) dx = 36$

What is the value of $\int_5^{20} (3p(x) - 4) dx$?

3. (2 points) Evaluate the following limit. Use proper notation throughout your evaluation of this limit.

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(3k + 2n)^2}{n^3}$$

4. (2 points) From section 5.2 we have the following property of definite integrals.

- If $f(x)$ is continuous and $m \leq f(x) \leq M$ for $a \leq x \leq b$, then $m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$

Use this property to carefully explain why the following inequality holds.

$$0.6 \leq \int_{-1}^2 \frac{1}{\sqrt{17+x^3}} dx \leq 0.75$$

5. (2 points) The area between the x -axis and the graph of $f(x) = \frac{1}{x^3 + 2}$ on the interval $[6, 11]$ can be written as a limit of Riemann sums in many different ways. I have shown how to do this for two of the six ways indicated below. Fill in the missing information for the remaining limits so that the only variables appearing are n and k . Do not evaluate these limits.

- (a) Using a limit of right Riemann sums,

$$AREA = \lim_{n \rightarrow \infty} \sum_{k=1}^n \left[\frac{1}{\left(6 + k \cdot \frac{5}{n}\right)^3 + 2} \cdot \frac{5}{n} \right]$$

- (b) Using a limit of right Riemann sums,

$$AREA = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} \left[\qquad \qquad \qquad \right]$$

- (c) Using a limit of left Riemann sums,

$$AREA = \lim_{n \rightarrow \infty} \sum_{k=1}^n \left[\qquad \qquad \qquad \right]$$

- (d) Using a limit of left Riemann sums,

$$AREA = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} \left[\qquad \qquad \qquad \right]$$

- (e) Using a limit of midpoint Riemann sums,

$$AREA = \lim_{n \rightarrow \infty} \sum_{k=1}^n \left[\qquad \qquad \qquad \right]$$

- (f) Using a limit of midpoint Riemann sums,

$$AREA = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} \left[\frac{1}{\left(6 + (k + 0.5) \cdot \frac{5}{n}\right)^3 + 2} \cdot \frac{5}{n} \right]$$