

CLASSIFYING SPACES FOR 1-TRUNCATED COMPACT LIE GROUPS

CHARLES REZK

ABSTRACT. A 1-truncated compact Lie group is any extension of a finite group by a torus. In this note we compute the homotopy types of $\text{Map}(BG, BH)$ and $(B_G H)^G$ for compact Lie groups G and H with H 1-truncated, showing that they are computed entirely in terms of spaces of homomorphisms from G to H . These results generalize the well-known case when H is finite, and the case of H compact abelian due to Lashof, May, and Segal.

1. INTRODUCTION

By a **1-truncated** compact Lie group H , we mean one whose homotopy groups vanish in dimensions 2 and greater. Equivalently, H is a compact Lie group with identity component H_0 a torus (isomorphic to some $U(1)^d$); i.e., an extension of a finite group by a torus.

The class of 1-truncated compact Lie groups includes (i) all finite groups, and (ii) all compact abelian Lie groups, both of which are included in the class (iii) all groups which are isomorphic to a product of a compact abelian Lie group with a finite group, or equivalently a product of a torus with a finite group.

The goal of this paper is to extend certain results, which were already known for finite groups, compact abelian Lie groups, or products thereof, to all 1-truncated compact Lie groups.

We write $\text{Hom}(G, H)$ for the space of continuous homomorphisms, equipped with the compact-open topology. Our first theorem relates this to the space of based maps between classifying spaces.

1.1. Theorem. *For G, H compact Lie groups with H 1-truncated, the evident map*

$$B: \text{Hom}(G, H) \rightarrow \text{Map}_*(BG, BH)$$

is a weak equivalence.

Using this, we will derive an unbased variant.

1.2. Theorem. *For G, H compact Lie groups with H 1-truncated, there is a weak equivalence*

$$\text{Hom}(G, H) \times_H EH \rightarrow \text{Map}(BG, BH).$$

Here H acts on $\text{Hom}(G, H)$ by conjugation: $h \cdot \phi = h\phi h^{-1}$.

When H is discrete, these are well-known and classical results. The case of H an abelian compact Lie group is proved in [LMS83]; both the finite and compact abelian Lie cases are discussed in [May90].

1.3. Remark. For G and H compact, there is a homeomorphism (7.1)

$$\text{Hom}(G, H) \approx \coprod_{[\phi: G \rightarrow H]} H/C_H(\phi)$$

Date: February 2, 2017.

The author was supported under NSF grant DMS-1406121.

where the coproduct is over conjugacy classes of homomorphisms, and $C_H(\phi)$ is the centralizer of $\phi(G)$ in H . When H is a 1-truncated compact Lie group, we see from (1.1) that $\text{Map}_*(BG, BH)$ is also weakly equivalent to this coproduct, and from (1.2) that there is a weak equivalence

$$\text{Map}(BG, BH) \approx \coprod_{[\phi: G \rightarrow H]} BC_H(\phi).$$

Finally, we will give a description of the fixed points of the equivariant classifying space $B_G H$, which represents G -equivariant H -principal bundles, in the case that G and H are compact Lie and H is 1-truncated.

1.4. Theorem. *For G, H compact Lie groups with H 1-truncated, the map*

$$\pi^*: B_G H \rightarrow \text{Map}(EG, B_G H)$$

*induced by restriction along $\pi: EG \rightarrow *$ is a G -equivariant weak equivalence.*

The case of H finite or compact abelian is proved in [May90].

1.5. Remark. For any closed subgroup $G' \leq G$, taking G' fixed points gives rise to a map

$$(B_G H)^{G'} \rightarrow \text{Map}(EG, B_G H)^{G'} \xleftarrow{\sim} \text{Map}(EG, BH)^{G'} \approx \text{Map}(B_{G'} H, BH),$$

and (1.4) amounts to saying that for any G' this map is a weak equivalence. It is standard [LM86, Theorem 10] that, for arbitrary compact G and H , $(B_G H)^{G'}$ is weakly equivalent to $\coprod_{[\phi: G' \rightarrow H]} BC_H(\phi)$, while if H is also 1-truncated, (1.2) and (1.3) imply that $\text{Map}(B_{G'} H, BH)$ is also weakly equivalent to the same coproduct, thus giving an abstract weak equivalence $(B_G H)^{G'} \approx \text{Map}(B_{G'} H, BH)$. The point of (1.4) is to show that the map π^* exhibits this equivalence.

The map of (1.4) in a certain sense classifies the formation of the G -Borel quotient. That is, given a G -equivariant map $f: X \rightarrow B_G H$ classifying a G -equivariant principal H -bundle $P \rightarrow X$, the G -equivariant map $\pi^* f: X \rightarrow \text{Map}(EG, B_G H)$ is adjoint to a non-equivariant map $X \times_G EG \rightarrow B_G H$ which classifies the bundle $P \times_G EG \rightarrow X \times_G EG$; see [May90]. As a consequence of the theories of classifying spaces, we obtain the following.

1.6. Corollary. *Let G and H be compact Lie groups with H 1-truncated. Then for a paracompact G -space X , formation of G -Borel quotient gives rise to a bijection between (i) equivalence classes of G -equivariant principal H -bundles over X , and (ii) equivalence classes of principal H -bundles over $X \times_G EG$.*

1.7. Organization of the paper. The proof of (1.1) is the probably the most interesting part of the paper. It is carried out in §§2–4. The key ingredient is the use of the nerve $N(H, V)$ of the “exponential crossed module” (2.3) of the 1-truncated compact Lie group H . We first show that the simplicial space $N(H, V)$ is a Reedy fibrant model for the usual simplicial nerve NH of H (3.8), and so can be used to compute maps BG to BH , in terms of maps of simplicial spaces from NG to $N(H, V)$. The proof is completed (§4) by showing that, in a certain sense, the difference between $\text{Hom}(G, H)$ and the space $\text{Map}_{s\text{Top}}(NG, N(H, V))$ of maps between simplicial spaces is measured precisely by the continuous 2-cocycles on G with values in V , modulo boundaries of 1-cocycles. Because G is compact, Haar measure gives a contracting homotopy (4.2) on the complex of continuous chains on G . A sketch by the author of this proof originally appeared as an answer to a question on the site MathOverflow¹.

Our approach gives a uniform proof of (1.1) for all 1-truncated compact Lie groups H ; furthermore, even in the case of abelian H , it is somewhat more direct than the one given in [LMS83].

We derived the unbased theorem (1.2) from the based version (1.1) in §5, by comparing associated fibrations over BH .

¹“Equivariant classifying spaces from classifying spaces”, <http://mathoverflow.net/q/223546>.

The result on equivariant classifying spaces (1.4) is proved in §6. The proof relies on an explicit model, built as the nerve of a certain topological category, of the restriction of the universal (G, H) -bundle to the fixed point subspace $(B_G H)^G \subseteq B_G H$. The explicit model we use appears to be essentially of the type described in [GMM12].

In §7, we give for the convenience of the reader a proof of the identification of $\text{Hom}(G, H)$ as mentioned in (1.3).

1.8. Acknowledgments. I thank Peter May for comments on a draft of this paper.

1.9. Conventions. In this paper, we write Top for the category of *compactly generated weak Hausdorff spaces* ($CGWH$), the standard convenient category of spaces. This category is cartesian closed, and we write $\text{Map}(X, Y)$ for the internal function object, i.e., continuous maps with the k -ification of the compact-open topology. We make use of the “usual” model structure on Top , in which weak equivalences are weak equivalences on homotopy groups, and fibrations are Serre fibrations.

2. NERVE OF A TOPOLOGICAL CROSSED MODULE

2.1. Crossed modules. Recall that a **crossed module** consists of

- groups H and V ,
- a homomorphism $\epsilon: V \rightarrow H$,
- a homomorphism $\alpha: H \rightarrow \text{Aut}(V)$,
- such that (i) $\epsilon(\alpha(h)(v)) = h\epsilon(v)h^{-1}$ and (ii) $\alpha(\epsilon(v))(v') = vv'v^{-1}$ for $h \in H, v, v' \in V$.

A **topological crossed module** is one in which V and H are topological groups, and ϵ and α are continuous. I’ll typically write (H, V) for the crossed module, leaving ϵ and α understood. Note that we will often consider crossed modules in which V is an abelian group, in which case we will switch to additive notation for V , though not for H .

2.2. Example. Given any group H , there is a unique crossed module which we will denote $(H, 0)$, in which 0 is the trivial group.

2.3. Example (Exponential crossed module). The following example is the crucial one for this paper. Suppose H is a 1-truncated compact Lie group. We set

- $V := T_e H$, the Lie algebra of H , which is a group under addition of vectors;
- $\epsilon := \exp: V \rightarrow H$, the exponential map; this is a homomorphism since H_0 is abelian;
- $\alpha := \text{ad}: H \rightarrow GL(V)$, the adjoint action.

We typically write the group law of V additively, so the identities for the crossed module structure become

$$\exp(\text{ad}(h)(v)) = h \exp(v) h^{-1}, \quad \text{ad}(\exp(v))(v') = v + v' - v = v'.$$

The following features of this case will be significant:

- (1) $\alpha = \text{ad}: H \rightarrow GL(V)$ factors through the quotient group H/H_0 ,
- (2) $\epsilon = \exp: V \rightarrow H$ is a covering map,
- (3) the underlying space of V is contractible.

2.4. Nerve of a crossed module. The **nerve** of a topological crossed module $N(H, V)$ is the simplicial space defined as follows; except for the topology, this is as in [Bro99, §3.1]. The space $N(H, V)_n$ in degree n is the space of tuples

$$((h_{ij})_{0 \leq i \leq j \leq n}, (v_{ijk})_{0 \leq i \leq j \leq k \leq n}), \quad h_{ij} \in H, \quad v_{ijk} \in V,$$

satisfying the identities

- (1) $h_{ii} = e$ and $v_{iij} = v_{ijj} = e$ for all $i \leq j$,
- (2) $h_{ik} = \epsilon(v_{ijk})h_{ij}h_{jk}$ for all $i \leq j \leq k$,

(3) $v_{ik\ell}v_{ijk} = v_{ij\ell} \alpha(h_{ij})(v_{jkl})$ for all $i \leq j \leq k \leq \ell$.

The action of simplicial operators $\delta: [n] \rightarrow [m]$ is the evident one: $(\delta h)_{ij} = h_{\delta(i), \delta(j)}$ and $(\delta v)_{ijk} = v_{\delta(i), \delta(j), \delta(k)}$. A standard argument shows that as a space $N(H, V)_n \approx H^n \times V^{\binom{n}{2}}$, e.g., via the projection to coordinates h_{0i} , $1 \leq i \leq n$ and v_{0ij} , $1 \leq i < j \leq n$.

Note that $N(H, V)_0 = *$, i.e., $N(H, V)$ is a *reduced* simplicial space.

2.5. *Example.* The nerve of $N(H, 0)$ is precisely the usual nerve of the group H ; we write $N(H) := N(H, 0)$.

2.6. **Simplicial spaces and the Reedy model structure.** We write $s\text{Top}$ for the category of simplicial spaces, i.e., functors $\Delta^{\text{op}} \rightarrow \text{Top}$. We are going to use the **Reedy model structure** on $s\text{Top}$. We will need to use the following features of this model structure:

- (1) Weak equivalences $f: X \rightarrow Y$ in $s\text{Top}$ are precisely the levelwise weak equivalences, i.e., $f_n: X_n \rightarrow Y_n$ is a weak equivalence for all $n \geq 0$.
- (2) An object X is cofibrant (**Reedy cofibrant**) if and only if the latching space inclusions $\gamma_n: L_n X \rightarrow X_n$ are cofibrations in Top .
- (3) An object Y is fibrant (**Reedy fibrant**) if and only if the matching space projections $\delta_n: Y_n \rightarrow M_n Y$ are fibrations in Top .
- (4) The model structure is topological. In particular, if X is a cofibrant simplicial space and $Y \rightarrow Y'$ is a weak equivalence between fibrant simplicial spaces, then $\text{Map}_{s\text{Top}}(X, Y) \rightarrow \text{Map}_{s\text{Top}}(X, Y')$ is a weak equivalence of spaces.

We will need to examine latching and matching spaces in a bit more detail.

2.7. **Latching and matching spaces.** We recall the notion of latching and matching spaces. For simplicial spaces $X: \Delta^{\text{op}} \rightarrow \text{Top}$ and all $n \geq 0$, we have natural maps of spaces

$$L_n X \xrightarrow{\gamma_n} X_n \xrightarrow{\delta_n} M_n X$$

where

$$L_n X = \text{colim}_{(\Delta_{[n]}^{\text{op}})_{<n}} X, \quad M_n = \text{lim}_{(\Delta_{[n]'}^{\text{op}})_{<n}} X,$$

called the **latching** and **matching** spaces of X .

2.8. **Latching spaces for the nerve of a group.**

2.9. **Proposition.** *Let G be a topological group, and $NG \in s\text{Top}$ its nerve. Then for each $n \geq 0$, the latching inclusion $\gamma_n: L_n(NG) \rightarrow (NG)_n$ is isomorphic to the inclusion*

$$\{(g_1, \dots, g_n) \mid \exists i, g_i = e\} \rightarrow G^n.$$

In particular, NG is Reedy cofibrant if $\{e\} \rightarrow G$ is a cofibration in Top .

Proof. Standard. □

2.10. **Matching spaces for the nerve of a crossed module.** We describe the matching projections for the nerve of a topological crossed module.

2.11. **Proposition.** *Consider $N := N(H, V)$ the nerve of a topological crossed module. We write $M_n := M_n N$ for its matching spaces.*

- (0) $\delta_0: N_0 \rightarrow M_0$ is the isomorphism of 1-point spaces.
- (1) $\delta_1: N_1 \rightarrow M_1$ is the projection $H \rightarrow *$.

(2) $M_2 \approx H^{\times 3}$, and there is a pullback square

$$\begin{array}{ccc} N_2 & \longrightarrow & V \\ \delta_2 \downarrow & & \downarrow \epsilon \\ M_2 & \xrightarrow{(h_{01}, h_{02}, h_{12}) \mapsto h_{02} h_{12}^{-1} h_{02}^{-1}} & H \end{array}$$

(3) There is a commutative diagram

$$\begin{array}{ccccc} N_3 & \longrightarrow & \{e\} & & \\ \delta_3 \downarrow & & \downarrow & & \\ M_3 & \longrightarrow & \text{Ker } \epsilon & \longrightarrow & \{e\} \\ \downarrow & & \downarrow & & \downarrow \\ H^{\times 3} \times V^{\times 3} & \xrightarrow{\pi} & V & \xrightarrow{\epsilon} & H \end{array}$$

in which all squares are pullback squares, and π is given by

$$(h_{01}, h_{12}, h_{23}, v_{012}, v_{013}, v_{023}, v_{123}) \mapsto v_{023} v_{012} \alpha(h_{01}) (v_{123})^{-1} v_{013}^{-1}.$$

(≥ 4) $\delta_n: N_n \rightarrow M_n$ is an isomorphism for $n \geq 4$.

Proof. Straightforward. In (3), one shows directly that the right-hand lower square, bottom rectangle, and left rectangle are pullbacks. \square

Recall that a simplicial space X is **Reedy fibrant** if each of the maps $\delta_n: X_n \rightarrow M_n X$ is a fibration of spaces.

2.12. Corollary. *If (H, V) is a topological crossed module such that ϵ is a covering map, then $N(H, V)$ is Reedy fibrant.*

Proof. Immediate using (2.11). Note that the condition that ϵ be a covering map in (2.11)(3) implies that δ_3 is an open and closed embedding. \square

In particular, (2.12) applies to our main example (2.3).

3. MAPS BETWEEN REDUCED SIMPLICIAL SPACES

A simplicial space $X \in s\text{Top}$ is said to be **reduced** if $X_0 \approx *$. We write $s\text{Top}^{\text{red}} \subset s\text{Top}$ for the full subcategory of reduced simplicial spaces. Note that reduced simplicial spaces are canonically based, so that we may in fact regard $s\text{Top}^{\text{red}}$ as a full subcategory of simplicial based spaces $s\text{Top}_*$.

3.1. Realization of reduced simplicial spaces. We recall the geometric realization functor $\|-\|: s\text{Top} \rightarrow \text{Top}$, defined so that $\|X\|$ is the coend of the functor $\Delta^{\text{op}} \times \Delta \rightarrow \text{Top}$ given by $([m], [n]) \mapsto X_m \times \Delta^n$, where Δ^n is the topological n -simplex.

3.2. Proposition. *The restriction of the geometric realization functor to a functor $\|-\|: s\text{Top}^{\text{red}} \rightarrow \text{Top}_*$ admits a right adjoint $\nabla: \text{Top}_* \rightarrow s\text{Top}^{\text{red}}$, defined by*

$$(\nabla Y)_n := \text{Map}_*(\Delta^n / \text{Sk}_0 \Delta^n, Y),$$

where $\text{Sk}_0 \Delta^n \subseteq \Delta^n$ is the set of vertices of the simplex. The adjunction is compatible with the topological enrichment, and so gives a natural homeomorphism

$$\text{Map}_{s\text{Top}}(X, \nabla Y) \approx \text{Map}_*(\|X\|, Y)$$

for $X \in s\text{Top}^{\text{red}}$ and $Y \in \text{Top}_*$.

Proof. This is a straightforward consequence of the observation that for reduced simplicial spaces X , $\|X\|$ is seen to be isomorphic to the coend (in Top_*) of $([m], [n]) \mapsto X_m \wedge (\Delta^n / \text{Sk}_0 \Delta^n)$. \square

We also note the following.

3.3. Proposition. *For any $Y \in \text{Top}_*$, the simplicial space ∇Y is Reedy fibrant.*

Proof. The matching space projection has the form $\text{Map}_*(\Delta^n / \text{Sk}_0 \Delta^n, Y) \rightarrow \text{Map}_*(\partial \Delta^n / \text{Sk}_0 \Delta^n, Y)$, which is clearly a fibration. \square

For a topological group H , we consider the classifying space $BH := \|NH\|$.

3.4. Proposition. *If H is a topological group with identity element a non-degenerate basepoint (i.e., $\{e\} \rightarrow H$ has the HEP), then the map*

$$\eta: NH \rightarrow \nabla \|NH\| = \nabla BH$$

given by the unit map of the adjunction of (3.2) is a levelwise weak equivalence of simplicial spaces.

Proof. In degree 0, η is the isomorphism of one-point spaces. In degree 1 it has the form

$$H \rightarrow \text{Map}_*(\Delta^1 / \{0, 1\}, \|NH\|) \approx \Omega BH.$$

A standard argument (e.g., using the usual simplicial model for the universal fibration [May75]) shows that this is a weak equivalence.

For $n \geq 2$, we reduce to the $n = 1$ case using the fact that $I_n / \text{Sk}_0 \Delta^n \rightarrow \Delta^n / \text{Sk}_0 \Delta^n$ is a homotopy equivalence of pointed spaces, and thus

$$\text{Map}_*(\Delta^n / \text{Sk}_0 \Delta^n, \|NH\|) \rightarrow \text{Map}_*(I_n / \text{Sk}_0 \Delta^n, \|NH\|) \approx (\Omega BG)^{\times n}$$

is a weak equivalence, where $I_n \subseteq \Delta^n$ is the union of the edges with vertices $\{k-1, k\}$ for all $k = 1, \dots, n$. \square

3.5. $\text{Map}(X, N(H, V))$ computes $\text{Map}_*(\|X\|, BH)$. Now we fix a 1-truncated compact Lie group H and the corresponding exponential crossed module (H, V) of (2.3). We have a map of reduced simplicial spaces

$$NH \xrightarrow{(\iota, \eta)} N(H, V) \times \nabla \|NH\|$$

in which ι is the evident inclusion $NH = N(H, 0) \subseteq N(H, V)$, and η the unit map of the adjunction of (3.2). Observe that both ι and η are levelwise weak equivalences (ι because V is contractible, η by (3.4)). Furthermore, both $N(H, V)$ (2.12) and $\nabla \|NH\|$ (3.3) are Reedy fibrant.

Using the Reedy model structure on simplicial spaces, we can factor the above map as

$$(3.6) \quad NH \xrightarrow{j} (NH)^f \xrightarrow{(\iota', \eta')} N(H, V) \times \nabla \|NH\|$$

so that $(NH)^f$ is Reedy fibrant and j is a levelwise weak equivalence, whence ι' and η' are also levelwise weak equivalences.

3.7. Proposition. *For X a Reedy cofibrant simplicial space with $X_0 = *$, and (H, V) the exponential crossed module of a 1-truncated compact Lie group H , we have that $\text{Map}_{s\text{Top}}(X, N(H, V))$ is weakly equivalent to $\text{Map}_*(\|X\|, \nabla \|NH\|)$. Furthermore, $\iota_*: \text{Map}_{s\text{Top}}(X, NH) \rightarrow \text{Map}_{s\text{Top}}(X, N(H, V))$ is a weak equivalence of spaces if and only if $\eta_*: \text{Map}_{s\text{Top}}(X, NH) \rightarrow \text{Map}_{s\text{Top}}(X, \nabla \|NH\|)$ is.*

Proof. Straightforward using the factorization (3.6), the fact that Reedy model structure is compatible with the topological enrichment, and the adjunction (3.2). \square

3.8. Corollary. *If (H, V) is as above, and G is a topological group such that $\{e\} \rightarrow G$ is a cofibration, then $\text{Map}_*(BG, BH)$ is weakly equivalent to $\text{Map}_{s\text{Top}}(NG, N(H, V))$, and*

$$B: \text{Hom}(G, H) \rightarrow \text{Map}_*(BG, BH)$$

is a weak equivalence if and only if

$$\iota_*: \text{Map}_{s\text{Top}}(NG, NH) \rightarrow \text{Map}_{s\text{Top}}(NG, N(H, V))$$

is a weak equivalence.

Proof. Use (3.7) with $X = NG$, which is Reedy cofibrant by (2.9). It is straightforward to see that $\text{Map}_{s\text{Top}}(NG, NH) \rightarrow \text{Hom}(G, H)$ (evaluation at spaces in degree 1) is a homeomorphism, and so the map B coincides with ι_* . \square

3.9. Remark. If H is a discrete group, then NH is already Reedy fibrant, in which case we can immediately derive the well-known fact that $B: \text{Hom}(G, H) \rightarrow \text{Map}_*(BG, BH)$ is a weak equivalence for any such topological group G .

4. PROOF OF (1.1): BASED MAPPING SPACE

As above, we assume that H is a 1-truncated compact Lie group. We will now also assume that G is a compact Lie group. By (3.8), we have reduced (1.1) to showing that $\text{Map}_{s\text{Top}}(NG, NH) \rightarrow \text{Map}_{s\text{Top}}(NG, N(H, V))$ is a weak equivalence.

Let $E := \text{Map}_{s\text{Top}}(NG, N(H, V))$. Using (2.11) and the identification of the latching inclusions $L_n NG \rightarrow G^n$ (2.9), we see that E is precisely the space of pairs

$$(\zeta, \nu) \in \text{Map}(G, H) \times \text{Map}(G \times G, V)$$

such that

- (1) $\zeta(e) = e$ and $\nu(g, e) = 0 = \nu(e, g)$ for $g \in G$,
- (2) $\zeta(g_1 g_2) = \exp[\nu(g_1, g_2)]\zeta(g_1)\zeta(g_2)$ for $g_1, g_2 \in G$,
- (3) $\nu(g_1 g_2, g_3) + \nu(g_1, g_2) = \nu(g_1, g_2 g_3) + \text{ad}(\zeta(g_1))[\nu(g_2, g_3)]$ for $g_1, g_2, g_3 \in G$.

Explicitly, this corresponds to the map $NG \rightarrow N(H, V)$ which (in the notation of §2.1) sends $(g_{ij}) \in (NG)_n$ to $(h_{ij}, v_{ijk}) \in N(H, V)_n$ with $h_{ij} = \zeta(g_{ij})$ and $v_{ijk} = \nu(g_{ij}, g_{jk})$.

Let $E^0 := \text{Map}_{s\text{Top}}(NG, NH)$. The map $E^0 \rightarrow E$ is precisely inclusion into the subspace consisting of points of the form $(\zeta, 0)$.

For a continuous map $\zeta: G \rightarrow H$, we write $\bar{\zeta}: G \rightarrow H/H_0$ for the composite with the quotient map $H \rightarrow H/H_0$. Note that if $(\zeta, \nu) \in E$, then $\bar{\zeta}$ is a continuous homomorphism of groups. Since H/H_0 is discrete, we obtain coproduct decompositions

$$E = \coprod_{\gamma} E_{\gamma}, \quad E^0 = \coprod_{\gamma} E_{\gamma}^0, \quad \gamma \in \text{Hom}(G, H/H_0).$$

Thus, we must show that for each such γ , the inclusion $E_{\gamma}^0 \subseteq E_{\gamma}$ is a weak equivalence. In fact, we can give an explicit (strong) deformation retraction of E_{γ} to E_{γ}^0 , which relies on the existence of a contracting homotopy of the complex $C^*(G, V_{\text{ad } \gamma})$ of normalized continuous cochains on G with values in the representation $\text{ad } \gamma: G \rightarrow \text{Aut}(V)$, which may be constructed explicitly using an invariant measure on the compact group G . We spell out the details we need below.

Fix $\gamma \in \text{Hom}(G, H/H_0)$. Let $C_{\gamma}^1 \subseteq \text{Map}(G, V)$ be the subspace of functions $\mu: G \rightarrow V$ such that

$$\mu(e) = 0.$$

Let $Z_{\gamma}^2 \subseteq \text{Map}(G \times G, V)$ be the subspace of functions $\nu: G \times G \rightarrow V$ such that

$$\nu(g, e) = 0 = \nu(e, g), \quad g \in G,$$

and

$$\nu(g_1 g_2, g_3) + \nu(g_1, g_2) = \nu(g_1, g_2 g_3) + \text{ad } \gamma(g_1)[\nu(g_2, g_3)], \quad g_1, g_2, g_3 \in G.$$

Both Z_γ^2 and C_γ^1 are topological real vector spaces. Define continuous and linear maps

$$d: C_\gamma^1 \rightarrow Z_\gamma^2, \quad H: Z_\gamma^2 \rightarrow C_\gamma^1$$

by

$$\begin{aligned} d\mu(g_1, g_2) &:= \mu(g_1) - \mu(g_1 g_2) + \text{ad } \gamma(g_1)\mu(g_2), \\ H\nu(g) &:= \int_G x^{-1}\nu(x, g) dx, \end{aligned}$$

where we use right-invariant Haar measure on G normalized so that $\int_G dx = 1$.

4.1. Lemma. *The composite $dH: Z_\gamma^2 \rightarrow Z_\gamma^2$ is the identity map.*

Proof. For $g \in G$ and $v \in V$ we write “ gv ” for $\text{ad } \gamma(g)(v)$ below. Given $\nu \in Z_\gamma^2$ we have

$$\begin{aligned} dH\nu(g_1, g_2) &= \int_G x^{-1}\nu(x, g_1) - x^{-1}\nu(x, g_1 g_2) + g_1 x^{-1}\nu(x, g_2) dx \\ &= \int_G x^{-1}\nu(x, g_1) - x^{-1}[\nu(xg_1, g_2) + \nu(x, g_1) - x\nu(g_1, g_2)] + g_1 x^{-1}\nu(x, g_2) dx \\ &= \nu(g_1, g_2) - \int_G g_1(xg_1)^{-1}\nu(xg_1, g_2) dx + \int_G g_1 x^{-1}\nu(x, g_2) dy = \nu(g_1, g_2), \end{aligned}$$

the last cancellation by right-invariance of the measure. \square

4.2. Proposition. *The inclusion $E_\gamma^0 \subseteq E_\gamma$ admits a strong deformation retraction.*

Proof. Define $K_t: E_\gamma \rightarrow E_\gamma$ for $0 \leq t \leq 1$ by $K_t(\zeta, \nu) := (\zeta_t, \nu_t)$, with

$$\begin{aligned} \zeta_t(g) &:= \exp[t H\nu(g)]\zeta(g), \\ \nu_t(g_1, g_2) &:= \nu(g_1, g_2) - t dH\nu(g_1, g_2). \end{aligned}$$

We have $K_0 = \text{id}_{E_\gamma}$, $K_t|_{E_\gamma^0} = \text{id}_{E_\gamma^0}$, and $K_1(E_\gamma) \subseteq E_\gamma^0$ as desired, the last using (4.1). \square

The proof of (1.1) follows, using (3.7) and the remarks above.

4.3. Remark. If H is an abelian group, then $\text{ad}: H \rightarrow \text{Aut}(V)$ is trivial. In this case, the proof of (4.2) directly gives a deformation retraction of $E^0 \subseteq E$.

5. PROOF OF (1.4): UNBASED MAPPING SPACE

Given simplicial spaces X and Y , we have an internal function object $\underline{\text{Map}}(X, Y) \in s\text{Top}$, characterized so that $\underline{\text{Map}}(X, -)$ is the right adjoint to $(-) \times X$. We have that

$$\underline{\text{Map}}(X, Y)_n = \text{Map}_{s\text{Top}}(X \times N[n], Y),$$

where $[n]$ is the n -arrow category. In particular, $\underline{\text{Map}}(X, Y)_0 \approx \text{Map}_{s\text{Top}}(X, Y)$.

Formation of the internal function object is compatible with realization: there are canonical maps

$$(5.1) \quad \rho: \|\underline{\text{Map}}(X, Y)\| \rightarrow \text{Map}(\|X\|, \|Y\|)$$

natural in X and Y . This map exists exactly because the realization functor $\|- \| : s\text{Top} \rightarrow \text{Top}$ preserves finite products, and is characterized as the map adjoint to

$$\|\underline{\text{Map}}(X, Y)\| \times \|X\| \xleftarrow{\sim} \|\underline{\text{Map}}(X, Y) \times X\| \xrightarrow{|\text{eval}|} \|Y\|.$$

Given topological groups G and H , we consider the function object $\underline{\text{Map}}(NG, NH)$. We have an evident isomorphism

$$\underline{\text{Map}}(NG, NH) \approx N\text{Fun}(G, H),$$

where $\text{Fun}(G, H)$ is the internal category in Top of functors and natural transformations from G to H . Explicitly, this has

- objects $\phi \in \text{Hom}(G, H)$, and
- morphisms $\phi_0 \xrightarrow{h} \phi_1$ where $h \in H$, $\phi_1 = h\phi_0h^{-1}$.

and thus homeomorphisms $N\text{Fun}(G, H)_n = \underline{\text{Map}}(NG, NH)_n = \text{Hom}(G, H) \times H^{\times n}$.

Write $(H \curvearrowright H)$ for the translation category of the left action of H on itself, viewed as a category object in Top . This has

- objects $h_0 \in H$,
- morphisms $h_0 \xrightarrow{h} h_1$ where $h \in H$, $h_1 = hh_0$.

We have homeomorphisms $N(H \curvearrowright H)_n = H^{\times(n+1)}$. The group H acts on the category $(H \curvearrowright H)$ by $\delta \cdot h_0 = h_0\delta^{-1}$ (on objects) and $\delta \cdot (h_0 \xrightarrow{h} h_1) = h_0\delta^{-1} \xrightarrow{h} h_1\delta^{-1}$ (on morphisms), where $\delta \in H$.

We let $EH := \|N(H \curvearrowright H)\|$, a contractible H space with free H -action.

5.2. Lemma. *There is a homeomorphism $\|\underline{\text{Map}}(NG, NH)\| \approx (\text{Hom}(G, H) \times EH)/H$, where H acts on $\text{Hom}(G, H)$ by conjugation.*

Proof of (1.2). We have a commutative diagram

$$\begin{array}{ccc} \|\underline{\text{Map}}(NG, NH)\| & \xrightarrow{\rho} & \text{Map}(\|NG\|, \|NH\|) \\ \alpha \downarrow & & \downarrow \beta \\ \|\underline{\text{Map}}(*, NH)\| & \xrightarrow{\approx} & \text{Map}(\|*\|, \|NH\|) \end{array}$$

where the vertical maps are induced by restriction along $* \rightarrow NG$, and the lower horizontal map is the evident homeomorphism (both source and target are homeomorphism BH). We claim that ρ is a weak equivalence.

By (5.2) we see that $\alpha: (\text{Hom}(G, H) \times EH)/H \rightarrow BH$ is a fiber bundle with fiber H . Since $\beta: \text{Map}(BG, BH) \rightarrow BH$ is also a fibration, and the base space BH is path connected, ρ is a weak equivalence if and only if its restriction to the fiber over the base point is, which is precisely the weak equivalence $\text{Hom}(G, H) \rightarrow \text{Map}_*(BG, BH)$ of (1.1). \square

6. PROOF OF (1.4): EQUIVARIANT CLASSIFYING SPACE

6.1. Recollections on equivariant bundles. A G -equivariant principal H bundle (or (G, H) -bundle), is a principal H -bundle $\pi: P \rightarrow X$, together with actions of G on P and X , compatible with π , so that G acts via maps of principal H -bundles. We will always assume that both G and H are compact Lie groups.

This definition is somewhat anomalous, in that (G, H) -bundles are not characterized by a property which is local in X . Thus, we say that a (G, H) -bundle is **locally trivial** if it looks locally like

$$(G \times H) \times_{\Lambda} U \rightarrow G \times_{G'} U,$$

where $G' \leq G$ is a closed subgroup, $\Lambda_{\phi} := \{(g, \phi(g)) \mid g \in G'\}$ is the graph of some homomorphism $\phi: G' \rightarrow H$, and $\Lambda \xrightarrow{\sim} G'$ acts on a space U . The key result is that if G and H are compact and X is completely regular², then any (G, H) -bundle over X is locally trivial [Las82, Cor. 1.5].

A (G, H) -bundle $P \rightarrow X$ is **numerable** if it admits a locally trivializing cover which itself admits a subordinate partition of unity by G -invariant functions. Over a paracompact base X , every locally trivial bundle is numerable [Las82, Cor. 1.13]. There is a universal (G, H) -bundle $E_G H \rightarrow B_G H$, which classifies equivalence classes of numerable bundles: see [Las82], [LM86], and also [LU14] for a recent and more general treatment.

²Completely regular = points are closed, and any point and disjoint closed subset are separated by a real valued function.

We will be mainly concerned with the case of (G, H) -bundles $\pi: P \rightarrow X$ such that G acts trivially on X . In such a case there is a natural function

$$\tau: P \rightarrow \text{Hom}(G, H)$$

defined so that $\tau(p)(\gamma) \in H$ is the unique $\delta \in H$ such that $(\gamma, \delta) \cdot p = p$. When $P \rightarrow X$ is locally trivial, the map τ is seen to be continuous. Observe that τ is $G \times H$ -equivariant, where this group acts on $\text{Hom}(G, H)$ by conjugation: $(\gamma, \delta) \cdot \phi = \delta\phi(\gamma)^{-1}\phi(\gamma)\delta^{-1}$.

6.2. Lemma. *For any locally trivial (G, H) -bundle $\pi: P \rightarrow X$ over a G -fixed base X , the map $\tau: P \rightarrow \text{Hom}(G, H)$ is a Serre fibration.*

Proof. This will follow by showing that $(\tau, \pi): P \rightarrow \text{Hom}(G, H) \times X$ is actually fiber bundle. Since π is locally trivial, we can reduce to the case when π has the form $\pi: (G \times H)/\Lambda_\phi \times U \rightarrow U$, where $\Lambda_\phi \leq G \times H$ is the graph of some homomorphism $\phi: G \rightarrow H$. Then

$$(\tau, \pi) = \rho \times \text{id}_U: (G \times H)/\Lambda_\phi \times U \rightarrow \text{Hom}(G, H) \times U,$$

where $\rho: (G \times H)/\Lambda_\phi \rightarrow \text{Hom}(G, H)$ sends $[\gamma, \delta] \mapsto \delta\phi(\gamma)^{-1}\phi(\delta\phi(\gamma)^{-1})^{-1}$. Because $\text{Hom}(G, H)$ is topologically a coproduct of orbits under H -conjugation (1.3), (7.1), we see that ρ is isomorphic to the composite of a projection map $(G \times H)/\Lambda_\phi \rightarrow H/C_H(\phi)$ (induced by $(\gamma, \delta) \mapsto \delta\phi(\gamma)^{-1}$) with an open and closed immersion, and thus is a fibration. \square

6.3. Outline of the proof. To prove that the map $B_G H \rightarrow \text{Map}(EG, B_G H)$ (induced by restriction along $EG \rightarrow *$) is a G -equivariant weak equivalence, it suffices to show that it induces a weak equivalence of spaces $(B_G H)^{G'} \rightarrow \text{Map}(EG, B_G H)^{G'}$ for all closed subgroups $G' \leq G$. Without loss of generality, we may assume $G' = G$, since, when the group action is restricted to the subgroup G' , $B_G H$ is a $B_{G'} H$ and EG is an EG' . Thus, we will show that $(B_G H)^G \rightarrow \text{Map}(EG, B_G H)^G$ is an equivalence, using the following.

6.4. Lemma. *Suppose given a (G, H) -bundle $P \rightarrow X$ over a space X with trivial G -action, together with maps:*

- (1) $\alpha: X \rightarrow (B_G H)^G \subseteq B_G H$ classifying the G -equivariant H -bundle $P \rightarrow X$, i.e., covered by a (G, H) -bundle map $P \rightarrow E_G H$, and
- (2) $\rho: X \rightarrow \text{Map}(BG, BH)$, whose adjoint $\tilde{\rho}: X \times BG = (X \times EG)/G \rightarrow BH$ classifies the H -bundle $(P \times EG)/G \rightarrow (X \times EG)/G$, i.e., covered by an H -bundle map $(P \times EG)/G \rightarrow EH$.

Then the diagram

$$(6.5) \quad \begin{array}{ccc} X & \xrightarrow{\alpha} & (B_G H)^G \\ \rho \downarrow & & \downarrow \\ \text{Map}(BG, BH) & \xrightarrow{\sim} & \text{Map}(EG, B_G H)^G \end{array}$$

commutes up to homotopy. (The bottom map is induced by a G -equivariant map $BH \rightarrow B_G H$, where G acts trivially on BH , and is a weak equivalence.)

Proof. The adjoints of both composite maps $X \rightarrow \text{Map}(EG, B_G H)^G$ are G -equivariant maps $(X \times EG)/G \rightarrow B_G H$, which in either case are covered by maps $(P \times EG)/G \rightarrow E_G H$ of (G, H) -bundles. The result follows from the universal property of $B_G H$, as the classifying space for such bundles. \square

The strategy is as follows. Fix compact Lie groups G and H , and take $\rho: X \rightarrow \text{Map}(BG, BH)$ in (6.4) to be isomorphic to the map $(\text{Hom}(G, H) \times EH)/H \rightarrow \text{Map}(BG, BH)$ described in §5, which for 1-truncated H gives the weak equivalence of (1.2). We will

- (1) construct a certain (G, H) -bundle $P \rightarrow X$ (where G acts trivially on X),

(2) prove that a map $\alpha: X \rightarrow B_G H$ classifying $P \rightarrow X$ induces a weak equivalence $X \xrightarrow{\sim} (B_G H)^G \subseteq B_G H$, and

(3) construct a bundle map $(P \times EG)/G \rightarrow EH$ covering the adjoint $\tilde{\rho}: X \times BG \rightarrow BH$ to ρ .

Thus by (6.4) both α and ρ fit in a homotopy commutative square (6.5). It follows that if H is 1-truncated, (1.2) implies that ρ is a weak equivalence, from which it follows that the right-hand vertical arrow is a weak equivalence, which is the desired result. Note: the hypothesis that H is 1-truncated is used only to show that ρ (which exists for arbitrary H) is a weak equivalence.

6.6. Step 1: Construction of $P \rightarrow X$. As in the previous section, we consider categories $\text{Fun}(G, H)$ and $(H \curvearrowright H)$ (internal to Top), where G and H are compact Lie groups. Consider the topological category C defined as the fiber product

$$C := \text{Fun}(G, H) \times_H (H \curvearrowright H)$$

via the evident restriction functors $\text{Fun}(G, H) \rightarrow \text{Fun}(\{e\}, H) = H$ and $(H \curvearrowright H) \rightarrow (H \curvearrowright *) = H$. (Here H represents a topological category with one object.)

The group $G \times H$ acts on C via

$$(\gamma, \delta) \cdot (\phi_0, h_0) = (\phi_0, \phi_0(\gamma)h_0\delta^{-1}),$$

on objects and

$$(\gamma, \delta) \cdot (\phi_0 \xrightarrow{h} \phi_1, h_0 \xrightarrow{h} h_1) = (\phi_0 \xrightarrow{h} \phi_1, \phi_0(\gamma)h_0\delta^{-1} \xrightarrow{h} \phi_1(\gamma)h_1\delta^{-1})$$

on morphisms, $(\gamma, \delta) \in G \times H$. (This works exactly because $h\phi_0(\gamma) = \phi_1(\gamma)h$.) The evident projection functor $C \rightarrow \text{Fun}(G, H)$ is invariant under the $G \times H$ action on C , with respect to the trivial action on $\text{Fun}(G, H)$.

We set $P := \|NC\|$ and $X := \|N\text{Fun}(G, H)\|$, with $P \rightarrow X$ induced by the evident projection functor. It is straightforward to show that the induced $G \times H$ -action on P is compatible with the projection map to X , and that H acts freely on P with $P/H \approx X$. In particular, $P \rightarrow X$ has the structure of a G -equivariant principal H -bundle.

We note an equivalent description of C , and hence of P . Let $C' := \text{Hom}(G, H) \times N(H \curvearrowright H)$, where $\text{Hom}(G, H)$ is viewed as a topological category with only identity maps. There is an isomorphism $C' \rightarrow C$ of topological categories, given on objects and morphisms by

$$(\phi, h_0) \mapsto (h_0\phi h_0^{-1}, h_0), \quad (\phi, h_0 \xrightarrow{h} h_1) \mapsto (h_0\phi h_0^{-1} \xrightarrow{h} h_1\phi h_1^{-1}, h_0 \xrightarrow{h} h_1).$$

The $G \times H$ -action on C' induced by this isomorphism is described by

$$(\gamma, \delta) \cdot (\phi, h_0) = (\delta\phi(\gamma)^{-1}\phi\phi(\gamma)\delta^{-1}, h_0\phi(\gamma)\delta^{-1}),$$

$$(\gamma, \delta) \cdot (\phi, h_0 \xrightarrow{h} h_1) = (\delta\phi(\gamma)^{-1}\phi\phi(\gamma)\delta^{-1}, h_0\phi(\gamma)\delta^{-1} \xrightarrow{h} h_1\phi(\gamma)\delta^{-1}).$$

In particular, the projection functor $C' \rightarrow \text{Hom}(G, H)$ induces a $G \times H$ -equivariant map $P \rightarrow \text{Hom}(G, H)$ (using the conjugation $G \times H$ action on $\text{Hom}(G, H)$), and this map is a non-equivariant weak equivalence, since $P \approx \|NC'\| \approx \text{Hom}(G, H) \times EH$.

6.7. Step 2: The weak equivalence $\alpha: X \rightarrow (B_G H)^G$. Choose any $X \rightarrow B_G H$ classifying the $P \rightarrow X$ constructed above (this exists because X is paracompact and completely regular, so numerable), and so covered by a $G \times H$ -equivariant map $P \rightarrow E_G H$. Since the action of G on X is trivial, these factor through $\alpha: X \rightarrow (B_G H)^G$ and $\alpha': P \rightarrow p^{-1}((B_G H)^G)$, where $p: E_G H \rightarrow B_G H$ is the universal bundle.

6.8. Lemma. *The map $\alpha': P \rightarrow p^{-1}((B_G H)^G)$ is a weak equivalence of underlying spaces.*

Proof. The map α' fits in a commutative diagram,

$$\begin{array}{ccc} P & \xrightarrow{\alpha'} & p^{-1}((B_G H)^G) \\ & \searrow \tau & \swarrow \tau \\ & \text{Hom}(G, H) & \end{array}$$

where by (6.2) both maps marked τ are Serre fibrations. The fibers of these τ s over $\phi \in \text{Hom}(G, H)$ are EH and $(E_G H)^{\Lambda_\phi}$ respectively, both of which spaces are contractible. Thus α' is a weak equivalence (as are both τ s). \square

It follows that $\alpha: X \rightarrow (B_G H)^G$ is a weak equivalence, as it is obtained by the quotient of α' by free H -actions.

6.9. Step 3: The bundle map covering $\rho: X \rightarrow \text{Map}(BG, BH)$. We have a commutative square of functors

$$\begin{array}{ccc} C \times (G \curvearrowright G) = \text{Fun}(G, H) \times_H (H \curvearrowright H) \times (G \curvearrowright G) & \longrightarrow & (H \curvearrowright H) \\ \downarrow & & \downarrow \\ \text{Fun}(G, H) \times (G \curvearrowright G) & \longrightarrow & H \end{array}$$

where the vertical arrows are the evident projections, the top horizontal arrow is given by

$$(\phi_0, h_0, g_0) \mapsto (\phi_0(g_0)h_0), \quad (\phi_0 \xrightarrow{h} \phi_1, h_0 \xrightarrow{h} h_1, g_0 \xrightarrow{g} g_1) \mapsto (\phi_0(g_0)h_0 \xrightarrow{h\phi_0(g)=\phi_1(g)h} \phi_1(g_1)h_1).$$

on objects and morphisms, and the bottom horizontal arrow is given by

$$(\phi_0, g_0) \mapsto *, \quad (\phi_0 \xrightarrow{h} \phi_1, g_0 \xrightarrow{g} g_1) \mapsto (* \xrightarrow{h\phi_0(g)=\phi_1(g)h} *).$$

The group G acts on the objects on the left-hand side of the square, where G acts on C as described above, by the tautological right action on $(G \curvearrowright G)$, and trivially on $\text{Fun}(G, H)$. The horizontal arrows are invariant under this G action.

Thus, taking geometric realizations of nerves and passing to quotients by G -actions, we obtain a commutative square

$$\begin{array}{ccc} (P \times EG)/G & \longrightarrow & EH \\ \downarrow & & \downarrow \\ (X \times EG)/G & \longrightarrow & BH \end{array}$$

which is evidently a map of H -bundles. Under the identification $(X \times EG)/G \approx X \times BG \approx N(\text{Fun}(G, H) \times G)$, we see that the bottom arrow is isomorphic to that obtained from the evaluation functor $\text{Fun}(G, H) \times G \rightarrow H$, and thus is adjoint to the map $\rho: X \rightarrow \text{Map}(BG, BH)$ described earlier.

7. THE SPACE OF HOMOMORPHISMS BETWEEN COMPACT LIE GROUPS

Recall that $\text{Hom}(G, H)$ denotes the space of homomorphisms equipped with the compact-open topology. We give a proof of the following fact, which is standard but not easily read from the literature with which I am familiar.

7.1. Proposition. *Let G and H be Lie groups, with G compact. The map*

$$(\phi, hC(\phi)) \mapsto h\phi h^{-1}: \coprod_{[\phi]} H/C(\phi) \rightarrow \text{Hom}(G, H),$$

where $C(\phi) = \{h \in H \mid \phi(g)h = h\phi(g) \forall g \in G\}$, and $[\phi]$ runs over a set of H -conjugacy classes in $\text{Hom}(G, H)$, is a homeomorphism. In particular, $\text{Hom}(G, H)$ is locally compact, and thus a CGWH space.

Proof. We quote a classical theorem of Montgomery-Zippin [MZ55, p. 216]: for every compact subgroup K of a Lie group L , there exists a neighborhood U of K such that every closed subgroup of L in U is L -conjugate to a subgroup of K . Applied to $L = G \times H$, and $K = \Lambda_\phi = \{(g, \phi(g)) \mid g \in G\}$, the graph of a continuous homomorphism $\phi: G \rightarrow H$, we obtain a neighborhood $U \subseteq G \times H$ of Λ_ϕ such that if $\Lambda_{\phi'} \in U$ for $\phi' \in \text{Hom}(G, H)$, then ϕ' is H -conjugate to ϕ [CF64, 38.1].

There exists a neighborhood V of $e \in H$ such that $\Lambda_\phi \subseteq \{(g, h) \mid h\phi(g)^{-1} \in V\} \subseteq U$. To see this, use the homeomorphism $\alpha: G \times H \rightarrow G \times H$, $\alpha(g, h) = (g, h\phi(g)^{-1})$, together with the tube lemma applied to $G \times \{e\} \subseteq \alpha(U)$.

By definition, the set $V' := \{f: G \rightarrow H \mid f(G) \subseteq V\}$ is an open subset of $C(G, H)$, the space of continuous maps $G \rightarrow H$ equipped with the compact-open topology. The space $C(G, H)$ is a topological group under pointwise multiplication in H ; to prove this, use the fact that G , H , and finite products thereof are locally compact, so that the relevant evaluation maps are continuous. Therefore, the translated subset $V'\phi$ is open in $C(G, H)$. Tracing through the definitions, we see that any continuous homomorphism $G \rightarrow H$ in $V'\phi$ must be conjugate to ϕ .

Thus, we have shown that conjugacy classes are open subsets of $\text{Hom}(G, H)$.

Now consider the map of the proposition. Each $H/C(\phi)$ maps bijectively to a conjugacy class in $\text{Hom}(G, H)$. As H is Hausdorff, so is $C(G, H)$ and hence so is the subspace $\text{Hom}(G, H)$. Therefore, each $H/C(\phi) \rightarrow \text{Hom}(G, H)$ gives a homeomorphism to its image, since $H/C(\phi)$ is compact. Because the image is also open, the homeomorphism of the proposition follows.

As an immediate consequence, we see that $\text{Hom}(G, H)$ is a coproduct of compact Hausdorff spaces, and thus locally compact. \square

REFERENCES

- [Bro99] Ronald Brown, *Groupoids and crossed objects in algebraic topology*, Homology Homotopy Appl. **1** (1999), 1–78 (electronic).
- [CF64] P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, Ergebnisse der Mathematik und ihrer Grenzgebiete, N. F., Band 33, Academic Press Inc., Publishers, New York, 1964.
- [GMM12] B. J. Guillou, J. P. May, and M. Merling, *Categorical models for equivariant classifying spaces* (2012), available at [arXiv:1201.5178](https://arxiv.org/abs/1201.5178).
- [Las82] R. K. Lashof, *Equivariant bundles*, Illinois J. Math. **26** (1982), no. 2, 257–271.
- [LM86] R. K. Lashof and J. P. May, *Generalized equivariant bundles*, Bull. Soc. Math. Belg. Sér. A **38** (1986), 265–271 (1987). MR885537
- [LMS83] R. K. Lashof, J. P. May, and G. B. Segal, *Equivariant bundles with abelian structural group*, Proceedings of the Northwestern Homotopy Theory Conference (Evanston, Ill., 1982), Contemp. Math., vol. 19, Amer. Math. Soc., Providence, RI, 1983, pp. 167–176.
- [LU14] Wolfgang Lück and Bernardo Uribe, *Equivariant principal bundles and their classifying spaces*, Algebr. Geom. Topol. **14** (2014), no. 4, 1925–1995.
- [May75] J. Peter May, *Classifying spaces and fibrations*, Mem. Amer. Math. Soc. **1** (1975), no. 1, 155, xiii+98.
- [May90] J. P. May, *Some remarks on equivariant bundles and classifying spaces*, Astérisque **191** (1990), 7, 239–253. International Conference on Homotopy Theory (Marseille-Luminy, 1988).
- [MZ55] Deane Montgomery and Leo Zippin, *Topological transformation groups*, Interscience Publishers, New York-London, 1955.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, IL
E-mail address: rezk@math.uiuc.edu