

Isogenies, power operations, and homotopy theory

Charles Rezk

University of Illinois at Urbana-Champaign

Seoul, August 18, 2014



Plan.

- Context: “power operations” in cohomology theories.
- Recent advances: Morava E -theories. Formal groups and isogenies.
- Applications and vistas.

Motivating example.

K-theory (Grothendieck; Atiyah-Hirzebruch; 1950s)

$$K(X) = K^0(X) := \left\{ \begin{array}{l} \text{isomorphism classes of} \\ \text{vector bundles } / X \end{array} \right\} / \sim$$
$$V \sim V_1 + V_2 \quad \text{if} \quad 0 \rightarrow V_1 \rightarrow V \rightarrow V_2 \rightarrow 0.$$

Functors on vector bundles give operations on $K(X)$, e.g.:

$$V, W \mapsto V \otimes W, \quad V \mapsto \Lambda^n V, \quad V \mapsto \text{Sym}^n V.$$

$K(X)$ is a Λ -ring (Grothendieck)

Functions $\lambda^n: K(X) \rightarrow K(X)$ satisfying axioms

$$\lambda^n(x + y) = \dots, \quad \lambda^n(xy) = \dots, \quad \lambda^m \lambda^n(x) = \dots$$

... = explicit polynomials in $\lambda^i(x), \lambda^j(y)$.

Compact Lie $G \curvearrowright X$.

Equivariant K -theory

$$K_G(X) = K(X // G) := \{G \text{ equivariant vb } / X\} / \sim .$$

(Atiyah, 1966) tensor power is an operation

$$V \mapsto V^{\otimes n}: \quad K_G(X) \rightarrow K_{G \times \Sigma_n}(X) \approx K_G(X) \otimes R\Sigma_n.$$

$K_G(\text{point}) = K(\text{point} // G) = RG =$ representation ring of G .

$\Sigma_n =$ symmetric group.

Λ -rings \iff representation theory of symmetric groups

Λ -ring structure is complicated to describe, but is easy for “nice” rings.

(Wilkerson, 1982)

R torsion free comm. ring:

$$\left\{ \begin{array}{c} \Lambda\text{-ring} \\ \text{structures on } R \end{array} \right\} \leftrightarrow \left\{ \begin{array}{c} \{\psi^p: R \rightarrow R\}_{p \text{ prime}} \text{ lifts of Frobenius,} \\ \psi^p \psi^q = \psi^q \psi^p. \end{array} \right\}$$

- Adams operations ψ^n , $n \geq 1$; $\psi^m \psi^n = \psi^{mn}$, ring homomorphisms
Adams congruence $\psi^p(x) \equiv x^p \pmod{p}$, p prime
- any Λ -ring has $\psi^p, \theta^p: R \rightarrow R$ satisfying

$$\psi^p \text{ is a ring homomorphism, } \psi^p(x) = x^p + p \theta^p(x)$$

(say θ^p is a *witness* to the p th Adams congruence)

Line bundle $L \rightarrow X \implies \psi^n(L) = L^{\otimes n}$ in $K(X)$

Multiplicative group scheme

$$\begin{aligned}\mathbb{G}_m &= \text{Spec}(\mathbb{Z}[T, T^{-1}]) \\ &\approx \text{Spec}(K(\text{pt} // U(1)))\end{aligned}$$

Adams operation \implies *isogeny* of \mathbb{G}_m :

$$(K(\text{pt} // U(1)) \xrightarrow{\psi^n} K(\text{pt} // U(1))) \iff \mathbb{G}_m \xrightarrow{[n]} \mathbb{G}_m$$

Isogeny: finite flat homomorphism of group schemes

Remarks.

- $\widehat{\mathbb{G}}_m = \text{Spf } K(BU(1))$, multiplicative *formal group*
- These properties useful in classical applications (e.g., Adams work on vector fields on spheres, image of J , ...)

Other examples of power operations

$h^*(-)$ = generalized cohomology theory, commutative ring valued

Would like to have

$$h^*(X) \xrightarrow{P^n} h_{\Sigma_n}^*(X) = h^*(X \times B\Sigma_n) \quad \text{refines of } n\text{th power } x \mapsto x^n$$

Do these exist? **Yes** if $h^*(-)$ represented by a *structured commutative ring spectrum* (= commutative S -algebra = E_∞ -ring spectrum = ...)

Examples.

- **(Steenrod, 1953)** reduced power operations in $H^*(-, \mathbb{F}_p)$
(Sq^i for $p = 2$, P^i for p odd)
- **(Voevodsky, 2001)** motivic reduced power operations
- **(Quillen, 1971)** power operations in bordism theories
based on $M \mapsto M^{\times n} \curvearrowright \Sigma_n$
used to prove $\pi_* MU$ classifies formal group laws

What is elliptic cohomology?

Theory	K -theory	elliptic cohomology
Group scheme	\mathbb{G}_m	elliptic curve
Cycles	vector bundles	???

??? = 2-dim conformal field theories? (**Segal, ...**)

Examples:

- **(Goerss-Hopkins-Miller)** tmf = “topological modular forms” associated to *universal* elliptic curve over $\mathcal{M}_{\mathrm{Ell}}$ structured comm ring spectrum \implies power operations!
- **(Lurie)** *Equivariant* elliptic cohomology theories

Open question: Which equivariant elliptic cohomology theories admit power operations?

Tate curve $T[[q]] = \mathbb{C}^\times / q^{\mathbb{Z}}$, defined over $\text{Spec } \mathbb{Z}[[q]]$.

Equivariant elliptic cohomology at Tate curve

$$\text{Ell}_{\text{Tate}}(X // G) \underset{\text{approx}}{:=} K\left(\mathcal{L}^{\text{ghost}}(X // G) // U(1)\right)$$

“ghost loops” = constant loops; RHS is K of “twisted sectors” (see e.g., Ruan 2000, Lupercio-Urbe 2002)

(Ganter, 2007, 2013) Power operations for Ell_{Tate}

$\text{Ell}_{\text{Tate}}(X // G)$ is an **elliptic Λ -ring**: two families of operations

$$\lambda^n: \text{Ell}_{\text{Tate}} \rightarrow \text{Ell}_{\text{Tate}}, \quad \mu^m: \text{Ell}_{\text{Tate}} \rightarrow \text{Ell}_{\text{Tate}} \otimes_{\mathbb{Z}[[q]]} \mathbb{Z}[[q^{1/m}]]$$

$\{\lambda^n\}$ are Λ -ring structure, $\{\mu^m\}$ are Λ -ring homomorphisms

- *Morava E -theories* are “designer cohomology theories” — manufactured using homotopy theory, not coming from “nature”
- some arise as completions of “natural” theories, e.g.

$$K_p^\wedge, \quad \mathrm{Ell}_{\mathrm{s.-s. point}}^\wedge$$

- have rich theory of power operations (Ando, Hopkins, Strickland, R.)

Goal: describe what we know about this theory (a lot)

Recall: Power operations for K -theory are “controlled” by isogenies of \mathbb{G}_m

Slogan

Power operations for Morava E -theories are “controlled” by “deformations” of Frobenius isogenies of 1-dimensional formal groups

Let $G_0/\mathbb{F}_p =$ one dimensional commutative formal group of height $n \in \{1, 2, \dots\}$.

(Morava, 1978; Goerss-Hopkins-Miller 1993–2004)

There exists a cohomology theory E_{G_0} (*Morava E -theory*) which

- is represented by a structured commutative ring spectrum
- is complex orientable; formal group $\mathrm{Spf}(E^0\mathbb{C}P^\infty) =$ *universal deformation* of G_0 (in sense of Lubin-Tate)

$$E_{G_0}^0(\mathrm{pt}) = \mathbb{Z}_p[[a_1, \dots, a_{n-1}]]$$

$$E_{G_0}^*(\mathrm{pt}) = E_{G_0}^0(\mathrm{pt})[u, u^{-1}], \quad u \in E_{G_0}^2(\mathrm{pt})$$

Formal groups and complex oriented theories

Formal group is object locally described by a formal group law.

Formal group law (commutative, 1-dimensional)

$S(x, y) \in R[[x, y]]$ satisfying axioms for abelian group:

$$\begin{aligned}S(x, 0) &= x = S(0, x), \\S(x, y) &= S(y, x), \\S(S(x, y), z) &= S(x, S(y, z)).\end{aligned}$$

For future reference, we note the p -series of G_0 :

$$[p](x) = \underbrace{S(x, S(x, \dots S(x, x)))}_{x \text{ appears } p \text{ times}}$$

Complex oriented cohomology theory

Ring-valued cohomology theory E such that $E^*(\mathbb{C}P^\infty) = E^*[[x]]$, and x restricts to fundamental class of $\mathbb{C}P^1 = S^2$.

Examples: $H^*(-, \mathbb{Z})$, K -theory, Ell, **Morava E -theories**,...

Deformations of formal groups

G_0/\mathbb{F}_p formal group of *height* n (i.e., $[p]_{G_0}(x) = cx^{p^n} + O(x^{p^n+1})$, $c \neq 0$)
 R = complete local ring, $\mathbb{F}_p \subset R/\mathfrak{m}$

Groupoid $\text{Def}_{G_0}^0(R)$ of deformations of G_0/\mathbb{F}_p to R

Deformation (G, α) :

- G is a formal group over R ,
- iso $\alpha: G_0 \xrightarrow{\sim} G_{R/\mathfrak{m}}$ of formal groups over \mathbb{F}_p

Isomorphism $(G, \alpha) \rightarrow (G', \alpha')$ of deformations:

- iso $f: G \rightarrow G'$ compatible with id of G_0

Classified up to canonical iso by Lubin and Tate:

(Lubin-Tate, 1966)

\exists *universal deformation* $(G_{\text{univ}}, \alpha_{\text{univ}})$ over $A \approx \mathbb{Z}_p[[a_1, \dots, a_{n-1}]]$

G_{univ} is the formal group of Morava E -theory E_{G_0}

Isogeny of formal groups over R

Homomorphism $f: G \rightarrow G'$ given locally over R by $f(x) = cx^n +$ higher degree terms, $c \in R^\times$. ($n = \deg f$)

G_0/\mathbb{F}_p has a distinguished family of *Frobenius isogenies*

$$\text{Frob}^r: G_0 \rightarrow G_0, \quad r \geq 0,$$

given locally by $\text{Frob}^r(x) = x^{p^r}$.

Category $\text{Def}_{G_0}(R)$ of deformations of Frobenius

Objects:

- deformations (G, α) to R (= objects of $\text{Def}_{G_0}^0(R)$)

Morphisms $(G, \alpha) \rightarrow (G', \alpha')$:

- isogenies $f: G \rightarrow G'$ compatible with $\text{Frob}^r: G_0 \rightarrow G_0$, some $r \geq 0$

The “pile” $\text{Def} = \text{Def}_{G_0}$

We have assignments

complete local ring $R \iff$ category $\text{Def}(R)$

local homomorphism $R \rightarrow R' \iff$ functor $\text{Def}(R) \rightarrow \text{Def}(R')$

If $\text{Def}(R)$ were a groupoid, we would call it a (pre-)stack

Def is the “pile” of deformations of powers of Frob

Sheaves on Def

A *sheaf of modules* on Def is a collection of functors

$$A_R: \text{Def}(R) \rightarrow (R\text{-modules})$$

with compatibility wrt base change along local homomorphisms $R \rightarrow R'$

Likewise, a *sheaf of commutative rings* on Def is ...

Notation: $\text{Mod}(\text{Def})$, $\text{Com}(\text{Def})$.

$\text{Mod}(\text{Def}) = \text{Mod}(\Gamma)$ for a certain ring Γ

(Ando-Hopkins-Strickland 2004; see R. 2009)

Power operations make Morava E -cohomology E_{G_0} a functor

$$E^*(-): \text{Spaces} \rightarrow \text{Com}^*(\text{Def})$$

Key step (**Strickland 1997, 1998**):

$E^0 B\Sigma_{p^r} / I$ classifies subgroups of rank p^r of deformations

Broader context: We have $E^*(X) = \pi_*(E^{X_+})$ where $A = E^{X_+}$ is

- (i) a structured commutative E -algebra spectrum,
- (ii) $K(n)$ -local ($\Leftrightarrow \pi_* A$ complete wrt (a_1, \dots, a_{n-1}) in a suitable sense)

The real theorem is

(**ibid**)

π_* lifts to a functor

$$\pi_*: h\text{Com}(E)_{K(n)} \rightarrow \text{Com}^*(\text{Def})$$

on homotopy category of $K(n)$ -local commutative E -algebra spectra

$\text{Mod}(\text{Def}) =$ modules for a certain ring Γ

Height 1

$G_0 =$ multiplicative formal group; $E_{G_0} = K_p^\wedge$

$$\Gamma = \mathbb{Z}_p[\psi^p] \quad \text{gen. by Adams operation } \psi^p$$

Height 2 (R., arXiv:0812.1320)

$G_0/\mathbb{F}_2 =$ completion of s.-s. elliptic curve $y^2 + y = x^3$ over \mathbb{F}_2

$$\Gamma = \mathbb{Z}_2[[a]]\langle Q_0, Q_1, Q_2 \rangle \left/ \begin{array}{l} Q_0 a = a^2 Q_0 - 2a Q_1 + 6 Q_2 \\ Q_1 a = 3 Q_0 + a Q_2 \\ Q_2 a = -a Q_0 + 3 Q_1 \\ Q_1 Q_0 = 2 Q_2 Q_1 - 2 Q_0 Q_2 \\ Q_2 Q_0 = Q_0 Q_1 + a Q_0 Q_2 - 2 Q_1 Q_2 \end{array} \right)$$

(Y. Zhu, 2014) gives similar description at height 2, $p = 3$

There is a uniform description of Γ/p at height 2, all primes p

Properties of Γ

$n =$ height of G_0/\mathbb{F}_p

(Ando 1995)

$\text{Center}(\Gamma) = \mathbb{Z}_p[\tilde{T}_1, \dots, \tilde{T}_n]$, (*Hecke algebra*)

(R. arXiv:1204.4831)

Γ is *quadratic*, i.e.,

$$\Gamma \approx \text{Tensor alg.}(C_1) / (\text{ideal gen. by } C_2)$$

where C_1 and $C_2 \subseteq C_1 \otimes_{E_0} C_1$ are $E_0 = \mathbb{Z}_p[[a_1, \dots, a_{n-1}]]$ bimodules

(ibid)

Γ is *Koszul*: have Γ -bimodule resolution

$$0 \leftarrow \Gamma \leftarrow \Gamma \otimes_{E_0} C_0 \otimes_{E_0} \Gamma \leftarrow \cdots \leftarrow \Gamma \otimes_{E_0} C_n \otimes_{E_0} \Gamma \leftarrow 0,$$

each C_k is E_0 -bimod, free and f.g. as right E_0 -mod; $C_0 = E_0$

$\implies \text{gl. dim}(\Gamma) = 2n$

(R. *ibid*)

Γ is Koszul

- This was conjectured by Ando-Hopkins-Strickland
- It is purely a theorem about formal algebraic geometry
- Only general proof is a purely “topological” proof, using ingredients:
 - (1) $\Gamma =$ “primitives” of the Hopf algebra $\bigoplus_{m \geq 0} E_0(B\Sigma_m)$ (**Strickland**)
 - (2) bar complex of Γ in degree k is “primitives” in $\bigoplus_{m_1, \dots, m_k} E_0 B(\Sigma_{m_1} \wr \dots \wr \Sigma_{m_k})$
 - (3) vanishing results for Bredon homology of partition complexes with coeff. in appropriate Mackey functors (**Arone-Dwyer-Lesh 2013**)

Proof inspired by role of partition complexes as “derivatives of identity functor” in Goodwillie’s functor calculus

(R. 2012) purely alg. geom. proof in height 2 case, using results on moduli of subgroups of elliptic curves

Homotopy groups of $K(n)$ -local E -algebras have some more structure:

$$\pi_*: h\mathrm{Com}(E)_{K(n)} \rightarrow (T\text{-algebras})$$

“ T -algebras” = a complicated algebraic category (like Λ -rings)

(R. 2009)

R p -torsion free commutative E_* -algebra:

$$\left\{ \begin{array}{l} T\text{-algebra} \\ \text{structures on } R \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} A \in \mathrm{Com}^*(\mathrm{Def}) \text{ with } A(G_{\mathrm{univ}}) = R \\ \text{satisfying “Frobenius congruence”} \end{array} \right\}$$

Frobenius congruence: $Qx \equiv x^p \pmod{pR}$ for a certain $Q \in \Gamma$

There is a (non-additive) *witness* to the Frobenius congruence:

$$\theta: R \rightarrow R \quad \text{satisfying} \quad Qx = x^p + p\theta(x)$$

where R is a T -algebra

Application 1: nilpotence

Easy consequence of existence of “witness” θ such that
 $Q(x) = x^p + p\theta(x)$, $Q(x + y) = Q(x) + Q(y)$:

If $A \in \text{Com}(E)_{K(n)}$, then

$$x \in \pi_* A, \quad p^r x = 0 \quad \implies \quad x^{(p+1)^r} = 0.$$

Idea: deduce relation $\theta(px) = p^{p-1}x - Q(x) = (p^{p-1} - 1)x^p - p\theta(x)$.
If $px = 0$, then $0 = x\theta(px) = -x^{p+1}$.

Mathew-Noel-Naumann observe this, and use it (with Nilpotence Theorem of Devinatz-Hopkins-Smith) to give an easy proof of a conjecture of May:

(Mathew-Noel-Naumann 2014)

If $R =$ structured commutative ring spectrum, then the kernel of the Hurewicz map

$$\pi_* R \rightarrow H_*(R, \mathbb{Z})$$

consists of nilpotent elements

Application 2: units and orientations

$A = \text{structured commutative ring} \implies \text{units spectrum } \mathrm{gl}_1 A$

$$(\mathrm{gl}_1 A)^0(X) = (A^0(X))^\times$$

Question. Does there exist structured commutative ring map $MG \rightarrow A$, where $MG = \text{spectrum representing bordism}$ ($G \in \{U, SU, O, SO, \text{Spin}, \dots\}$)?

Answer (May-Quinn-Ray-Tornehave 1977). Yes iff the composite

$$g \rightarrow o \xrightarrow{J} \mathrm{gl}_1 S \rightarrow \mathrm{gl}_1 A$$

is null-homotopic as map of spectra, where $g = \text{infinite delooping of } G$

(Ando-Hopkins-R.; see Hopkins 2002)

There is a map of structured commutative ring spectra

$$M\text{String} \rightarrow \text{tmf}$$

which realizes the “Witten genus”; $\text{String} = \text{six-connected cover of Spin}$

Logarithmic operation: spectrum map $\ell: \mathrm{gl}_1 A \rightarrow A$

(tom Dieck 1989)

$A = K_p^\wedge$: exists $\ell: \mathrm{gl}_1 K_p^\wedge \rightarrow K_p^\wedge$, giving $\ell: K_p^\wedge(X)^\times \rightarrow K_p^\wedge(X)$ by

$$\begin{aligned} \ell(x) &= \log(x) - \frac{1}{p} \log(\psi^p(x)) & \log &= \text{Taylor exp. at 1} \\ &= \frac{1}{p} \log(x^p / \psi^p(x)) & \psi^p(x) &\equiv x^p \pmod{p} \\ &= \sum_{m \geq 1} (-1)^m \frac{p^{m-1}}{m} (\theta^p(x)/x)^m & \psi^p(x) &= x^p + p\theta^p(x) \end{aligned}$$

(R. 2006)

$E = E_{G_0}$, height $G_0 = n$; exists $\ell: \mathrm{gl}_1 E \rightarrow E$ giving $E^0(X)^\times \rightarrow E^0(X)$ by

$$\ell(x) = \sum_{k=0}^n (-1)^k p^{\binom{k}{2}-k} \log \tilde{T}_k(x)$$

where $\tilde{T}_k \in \mathbb{Z}_p[\tilde{T}_1, \dots, \tilde{T}_n] = \text{Center}(\Gamma)$

(Ando-Hopkins-R.)

Exists $MString \rightarrow tmf$ realizing Witten genus

Must construct null-homotopy of $\alpha: string \rightarrow gl_1 tmf$

Proof idea:

- Above techniques give “locally defined” logarithms $\ell_n: gl_1 tmf_p^\wedge \rightarrow tmf_{K(n)}$, $n = 1, 2$, all primes p
- Work one prime at a time; have “fracture squares”

$$\begin{array}{ccc}
 gl_1 tmf_p^\wedge & \xrightarrow{\ell_2} & tmf_{K(2)} \\
 \ell_1 \downarrow & & \downarrow \iota_1 \\
 tmf_{K(1)} & \xrightarrow{\gamma} & (tmf_{K(1)})_{K(2)}
 \end{array}$$

- $\text{Map}(string, tmf_{K(2)}) \approx *$, so reduce to $string \rightarrow \text{HoFib}(\gamma)$
- Explicit formulas for ℓ_1, ℓ_2 identify $\gamma = \iota_2 \circ (\text{id} - U)$, where $U: tmf_{K(1)} \rightarrow tmf_{K(1)}$ is topological lift of “Atkin operator” on p -adic modular forms

Application 3: derived indecomposables

Commutative ring k ; augmented comm. k -algebra $\pi: R \rightarrow k$

Indecomposables

$$Q_k(R) := I/I^2, \quad I = \text{Ker}(\pi: R \rightarrow k)$$

(“cotangent space at π ”)

$$T_k(R) := \text{Hom}_k(Q_k(R), k)$$

(“tangent space at π ”)

Commutative ring spectrum k ; augmented comm. k -algebra $\pi: R \rightarrow k$

(Basterra 1999, Basterra-Mandell 2005) Derived version

$$TQ_k(R) := “I/I^2” = \text{hocolim } \Omega_{nu}^n \Sigma_{nu}^n I,$$

nu = non-unital k -algebras

$$TT_k(R) := \underline{\text{Hom}}_k(TQ_k(R), k)$$

Also called *reduced topological André-Quillen homology/cohomology*

HA = Eilenberg-MacLane spectrum, representing $H^*(-, A)$

(Sullivan 1977)

X = simply connected f. type space; $H\mathbb{Q}^{X_+}$ = rational cochains spectrum

$$\pi_* TT_{H\mathbb{Q}}(H\mathbb{Q}^{X_+}) \approx \pi_* X \otimes \mathbb{Q}$$

(Mandell 2006)

X = simply connected f. type space; $H\overline{\mathbb{F}}_p^{X_+}$ = mod p cochains spectrum

$$\pi_* TT_{H\overline{\mathbb{F}}_p}(H\overline{\mathbb{F}}_p^{X_+})_p^\wedge \approx 0$$

(X_p^\wedge) can be recovered from $H\overline{\mathbb{F}}_p^{X_+}$, but not this way

Q: Are there structured commutative rings R that behave like $H\mathbb{Q}$?

Yes: $K(n)$ -local R , such as Morava E -theories

(Behrens-R., in progress)

E = Morava E -theory at height n ; $X = S^{2d-1}$ odd dimensional sphere

$$\pi_* TT_E(E^{X_+}) \approx E^* \Phi_n X$$

Bousfield-Kuhn functor $\Phi_n: \text{Spaces}_* \rightarrow \text{Spectra}_{K(n)}$

Φ_n carries part of the “ v_n -local homotopy groups of X ”

Spectral sequence computing derived tangent space

E = Morava E -theory, $\pi_* R$ smooth over $\pi_* E$,

$$E_{s,t}^2 = \text{Ext}_\Gamma^s(\omega^{-1/2} \otimes Q_{\pi_* E}(\pi_* R), \omega^{(t-1)/2} \otimes \text{nul}) \implies \pi_* TT_E(R)$$

$\omega^{t/2} \approx \tilde{E}^0(S^t)$, $\text{nul} = E_0$ with trivial Γ -action; $E_{s,t}^2 = 0$ if $s > n$

Combine

$$E_{s,t}^2 = \text{Ext}_\Gamma^s(\omega^{d-1}, \omega^{(t-1)/2} \otimes \text{nul}) \implies E^* \Phi_n S^{2d-1}$$

Recovers known calc at $n = 1$; collapses to $E^* \Phi_2 S^{2d-1} = \text{Ext}^2$ at $n = 2$

Q: Does equivariant elliptic cohomology admit power operations?

- Analogue of Def:

Isog = “pile” of all elliptic curves and isogenies between them
 $\implies \text{Mod}(\text{Isog}), \text{Com}(\text{Isog})$

- $\text{Mod}(\text{Isog})$ has analog of Koszul property
 $\text{Mod}(\text{Isog}_p)$ has homological dimension 2 rel to $\text{Qcoh}(\mathcal{M}_{\text{Ell}})$
- Known power operations for Ell_{Tate} and $\text{Ell}_{\text{s.-s.}}^\wedge$ are consistent with this picture

Conjecturally, equivariant elliptic cohomologies which are etale over \mathcal{M}_{Ell} should be “classified” by the etale site of Isog

<http://www.math.uiuc.edu/~rezk/rezk-icm-2014-slides.pdf>

- Ando, "Isogenies of formal group laws and power operations in the cohomology theories E_n ", Duke J., 1995
- Ando, Hopkins, Strickland, "The sigma orientation is an H_∞ map", Amer. J. 2004; arXiv:math/0204053
- Arone, Dwyer, Lesh, "Bredon homology of partition complexes"; <http://arxiv.org/abs/1306.0056>
- Atiyah, "Power operations in K -theory", Q. J. Math Oxford, 1966
- Basterra, "André-Quillen cohomology of commutative S -algebras", J. Pure Appl. Alg., 1999
- Basterra, Mandell, "Homology and cohomology of E_∞ -ring spectra", Math. Z., 2005; arXiv:math/0407209
- Ganter, "Stringy power operations in Tate K -theory", Homology, Homotopy, Appl., 2013; arXiv:math/0701565
- Ganter, "Power operations in orbifold Tate K -theory"; arXiv:1301.2754
- Goerss, Hopkins, "Moduli spaces of commutative ring spectra", LMS Lecture Notes 315, 2004
- Hopkins, "Algebraic topology and modular forms", Proc. ICM, 2002
- Lubin, Tate, "Formal moduli for one-parameter formal Lie groups", Bull. Soc. Math. France, 1966
- Lupercio, Uribe, "Loop groupoids, gerbes, and twisted sectors on orbifolds", Contemp. Math. 310, 2002; arXiv:math/0110207
- Mathew, Noel, Naumann, "On a nilpotence conjecture of J. P. May"; <http://arxiv.org/abs/1403.2023>
- May, Quinn, Ray, Tornehave, " E_∞ -ring spaces and E_∞ -ring spectra", Springer LNM 577, 1977
- Morava, "Completions of complex cobordism", Springer LNM 658, 1978
- Rezk, "The units of a ring spectrum and a logarithmic cohomology operation", J. AMS, 2006; arXiv:math/0407022
- Rezk, "Power operations for Morava E -theory of height 2 at the prime 2"; arXiv:0812.1320.
- Rezk, "The congruence criterion for power operations in Morava E -theory", Homology, Homotopy Appl., 2009; arXiv:0902.2499
- Rezk, "Rings of power operations for Morava E -theories are Koszul"; <http://arxiv.org/abs/1204.4831>
- Ruan, "Stringy geometry and topology of orbifolds", Contemp. Math. 310, 2002; arXiv:math/0011149
- Strickland, "Finite subgroups of formal groups", J. Pure Appl. Alg., 1997
- Strickland, "Morava E -theory of symmetric groups", Topology, 1998; arXiv:math/9801125
- Sullivan, "Infinitesimal computations in topology", IHES, 1977
- tom Dieck, "The Artin-Hasse logarithm for λ -rings", Springer LNM 1370, 1989
- Wilkerson, " A -rings, binomial domains, and vector bundles over $\mathbb{C}P^\infty$ ", Comm. Alg., 1982
- Zhu, "The power operation structure on Morava E -theory of height 2 and prime 3", Alg. & Geom. Top., 2014