

Sums of powers of binary quadratic forms

Bruce Reznick

University of Illinois at Urbana-Champaign

Joint Mathematical Meetings

AMS Special Session on Gaussian Graphical Models and
Combinatorial Algebraic Geometry

Baltimore, ND January 4, 2017 (slightly corrected 1/10/17)

I'd like to begin with a simple question. Consider a sum of two cubes of quadratic forms:

$$\sum_{j=1}^2 (\alpha_{j0}x^2 + \alpha_{j1}xy + \alpha_{j2}y^2)^3 = \sum_{k=0}^6 c_k x^{6-k} y^k,$$

I'd like to begin with a simple question. Consider a sum of two cubes of quadratic forms:

$$\sum_{j=1}^2 (\alpha_{j0}x^2 + \alpha_{j1}xy + \alpha_{j2}y^2)^3 = \sum_{k=0}^6 c_k x^{6-k} y^k,$$

One can view the seven c_k 's as cubic polynomials in the six $\alpha'_{j\ell}$'s, and since $7 > 6$, we know that the c_k 's must be algebraically dependent. There are $\binom{n+6}{6}$ monomials in the c_j 's of degree n ; these are forms of degree $3n$ in the $\alpha'_{j\ell}$'s, which comprise a vector space of dimension $\binom{3n+5}{5}$. And, eventually,

$$\binom{n+6}{6} > \binom{3n+5}{5}$$

so there must be dependence at degree n .

Unfortunately,

$$\binom{1442 + 6}{6} > \binom{3 \times 1442 + 5}{5}$$

is the first time this occurs. Various smart computational algebraic geometers I've asked over the years have been unable to brute force this relation.

Unfortunately,

$$\binom{1442 + 6}{6} > \binom{3 \times 1442 + 5}{5}$$

is the first time this occurs. Various smart computational algebraic geometers I've asked over the years have been unable to brute force this relation.

Is there a better way to do this? The answer is yes, and it turns out that there is a relation of degree $n = 15$ in the c_j 's, admittedly with more than 1000 terms.

Unfortunately,

$$\binom{1442 + 6}{6} > \binom{3 \times 1442 + 5}{5}$$

is the first time this occurs. Various smart computational algebraic geometers I've asked over the years have been unable to brute force this relation.

Is there a better way to do this? The answer is yes, and it turns out that there is a relation of degree $n = 15$ in the c_j 's, admittedly with more than 1000 terms.

This quindecim form is derived from a simple characterization of binary sextics which are a sums of two cubes of binary quadratics.

Unfortunately,

$$\binom{1442 + 6}{6} > \binom{3 \times 1442 + 5}{5}$$

is the first time this occurs. Various smart computational algebraic geometers I've asked over the years have been unable to brute force this relation.

Is there a better way to do this? The answer is yes, and it turns out that there is a relation of degree $n = 15$ in the c_j 's, admittedly with more than 1000 terms.

This quindecim form is derived from a simple characterization of binary sextics which are a sums of two cubes of binary quadratics.

Let's go to the characterization. There are two equivalent statements.

Theorem

Suppose p is a binary sextic. Then p is a sum of two cubes of quadratics if and only if:

Theorem

Suppose p is a binary sextic. Then p is a sum of two cubes of quadratics if and only if:

(i) p is a perfect cube or $p = f_1 f_2 f_3$, where the f_i 's are linearly dependent but non-proportional quadratic forms.

Theorem

Suppose p is a binary sextic. Then p is a sum of two cubes of quadratics if and only if:

(i) p is a perfect cube or $p = f_1 f_2 f_3$, where the f_i 's are linearly dependent but non-proportional quadratic forms.

(ii) There exists an invertible linear change of variables after which p equals either $g(x^2, y^2)$ or $\ell^3 g$ for some linear form ℓ , where g is a cubic which is a sum of two cubes (i.e., $g \neq \ell_1^2 \ell_2$.)

Theorem

Suppose p is a binary sextic. Then p is a sum of two cubes of quadratics if and only if:

(i) p is a perfect cube or $p = f_1 f_2 f_3$, where the f_i 's are linearly dependent but non-proportional quadratic forms.

(ii) There exists an invertible linear change of variables after which p equals either $g(x^2, y^2)$ or $\ell^3 g$ for some linear form ℓ , where g is a cubic which is a sum of two cubes (i.e., $g \neq \ell_1^2 \ell_2$.)

The proof of (i) is part of a more general result about sums of two cubes of forms.

Theorem

Suppose p is a binary sextic. Then p is a sum of two cubes of quadratics if and only if:

(i) p is a perfect cube or $p = f_1 f_2 f_3$, where the f_i 's are linearly dependent but non-proportional quadratic forms.

(ii) There exists an invertible linear change of variables after which p equals either $g(x^2, y^2)$ or $\ell^3 g$ for some linear form ℓ , where g is a cubic which is a sum of two cubes (i.e., $g \neq \ell_1^2 \ell_2$.)

The proof of (i) is part of a more general result about sums of two cubes of forms.

The proof of (ii) relies on the ancient art of simultaneous diagonalization: if q and r are two binary quadratic forms, then either they share a common factor, or they can be simultaneously diagonalized.

Theorem

Suppose p is a binary sextic. Then p is a sum of two cubes of quadratics if and only if:

(i) p is a perfect cube or $p = f_1 f_2 f_3$, where the f_i 's are linearly dependent but non-proportional quadratic forms.

(ii) There exists an invertible linear change of variables after which p equals either $g(x^2, y^2)$ or $\ell^3 g$ for some linear form ℓ , where g is a cubic which is a sum of two cubes (i.e., $g \neq \ell_1^2 \ell_2$.)

The proof of (i) is part of a more general result about sums of two cubes of forms.

The proof of (ii) relies on the ancient art of simultaneous diagonalization: if q and r are two binary quadratic forms, then either they share a common factor, or they can be simultaneously diagonalized.

(Also note: an even form under $(x, y) \mapsto (x + y, x - y)$ becomes a symmetric form, and vice versa.)

Theorem (Either mine, or very old and obscure, or both)

Theorem (Either mine, or very old and obscure, or both)

Suppose $F \in \mathbb{C}[x_1, \dots, x_n]$. Then $F = G^3 + H^3$ for forms G, H if and only if either $F = K^3$, or $F = G_1 G_2 G_3$, where the G_j 's are non-proportional, but linearly dependent factors.

Theorem (Either mine, or very old and obscure, or both)

Suppose $F \in \mathbb{C}[x_1, \dots, x_n]$. Then $F = G^3 + H^3$ for forms G, H if and only if either $F = K^3$, or $F = G_1 G_2 G_3$, where the G_j 's are non-proportional, but linearly dependent factors.

Proof.

First $G^3 + H^3 = (G + H)(G + \omega H)(G + \omega^2 H)$, where $\omega = e^{\frac{2\pi i}{3}}$, and if two of the factors $G + \omega^j H$ are proportional, then so are G and H , and hence F is a cube. In any event, please observe that $(G + H) + \omega(G + \omega H) + \omega^2(G + \omega^2 H) = 0$.

Theorem (Either mine, or very old and obscure, or both)

Suppose $F \in \mathbb{C}[x_1, \dots, x_n]$. Then $F = G^3 + H^3$ for forms G, H if and only if either $F = K^3$, or $F = G_1 G_2 G_3$, where the G_j 's are non-proportional, but linearly dependent factors.

Proof.

First $G^3 + H^3 = (G + H)(G + \omega H)(G + \omega^2 H)$, where $\omega = e^{\frac{2\pi i}{3}}$, and if two of the factors $G + \omega^j H$ are proportional, then so are G and H , and hence F is a cube. In any event, please observe that $(G + H) + \omega(G + \omega H) + \omega^2(G + \omega^2 H) = 0$.

Conversely, if F has such a factorization, there exist $0 \neq \alpha, \beta \in \mathbb{C}$ so that $F = G_1 G_2 (\alpha G_1 + \beta G_2)$. It is easily checked that

$$3\alpha\beta(\omega - \omega^2)F = (\omega^2\alpha G_1 - \omega\beta G_2)^3 - (\omega\alpha G_1 - \omega^2\beta G_2)^3.$$

Note that $3\alpha\beta(\omega - \omega^2) = 3\sqrt{-3} \alpha\beta \neq 0$. □

In any particular case, if $\deg F = 3r$, there are, up to multiple, only $\frac{(3r)!}{3!(r!)^3}$ ways to write F as a product of three factors of degree r , so checking this condition is algorithmic.

In any particular case, if $\deg F = 3r$, there are, up to multiple, only $\frac{(3r)!}{3!(r!)^3}$ ways to write F as a product of three factors of degree r , so checking this condition is algorithmic.

In particular, if $F(x, y)$ is a binary cubic form, then it has three linear factors $\ell_j(x, y) = \alpha_j x + \beta_j y$, and these are always dependent. Thus, as Sylvester and our 19th century predecessors knew, a binary cubic F is a sum of two cubes unless it has a square factor (and isn't a cube). We use this a lot.

The second case uses a simple old lemma whose proof is omitted.

Lemma

Two quadratic forms $q_1(x, y)$ and $q_2(x, y)$ either have a common linear factor, or can be simultaneously diagonalized; that is, $q_j(ax + by, cx + dy) = \rho_j x^2 + \sigma_j y^2$.

The second case uses a simple old lemma whose proof is omitted.

Lemma

Two quadratic forms $q_1(x, y)$ and $q_2(x, y)$ either have a common linear factor, or can be simultaneously diagonalized; that is, $q_j(ax + by, cx + dy) = \rho_j x^2 + \sigma_j y^2$.

Thus, if $p = q_1^t + q_2^t$, where q_j is quadratic, then either the q_j 's have a common linear factor (and $p = \ell^t g$, where g is a sum of two linear t -th powers), or after a linear change of variables,

$$p(ax + by, cx + dy) = \sum_{j=1}^2 (\rho_j x^2 + \sigma_j y^2)^t;$$

That is, $p(ax + by, cx + dy) = g(x^2, y^2)$, where g again is a sum of two linear t -th powers (typical for $t = 3$, not for $t > 3$.)

Checking if p is even after a change of variables is also algorithmic.

$$\begin{aligned} p(x, y) &= \prod_{j=0}^{2d-1} (x - \lambda_j y) \implies \\ p(ax + by, cx + dy) &= p(a, -c) \prod_{j=0}^{2d-1} \left(x - \left(\frac{\lambda_j d - b}{a - \lambda_j c} \right) y \right) \\ &:= p(a, -c) \prod_{j=0}^{2d-1} (x - \mu_j y). \end{aligned}$$

Thus, the roots of p (taking ∞ if $y \mid p$) are mapped by a Möbius transformation. If $\tilde{p}(x, y) = p(ax + by, cx + dy)$ is even, then $T(z) = -z$ is an involution on the roots, say $T(\mu_{2j}) = \mu_{2j+1}$. It follows that there is an involutory Möbius transformation U permuting the d pairs of roots of p ; to be specific:

$$\lambda_{2j+1} = \frac{2ad - (ad + bc)\lambda_{2j}}{(ad + bc) - 2cd\lambda_{2j}}.$$

The algorithm is this: Given p , find the roots λ_j , and for each quadruple $\lambda_{i_1}, \lambda_{i_2}, \lambda_{i_3}, \lambda_{i_4}$, define the Möbius transformation U so that $U(\lambda_{i_1}) = \lambda_{i_2}$, $U(\lambda_{i_2}) = \lambda_{i_1}$ and $U(\lambda_{i_3}) = \lambda_{i_4}$ and see if it permutes the others. There are instances in which more than one U may work; for example, if p is both even and symmetric.

The algorithm is this: Given p , find the roots λ_j , and for each quadruple $\lambda_{i_1}, \lambda_{i_2}, \lambda_{i_3}, \lambda_{i_4}$, define the Möbius transformation U so that $U(\lambda_{i_1}) = \lambda_{i_2}$, $U(\lambda_{i_2}) = \lambda_{i_1}$ and $U(\lambda_{i_3}) = \lambda_{i_4}$ and see if it permutes the others. There are instances in which more than one U may work; for example, if p is both even and symmetric.

Don't get me wrong. Complications abound. Here's a simple one. Consider the even sextic

$$p(x, y) = x^6 - x^4y^2 - x^2y^4 + y^6 = (x^2 - y^2)^2(x^2 + y^2).$$

Here, $p(x, y) = g(x^2, y^2)$, where $g(x, y) = (x - y)^2(x + y)$ (having a square factor) is unfortunately not a sum of two cubes. On the other hand, if $\gamma = \frac{2}{\sqrt{3}}i$, then

$$\begin{aligned} p(x, y) &= (x^2 + 2xy + y^2)(x^2 + y^2)(x^2 - 2xy + y^2) \implies \\ 2p(x, y) &= (x^2 + \gamma xy + y^2)^3 + (x^2 - \gamma xy + y^2)^3. \end{aligned}$$

Now let's suppose our given cubic p is a sum of two cubes, factor it and expand it in the usual way. Write p as

$$\sum_{k=0}^6 c_k x^{6-k} y^k = c_0 \left(x^6 + \sum_{k=1}^6 e_k x^{6-k} y^k \right) = c_0 \prod_{j=1}^6 (x + r_j y),$$

where the e_k 's are the elementary symmetric functions.

Now let's suppose our given cubic p is a sum of two cubes, factor it and expand it in the usual way. Write p as

$$\sum_{k=0}^6 c_k x^{6-k} y^k = c_0 \left(x^6 + \sum_{k=1}^6 e_k x^{6-k} y^k \right) = c_0 \prod_{j=1}^6 (x + r_j y),$$

where the e_k 's are the elementary symmetric functions.

There are 15 ways to divide the 6 r_j 's into 3 pairs of roots, and the condition that the quadratic factors be dependent for some choice of factorization is equivalent to the vanishing of

$$H(r) := \prod_{\ell=1}^{15} \begin{vmatrix} 1 & 1 & 1 \\ r_{\sigma_\ell(1)} + r_{\sigma_\ell(2)} & r_{\sigma_\ell(3)} + r_{\sigma_\ell(4)} & r_{\sigma_\ell(5)} + r_{\sigma_\ell(6)} \\ r_{\sigma_\ell(1)} r_{\sigma_\ell(2)} & r_{\sigma_\ell(3)} r_{\sigma_\ell(4)} & r_{\sigma_\ell(5)} r_{\sigma_\ell(6)} \end{vmatrix}.$$

Now let's suppose our given cubic p is a sum of two cubes, factor it and expand it in the usual way. Write p as

$$\sum_{k=0}^6 c_k x^{6-k} y^k = c_0 \left(x^6 + \sum_{k=1}^6 e_k x^{6-k} y^k \right) = c_0 \prod_{j=1}^6 (x + r_j y),$$

where the e_k 's are the elementary symmetric functions.

There are 15 ways to divide the 6 r_j 's into 3 pairs of roots, and the condition that the quadratic factors be dependent for some choice of factorization is equivalent to the vanishing of

$$H(r) := \prod_{\ell=1}^{15} \begin{vmatrix} 1 & 1 & 1 \\ r_{\sigma_\ell(1)} + r_{\sigma_\ell(2)} & r_{\sigma_\ell(3)} + r_{\sigma_\ell(4)} & r_{\sigma_\ell(5)} + r_{\sigma_\ell(6)} \\ r_{\sigma_\ell(1)} r_{\sigma_\ell(2)} & r_{\sigma_\ell(3)} r_{\sigma_\ell(4)} & r_{\sigma_\ell(5)} r_{\sigma_\ell(6)} \end{vmatrix}.$$

This is an I-really-hope-it's-symmetric (and it is) polynomial of degree 45 in the r_j 's.

Mathematica can compute $H(r)$ without too much difficulty, and in 11657.87 seconds transform it into a function in the e_k 's of degree 15. Now write $e_k = c_k/c_0$, make the substitution and multiply by c_0^{15} to get the relation. It has 1360 terms, so I won't write it here. (I also need to express it in terms of the fundamental invariants of the binary sextic, and haven't done so yet.) It is *isobaric* in the old sense, each monomial $\prod c_k^{m_k}$ has $\sum m_k = 15$, $\sum km_k = 45$.

Mathematica can compute $H(r)$ without too much difficulty, and in 11657.87 seconds transform it into a function in the e_k 's of degree 15. Now write $e_k = c_k/c_0$, make the substitution and multiply by c_0^{15} to get the relation. It has 1360 terms, so I won't write it here. (I also need to express it in terms of the fundamental invariants of the binary sextic, and haven't done so yet.) It is *isobaric* in the old sense, each monomial $\prod c_k^{m_k}$ has $\sum m_k = 15$, $\sum km_k = 45$. You can use it to check that a generic pencil of sextic contains a finite number of sums of two cubes, but this method doesn't distinguish repeated factors. So it says that C^2 should be a sum of two cubes for any cubic C , because $C = l_1 l_2 l_3$ and $\{l_1^2, l_2 l_3, l_2 l_3\}$ is dependent. Bad. On the other hand, $(x^3 + y^3)^2 + ax^5 y$ is provably a sum of two cubes if and only if

$$a^3 \in \left\{ \frac{3^6}{5^5} \cdot (13 \pm 5\sqrt{145}) \right\}.$$

Mathematica can compute $H(r)$ without too much difficulty, and in 11657.87 seconds transform it into a function in the e_k 's of degree 15. Now write $e_k = c_k/c_0$, make the substitution and multiply by c_0^{15} to get the relation. It has 1360 terms, so I won't write it here. (I also need to express it in terms of the fundamental invariants of the binary sextic, and haven't done so yet.) It is *isobaric* in the old sense, each monomial $\prod c_k^{m_k}$ has $\sum m_k = 15$, $\sum km_k = 45$. You can use it to check that a generic pencil of sextic contains a finite number of sums of two cubes, but this method doesn't distinguish repeated factors. So it says that C^2 should be a sum of two cubes for any cubic C , because $C = l_1 l_2 l_3$ and $\{l_1^2, l_2 l_3, l_2 l_3\}$ is dependent. Bad. On the other hand, $(x^3 + y^3)^2 + ax^5 y$ is provably a sum of two cubes if and only if

$$a^3 \in \left\{ \frac{3^6}{5^5} \cdot (13 \pm 5\sqrt{145}) \right\}.$$

Not sure what that means.

Here is a nice conjecture due to Boris Shapiro. Let $H_m(\mathbb{C}^2)$ denote the vector space of binary forms of degree m with complex coefficients, and suppose $m = de$, $d, e \in \mathbb{N}$.

Here is a nice conjecture due to Boris Shapiro. Let $H_m(\mathbb{C}^2)$ denote the vector space of binary forms of degree m with complex coefficients, and suppose $m = de$, $d, e \in \mathbb{N}$.

Conjecture

Every $p \in H_{de}(\mathbb{C}^2)$ can be written as a sum of d d -th powers of forms in $H_e(\mathbb{C}^2)$.

Here is a nice conjecture due to Boris Shapiro. Let $H_m(\mathbb{C}^2)$ denote the vector space of binary forms of degree m with complex coefficients, and suppose $m = de$, $d, e \in \mathbb{N}$.

Conjecture

Every $p \in H_{de}(\mathbb{C}^2)$ can be written as a sum of d d -th powers of forms in $H_e(\mathbb{C}^2)$.

- The first point to make is that this assertion is a *universal* statement, not a *generic* one.

Here is a nice conjecture due to Boris Shapiro. Let $H_m(\mathbb{C}^2)$ denote the vector space of binary forms of degree m with complex coefficients, and suppose $m = de$, $d, e \in \mathbb{N}$.

Conjecture

Every $p \in H_{de}(\mathbb{C}^2)$ can be written as a sum of d d -th powers of forms in $H_e(\mathbb{C}^2)$.

- The first point to make is that this assertion is a *universal* statement, not a *generic* one.
- If $e = 1$, this conjecture is a familiar statement to those who work with Waring rank, and the binary forms of degree d which require d d -th powers of linear forms are precisely those of the shape $(ax + by)^{d-1}(a'x + b'y)$, $ab' \neq a'b$; ie $\ell^{d-1}\ell'$.

Here is a nice conjecture due to Boris Shapiro. Let $H_m(\mathbb{C}^2)$ denote the vector space of binary forms of degree m with complex coefficients, and suppose $m = de$, $d, e \in \mathbb{N}$.

Conjecture

Every $p \in H_{de}(\mathbb{C}^2)$ can be written as a sum of d d -th powers of forms in $H_e(\mathbb{C}^2)$.

- The first point to make is that this assertion is a *universal* statement, not a *generic* one.
- If $e = 1$, this conjecture is a familiar statement to those who work with Waring rank, and the binary forms of degree d which require d d -th powers of linear forms are precisely those of the shape $(ax + by)^{d-1}(a'x + b'y)$, $ab' \neq a'b$; ie $\ell^{d-1}\ell'$.
- If $d = 1$, there is nothing to prove.

- If $d = 2$, then $m = 2e$ is even, and p can be factored into linear factors, so that $p = fg$ for $f, g \in H_e(\mathbb{C}^2)$ and

$$p = fg = \left(\frac{f+g}{2}\right)^2 - \left(\frac{f-g}{2}\right)^2$$

is a sum of two squares (one could write this with “ i ” inside.)

- If $d = 2$, then $m = 2e$ is even, and p can be factored into linear factors, so that $p = fg$ for $f, g \in H_e(\mathbb{C}^2)$ and

$$p = fg = \left(\frac{f+g}{2}\right)^2 - \left(\frac{f-g}{2}\right)^2$$

is a sum of two squares (one could write this with “ i ” inside.)

- The conjecture is true generically. Using the classical Lasker-Wakeford approach, if $de + 1 = k(e + 1) + s$, $0 \leq s \leq e$, then

$$\sum_{j=1}^k (\alpha_{j0}x^e + \dots)^d + (\beta_0x^e + \dots + \beta_{s-1}x^{e-(s-1)}y^{s-1})^d$$

is a canonical form for binary forms of degree de , and

$$\left\lceil \frac{de+1}{e+1} \right\rceil \leq \left\lceil \frac{de+d}{e+1} \right\rceil = d.$$

- The conjecture is true if you remove the restriction to forms (but lose the information about degrees). In fact, every polynomial is a sum of d d -th powers of polynomials by a result of Newman-Slater. Let ζ_d denote a primitive d -th root of unity and p be a polynomial in any number of variables. Then the usual orthogonality properties of roots of unity imply that

$$d^2 p = \sum_{k=0}^{d-1} \zeta_d^{-k} (1 + \zeta_d^k p)^d.$$

- The conjecture is true if you remove the restriction to forms (but lose the information about degrees). In fact, every polynomial is a sum of d d -th powers of polynomials by a result of Newman-Slater. Let ζ_d denote a primitive d -th root of unity and p be a polynomial in any number of variables. Then the usual orthogonality properties of roots of unity imply that

$$d^2 p = \sum_{k=0}^{d-1} \zeta_d^{-k} (1 + \zeta_d^k p)^d.$$

- I now sketch an algorithmic proof for the simplest non-obvious case — $d = 3$, $e = 2$ — that is, every complex binary sextic is a sum of three cubes of quadratic forms.

Part of an ongoing project with Hal Schenck and Boris Shapiro I think.

Part of an ongoing project with Hal Schenck and Boris Shapiro I think.

Theorem

There is an algorithm for writing every binary sextic in $\mathbb{C}[x, y]$ as a sum of three cubes of quadratic forms.

Part of an ongoing project with Hal Schenck and Boris Shapiro I think.

Theorem

There is an algorithm for writing every binary sextic in $\mathbb{C}[x, y]$ as a sum of three cubes of quadratic forms.

Part of an ongoing project with Hal Schenck and Boris Shapiro I think.

Theorem

There is an algorithm for writing every binary sextic in $\mathbb{C}[x, y]$ as a sum of three cubes of quadratic forms.

Write the binary sextic (warning: different notation) as

$$p(x, y) = \sum_{k=0}^6 \binom{6}{k} a_k x^{6-k} y^k.$$

Given $p \neq 0$, we may always make an invertible change of variables to ensure that $p(0, 1)p(1, 0) \neq 0$; hence, assume $a_0 a_6 \neq 0$.

Part of an ongoing project with Hal Schenck and Boris Shapiro I think.

Theorem

There is an algorithm for writing every binary sextic in $\mathbb{C}[x, y]$ as a sum of three cubes of quadratic forms.

Write the binary sextic (warning: different notation) as

$$p(x, y) = \sum_{k=0}^6 \binom{6}{k} a_k x^{6-k} y^k.$$

Given $p \neq 0$, we may always make an invertible change of variables to ensure that $p(0, 1)p(1, 0) \neq 0$; hence, assume $a_0 a_6 \neq 0$.

By an observation of *ad hoc*,

$$\begin{aligned} q(x, y) &= x^2 + \frac{2a_1}{a_0} x y + \frac{5a_0 a_2 - 4a_1^2}{a_0^2} y^2 \\ \implies a_0 q^3(x, y) &= a_0 x^6 + 6a_1 x^5 y + 15a_2 x^4 y^2 + \dots \end{aligned}$$

Thus there always exists a cubic c such that

$$p(x, y) - a_0q(x, y)^3 = y^3c(x, y).$$

Thus there always exists a cubic c such that

$$p(x, y) - a_0q(x, y)^3 = y^3c(x, y).$$

Usually, $(p - a_0q^3)/y^3 = c$ is a sum of 2 cubes of linear forms, from which it follows that p is a sum of 3 cubes. As we've seen, this only fails if c has a square factor. The discriminant of $c(x, y)$ is a non-zero polynomial in the a_i 's of degree 18, divided by a_0^{14} , assuming that Mathematica is reliable, so this works for general p .

Thus there always exists a cubic c such that

$$p(x, y) - a_0q(x, y)^3 = y^3c(x, y).$$

Usually, $(p - a_0q^3)/y^3 = c$ is a sum of 2 cubes of linear forms, from which it follows that p is a sum of 3 cubes. As we've seen, this only fails if c has a square factor. The discriminant of $c(x, y)$ is a non-zero polynomial in the a_i 's of degree 18, divided by a_0^{14} , assuming that Mathematica is reliable, so this works for general p .

We now consider the remaining cases in which this first approach fails. Such a failure will have the shape

$$p(x, y) = (ax^2 + bxy + cy^2)^3 + y^3(rx + sy)^2(tx + uy)$$

where $ru - st \neq 0$, so that $c(x, y)$ genuinely is not a sum of two cubes.

Let $p_T(x, y) = p(x, Tx + y)$ and write

$$p_T(x, y) = \sum_{k=0}^6 \binom{6}{k} a_k(T) x^{6-k} y^k.$$

Here, a_k is a polynomial in T of degree $6 - k$ and $a_6(T) = a_6 \neq 0$. There are at most 6 values of T which must be avoided to ensure that $a_0(T) \neq 0$.

Let $p_T(x, y) = p(x, Tx + y)$ and write

$$p_T(x, y) = \sum_{k=0}^6 \binom{6}{k} a_k(T) x^{6-k} y^k.$$

Here, a_k is a polynomial in T of degree $6 - k$ and $a_6(T) = a_6 \neq 0$. There are at most 6 values of T which must be avoided to ensure that $a_0(T) \neq 0$.

Repeating the same construction as above to p_T , we find that the discriminant is a polynomial of degree 72 in T with coefficients in $\{a, b, c, r, s, t, u\}$ and tens of thousands of terms. It turns out, tediously, that for every *non-trivial* choice of (a, b, c, d, r, s, t, u) , this discriminant gives a non-zero polynomial in T . (Cased out, not trusting in "Solve".)

Let $p_T(x, y) = p(x, Tx + y)$ and write

$$p_T(x, y) = \sum_{k=0}^6 \binom{6}{k} a_k(T) x^{6-k} y^k.$$

Here, a_k is a polynomial in T of degree $6 - k$ and $a_6(T) = a_6 \neq 0$. There are at most 6 values of T which must be avoided to ensure that $a_0(T) \neq 0$.

Repeating the same construction as above to p_T , we find that the discriminant is a polynomial of degree 72 in T with coefficients in $\{a, b, c, r, s, t, u\}$ and tens of thousands of terms. It turns out, tediously, that for every *non-trivial* choice of (a, b, c, d, r, s, t, u) , this discriminant gives a non-zero polynomial in T . (Cased out, not trusting in “Solve”.)

Hence by avoiding finitely many values of T , the previous argument will work successfully on p_T to give it as a sum of three cubes. We then reverse the invertible transformations and get an expression for p itself.

For example, suppose $p(x, y) = x^6 + x^5y + x^4y^2 + x^3y^3 + x^2y^4 + xy^5 + y^6$. Then

$$p(x, y) - \left(x^2 + \frac{1}{3}xy + \frac{2}{9}y^2\right)^3 = \frac{7}{729}y^3(54x^3 + 81x^2y + 99xy^2 + 103y^3).$$

For example, suppose $p(x, y) = x^6 + x^5y + x^4y^2 + x^3y^3 + x^2y^4 + xy^5 + y^6$. Then

$$p(x, y) - \left(x^2 + \frac{1}{3}xy + \frac{2}{9}y^2\right)^3 = \frac{7}{729}y^3(54x^3 + 81x^2y + 99xy^2 + 103y^3).$$

An application of Sylvester's algorithm shows that

$$54x^3 + 81x^2y + 99xy^2 + 103y^3 = m_1(78x + (173 - \sqrt{20153})y)^3 + m_2(78x + (173 + \sqrt{20153})y)^3,$$
$$m_1 = \frac{20153 + 134\sqrt{20153}}{354209128}, \quad m_2 = \frac{20153 - 134\sqrt{20153}}{354209128}$$

For example, suppose $p(x, y) = x^6 + x^5y + x^4y^2 + x^3y^3 + x^2y^4 + xy^5 + y^6$. Then

$$p(x, y) - \left(x^2 + \frac{1}{3}xy + \frac{2}{9}y^2\right)^3 = \frac{7}{729}y^3(54x^3 + 81x^2y + 99xy^2 + 103y^3).$$

An application of Sylvester's algorithm shows that

$$54x^3 + 81x^2y + 99xy^2 + 103y^3 = m_1(78x + (173 - \sqrt{20153})y)^3 + m_2(78x + (173 + \sqrt{20153})y)^3, \\ m_1 = \frac{20153 + 134\sqrt{20153}}{354209128}, \quad m_2 = \frac{20153 - 134\sqrt{20153}}{354209128}$$

This gives a simple sextic p as a sum of three cubes in an ugly way and gives no hint about the existence of the formula

$$p(x, y) = \sum_{\pm} \left(\frac{9 \pm \sqrt{-3}}{18}\right) \left(x^2 + \frac{1 \pm \sqrt{-3}}{2}xy + y^2\right)^3.$$

An alternative approach is to observe that for a sextic p , there is usually a quadratic q so that $p - q^3$ is even. (Look at the coefficients of x^5y, x^3y^3, xy^5 and solve the equations for the coefficients of q .) Then $p - q^3$ is a cubic in $\{x^2, y^2\}$ and so is usually a sum of two cubes of even quadratic form. If this doesn't work, apply it to p_T .

An alternative approach is to observe that for a sextic p , there is usually a quadratic q so that $p - q^3$ is even. (Look at the coefficients of x^5y, x^3y^3, xy^5 and solve the equations for the coefficients of q .) Then $p - q^3$ is a cubic in $\{x^2, y^2\}$ and so is usually a sum of two cubes of even quadratic form. If this doesn't work, apply it to p_T .

We do not know how to completely characterize the sets of sums of three cubes for a given p and what other symmetries those sets might have.

An alternative approach is to observe that for a sextic p , there is usually a quadratic q so that $p - q^3$ is even. (Look at the coefficients of x^5y, x^3y^3, xy^5 and solve the equations for the coefficients of q .) Then $p - q^3$ is a cubic in $\{x^2, y^2\}$ and so is usually a sum of two cubes of even quadratic form. If this doesn't work, apply it to p_T .

We do not know how to completely characterize the sets of sums of three cubes for a given p and what other symmetries those sets might have.

Or if a real binary sextic is a sum of three cubes of real quadratic forms. (Question of Lek-Heng Lim at a conference in China.)

An alternative approach is to observe that for a sextic p , there is usually a quadratic q so that $p - q^3$ is even. (Look at the coefficients of x^5y, x^3y^3, xy^5 and solve the equations for the coefficients of q .) Then $p - q^3$ is a cubic in $\{x^2, y^2\}$ and so is usually a sum of two cubes of even quadratic form. If this doesn't work, apply it to p_T .

We do not know how to completely characterize the sets of sums of three cubes for a given p and what other symmetries those sets might have.

Or if a real binary sextic is a sum of three cubes of real quadratic forms. (Question of Lek-Heng Lim at a conference in China.)

This heavy reliance on tools from *École de calcul ad hoc* can only take you so far. There are two natural next steps; based on the observation that $8 = 4 \times 2$ and $9 = 3 \times 3$. Is every binary octic a sum of four 4th powers of quadratic forms? Is every binary nonic a sum of three cubes of cubic (thanks GM!) forms? One more fun fact: according to the Oxford English Dictionary (as well as wikipedia), an obsolete term for the 4th power is *zenzizenzic*.

The octic (or *zenzizenzizenic*) case is interesting because a canonical form for the octics gives them as a sum of three fourth powers, and the Conjecture is still true even if some singular cases require one more.

The octic (or *zenzizenzizenic*) case is interesting because a canonical form for the octics gives them as a sum of three fourth powers, and the Conjecture is still true even if some singular cases require one more.

To sum up:

Conjecture

Every $p \in H_m(\mathbb{C}^2)$ can be written as a sum of d d -th powers of forms in $H_e(\mathbb{C}^2)$. This is true for $d = 1, d = 2, e = 1$ and for $(d, e) = (3, 2)$.

The octic (or *zenzizenzizenic*) case is interesting because a canonical form for the octics gives them as a sum of three fourth powers, and the Conjecture is still true even if some singular cases require one more.

To sum up:

Conjecture

Every $p \in H_m(\mathbb{C}^2)$ can be written as a sum of d d -th powers of forms in $H_e(\mathbb{C}^2)$. This is true for $d = 1, d = 2, e = 1$ and for $(d, e) = (3, 2)$.

Thank you for your attention. I also very much to thank Josephine, Rainer and Seth for the opportunity to speak here.

Thank you for your attention. I also very much to thank Josephine, Rainer and Seth for the opportunity to speak here.

$$\int_{\Omega} f_u dt.$$