



ELSEVIER

Contents lists available at ScienceDirect

Linear Algebra and its Applications

www.elsevier.com/locate/laa



On the Choi–Lam analogue of Hilbert’s 1888 theorem for symmetric forms

Charu Goel^{a,*}, Salma Kuhlmann^a, Bruce Reznick^b

^a Department of Mathematics and Statistics, University of Konstanz, Universitätsstrasse 10, 78457, Konstanz, Germany

^b Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, United States

ARTICLE INFO

Article history:

Received 25 September 2015

Accepted 18 January 2016

Available online xxxx

Submitted by P. Semrl

MSC:

11E76

11E25

05E05

ABSTRACT

A famous theorem of Hilbert from 1888 states that for given n and d , every positive semidefinite (psd) real form of degree $2d$ in n variables is a sum of squares (sos) of real forms if and only if $n = 2$ or $d = 1$ or $(n, 2d) = (3, 4)$. In 1976, Choi and Lam proved the analogue of Hilbert’s Theorem for symmetric forms by assuming the existence of psd not sos symmetric n -ary quartics for $n \geq 5$. In this paper we complete their proof by constructing explicit psd not sos symmetric n -ary quartics for $n \geq 5$.

© 2016 Elsevier Inc. All rights reserved.

Keywords:

Positive polynomials

Sums of squares

Symmetric forms

1. Introduction

A real form (homogeneous polynomial) f is called *positive semidefinite* (psd) if it takes only non-negative values and it is called a *sum of squares* (sos) if there exist other forms h_j so that $p = h_1^2 + \cdots + h_k^2$. The question whether every psd form of given degree

* Corresponding author.

E-mail addresses: charu.goel@uni-konstanz.de (C. Goel), salma.kuhlmann@uni-konstanz.de (S. Kuhlmann), reznick@math.uiuc.edu (B. Reznick).

and given number of variables can be written as a sum of squares of real forms has many ramifications and has been studied extensively. Since a psd form always has even degree, it is sufficient to consider this question for even degree forms. We refer to this question as (Q). The first significant result in this direction was given by D. Hilbert [9] in 1888. His celebrated theorem states that for given n and d , every psd form of degree $2d$ in n variables is sos if and only if $n = 2$ or $d = 1$ or $(n, 2d) = (3, 4)$.

The above answer to (Q) can be summarized by the following chart:

deg \ var	2	3	4	5	6	...
2	✓	✓	✓	✓	✓	...
4	✓	✓	×	×	×	...
6	✓	×	×	×	×	...
8	✓	×	×	×	×	...
⋮	⋮	⋮	⋮	⋮	⋮	⋮

where a tick (✓) denotes a positive answer to (Q), whereas a cross (×) denotes a negative answer to (Q).

Let $\mathcal{P}_{n,2d}$ and $\Sigma_{n,2d}$ denote the cone of psd and sos n -ary $2d$ -ic forms (i.e. forms of degree $2d$ in n variables) respectively. Hilbert made a careful study of quaternary quartics and ternary sextics, and demonstrated that $\Sigma_{3,6} \subsetneq \mathcal{P}_{3,6}$ and $\Sigma_{4,4} \subsetneq \mathcal{P}_{4,4}$. Moreover he showed that

$$\begin{aligned} &\text{if } \Sigma_{4,4} \subsetneq \mathcal{P}_{4,4} \text{ and } \Sigma_{3,6} \subsetneq \mathcal{P}_{3,6}, \text{ then} \\ &\Sigma_{n,2d} \subsetneq \mathcal{P}_{n,2d} \text{ for all } n \geq 3, 2d \geq 4 \text{ and } (n, 2d) \neq (3, 4). \end{aligned} \tag{1}$$

So it is sufficient to produce psd not sos forms in these two crucial cases of quaternary quartics and ternary sextics to get psd not sos forms in all remaining cases as in assertion (1) above. In those two cases Hilbert described a method to produce examples of psd not sos forms, which was “elaborate and unpractical” (see [3, p. 387]), so no explicit examples appeared in literature for next 80 years. Explicit examples with $(n, 2d) = (3, 6)$ were found by T.S. Motzkin [12] in 1967 and R.M. Robinson [14] in 1969; Robinson also found an explicit example with $(n, 2d) = (4, 4)$. M.D. Choi and T.Y. Lam [2–4] produced many more examples in the mid 1970’s. More examples were given later by B. Reznick [13] and K. Schmüdgen [15].

In 1976, Choi and Lam [3] considered the question when a psd form is sos for the special case when the form considered is moreover symmetric ($f(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = f(x_1, \dots, x_n) \forall \sigma \in S_n$). Let $S\mathcal{P}_{n,2d}$ and $S\Sigma_{n,2d}$ denote the set of symmetric psd and symmetric sos n -ary $2d$ -ic forms respectively. They demonstrated that it is enough to find symmetric psd not sos forms in the two crucial cases of n -ary quartics for $n \geq 4$ and ternary sextics to obtain symmetric psd not sos n -ary $2d$ -ics for all $n \geq 3, 2d \geq 4$ and $(n, 2d) \neq (3, 4)$ (see Proposition 2.2). They showed that the answer to this question is the same as the answer to (Q), by assuming the existence of psd not sos symmetric n -ary

quartics for $n \geq 5$. For the convenience of the reader we include the following citation from [3],

“the construction of $f_{n,4} \in \mathcal{SP}_{n,4} \setminus S\Sigma_{n,4}$ ($n \geq 4$) requires considerable effort, so we shall not go into the full details here. Suffice it to record the special form $f_{4,4} = \sum x^2y^2 + \sum x^2yz - 2xyzw$. Here the two summations denote the full symmetric sums (w.r.t. the variables x, y, z, w); hence the summation lengths are respectively 6 and 12”.

We complete their proof by constructing explicit psd not sos symmetric n -ary quartics for $n \geq 5$ (see Theorems 2.8 and 2.9). These theorems will be further used in [7], where we consider the question when an even symmetric psd form is sos (a form is even symmetric if it is symmetric and in each of its terms every variable has even degree).

2. Analogue of Hilbert’s 1888 theorem for symmetric forms

We revisit the question: for which n and d , will every symmetric psd form of degree $2d$ in n variables be sos? We refer to this question as $\mathcal{Q}(S)$.

Choi and Lam in [3] claimed that the answer to $\mathcal{Q}(S)$, that classifies the pairs $(n, 2d)$ for which every symmetric psd n -ary $2d$ -ic form is sos, is:

$$\mathcal{SP}_{n,2d} = S\Sigma_{n,2d} \text{ if and only if } n = 2 \text{ or } d = 1 \text{ or } (n, 2d) = (3, 4). \tag{2}$$

One direction of (2) follows from Hilbert’s Theorem. Conversely for proving $\mathcal{SP}_{n,2d} \subseteq S\Sigma_{n,2d}$ only if $n = 2$ or $d = 1$ or $(n, 2d) = (3, 4)$, they showed that it is enough to find $f \in \mathcal{SP}_{n,2d} \setminus S\Sigma_{n,2d}$ for all pairs $(n, 4)$ with $n \geq 4$ and for the pair $(3, 6)$, i.e. they demonstrated that

$$\begin{aligned} &\text{if } S\Sigma_{n,4} \subsetneq \mathcal{SP}_{n,4} \text{ for all } n \geq 4 \text{ and } S\Sigma_{3,6} \subsetneq \mathcal{SP}_{3,6}, \text{ then} \\ &S\Sigma_{n,2d} \subsetneq \mathcal{SP}_{n,2d} \text{ for all } n \geq 3, 2d \geq 4 \text{ and } (n, 2d) \neq (3, 4). \end{aligned} \tag{3}$$

Lemma 2.1. *Let $f \in \mathcal{F}_{n,2d}$ be a psd not sos form and p an irreducible indefinite form of degree r in $\mathbb{R}[x_1, \dots, x_n]$. Then $p^2f \in \mathcal{F}_{n,2d+2r}$ is also a psd not sos form.*

Proof. Clearly p^2f is psd. If $p^2f = \sum_k h_k^2$, then for every real tuple \underline{a} with $p(\underline{a}) = 0$, it follows that $(p^2f)(\underline{a}) = 0$. This implies $h_k^2(\underline{a}) = 0 \forall k$ (since h_k^2 is psd), and so on the real variety $p = 0$, we have $h_k = 0$ as well.

So (using [1, Theorem 4.5.1]), for each k , there exists g_k so that $h_k = pg_k$. This gives $f = \sum_k g_k^2$, which is a contradiction. \square

Proposition 2.2. *If $S\Sigma_{n,4} \subsetneq \mathcal{SP}_{n,4}$ for all $n \geq 4$ and $S\Sigma_{3,6} \subsetneq \mathcal{SP}_{3,6}$, then $S\Sigma_{n,2d} \subsetneq \mathcal{SP}_{n,2d}$ for all $n \geq 3, d \geq 2$ and $(n, 2d) \neq (3, 4)$.*

Proof. Suppose we have forms $f \in \mathcal{SP}_{n,2d} \setminus \mathcal{S}\Sigma_{n,2d}$ for all pairs $(n, 4)$ with $n \geq 4$, and for the pair $(3, 6)$. Then we can construct symmetric n -ary forms of higher degree by taking $(x_1 + \dots + x_n)^{2i} f$, which can be seen to be in $\mathcal{SP}_{n,2d+2i} \setminus \mathcal{S}\Sigma_{n,2d+2i} \forall i \geq 0$, by i applications of [Lemma 2.1](#) with $p = x_1 + \dots + x_n$. \square

For the pair $(3, 6)$, Robinson [\[14\]](#) constructed the symmetric ternary sextic form $R(x, y, z) := x^6 + y^6 + z^6 - (x^4y^2 + y^4z^2 + z^4x^2 + x^2y^4 + y^2z^4 + z^2x^4) + 3x^2y^2z^2$ and showed that it is psd but not sos. For the pair $(4, 4)$, Choi and Lam [\[3\]](#) gave the form $f_{4,4} \in \mathcal{SP}_{4,4} \setminus \mathcal{S}\Sigma_{4,4}$. So in view of [Proposition 2.2](#), it remains to find psd not sos symmetric n -ary quartics for $n \geq 5$.

We will now construct explicit forms $f \in \mathcal{SP}_{n,4} \setminus \mathcal{S}\Sigma_{n,4}$ for $n \geq 4$. For $n \geq 4$, consider the symmetric n -ary quartic (studied in [\[5\]](#))

$$L_n(\underline{x}) := m(n - m) \sum_{i < j} (x_i - x_j)^4 - \left(\sum_{i < j} (x_i - x_j)^2 \right)^2,$$

where $m = \lfloor \frac{n}{2} \rfloor$. We shall show that L_n is psd for all n and L_n is not sos for all odd $n \geq 5$.

We need an important result ([Theorem 2.3](#) below) of Choi, Lam and Reznick [\[5\]](#). The same argument was modified in [\[8, Theorem 2.3\]](#) to treat even symmetric n -ary octics for $n \geq 4$.

Theorem 2.3. *A symmetric n -ary quartic f is psd iff $f(\underline{x}) \geq 0$ for every $\underline{x} \in \mathbb{R}^n$ with at most two distinct coordinates (if $n \geq 4$), i.e. $\Lambda_{n,2} = \{\underline{x} \in \mathbb{R}^n \mid x_i \in \{r, s\}; r \neq s\}$ is a test set for symmetric n -ary quartics.*

Proof. See [\[16, Corollary 5.6\]](#) or [\[6, Corollary 3.11\]](#). \square

Remark 2.4. V. Timofte’s half degree principle [\[16\]](#) gives a complete generalization of the above theorem for both symmetric polynomials (i.e. invariant under the action of the group S_n) and even symmetric polynomials (i.e. invariant under the action of the group $S_n \times \mathbb{Z}_2^n$) of degree $2d$ in n variables. See [\[10\]](#) for an application of this principle to elementary symmetric functions.

For $n = 5$, $L_n(\underline{x})$ has been discussed by A. Lax and P.D. Lax. They showed [\[11, p. 72\]](#) that

$$A_5(\underline{x}) := \sum_{i=1}^5 \prod_{j \neq i} (x_i - x_j) = \frac{1}{8} L_5,$$

a psd symmetric quartic in five variables, is not sos.

Proposition 2.5. *L_n is psd for all n .*

Proof. In view of [Theorem 2.3](#), it is enough to prove that $L_n \geq 0$ on the test set $\Lambda_{n,2} = \{(\underbrace{r, \dots, r}_k, \underbrace{s, \dots, s}_{n-k}) \mid r \neq s \in \mathbb{R}; 0 \leq k \leq n\}$.

Now for $\underline{x} \in \Lambda_{n,2}$,

$$x_i - x_j = \begin{cases} \pm(r - s) \neq 0, & \text{for } k(n - k) \text{ terms,} \\ 0, & \text{otherwise} \end{cases}$$

so L_n takes the value

$$\begin{aligned} L_n(\underline{x}) &= m(n - m)k(n - k)(r - s)^4 - [k(n - k)(r - s)^2]^2 \\ &= k(n - k)(r - s)^4[(m - k)(n - m - k)], \end{aligned}$$

which is non-negative since there is no integer between m and $n - m$. \square

Definition 2.6. Let $\{0, 1\}^n$ be the set of all n -tuples $\underline{x} = (x_1, \dots, x_n)$ with $x_i \in \{0, 1\}$ for all $i = 1, \dots, n$. A subset $S \subset \{0, 1\}^n$ is called a **0/1 set** and $\underline{x} \in \{0, 1\}^n$ a **0/1 point**.

Lemma 2.7. Suppose $n \geq 4$ and $h(x_1, \dots, x_n)$ is a quadratic form that vanishes on all 0/1 points with m or $(m + 1)$ 1's, where $m = \lfloor \frac{n}{2} \rfloor$, i.e. $h(\underline{x}) = 0$ for all \underline{x} with m or $(m + 1)$ 1's and $\begin{cases} (m + 1) \text{ or } m \text{ 0's (respectively) for odd } n = 2m + 1; \\ m \text{ or } (m - 1) \text{ 0's (respectively) for even } n = 2m. \end{cases}$

Then h is identically zero.

Proof. Set $h(x_1, \dots, x_n) = \sum_{i=1}^n a_i x_i^2 + \sum_{i < j} a_{ij} x_i x_j$. Fix distinct i, j, k and let S , such that $|S| = m - 1$, be a set of indices not containing i, j, k . Then $h = 0$ on \underline{x} , where the 1's on \underline{x} occur precisely on $S \cup \{i\}$, $S \cup \{i, k\}$, $S \cup \{j\}$, $S \cup \{j, k\}$. So we have:

$$\begin{aligned} \text{on } S \cup \{i\} : \quad 0 &= \sum_{l \in S} a_l + a_i + \sum_{l < l' \in S} a_{ll'} + \sum_{l \in S} a_{il}, \\ \text{on } S \cup \{i, k\} : \quad 0 &= \sum_{l \in S} a_l + a_i + a_k + \sum_{l < l' \in S} a_{ll'} + \sum_{l \in S} a_{il} + \sum_{l \in S} a_{kl} + a_{ik}. \end{aligned}$$

Subtracting above two equations gives:

$$a_k + \sum_{l \in S} a_{kl} + a_{ik} = 0. \tag{4}$$

Doing the same with $S \cup \{j\}$ and $S \cup \{j, k\}$ gives:

$$a_k + \sum_{l \in S} a_{kl} + a_{jk} = 0. \tag{5}$$

Thus $a_{ik} = a_{jk}$ (from equations (4) and (5)).

Since i, j, k are arbitrary, $a_{ik} = a_{jk} = a_{jl}$ for any $l \neq i, j, k$. So all the coefficients of $x_i x_j$ (for $i \neq j$) in h are equal, say $a_{ij} = u; i \neq j$.

It follows from equation (4) that $a_k + mu = 0$. So $a_k = -mu \forall k$, which gives:

$$h(x_1, \dots, x_n) = u \left(-m \sum_{i=1}^n x_i^2 + \sum_{i < j} x_i x_j \right).$$

But then $h(\underbrace{1, \dots, 1}_m, 0, \dots, 0) = 0$ gives $u \left(-m(m) + \frac{m(m-1)}{2} \right) = 0$, which implies $u = 0$, which implies $h = 0$. \square

Theorem 2.8. *If $n \geq 5$ is odd, then L_n is not sos.*

Proof. Fix odd $n \geq 5, n = 2m + 1$. Then

$$L_{2m+1} = m(m+1) \sum_{i < j} (x_i - x_j)^4 - \left(\sum_{i < j} (x_i - x_j)^2 \right)^2.$$

If $L_{2m+1} = \sum_t h_t^2$, then $L_{2m+1}(\underline{x}) = 0 \Rightarrow$ each $h_t(\underline{x}) = 0$, for any $\underline{x} \in \mathbb{R}^n$.

In particular, $L_{2m+1}(\underline{x}) = 0$ when \underline{x} has m or $(m+1)$ 1's and $(m+1)$ or m 0's. So, $h_t(\underline{x}) = 0$ for \underline{x} with m or $(m+1)$ 1's and $(m+1)$ or m 0's respectively. Write

$$h_t(\underline{x}) = \sum_{i=1}^n a_i x_i^2 + \sum_{i < j} a_{ij} x_i x_j.$$

Then by Lemma 2.7, we get $h_t = 0$. Hence L_{2m+1} is not sos. \square

Now we construct $f \in SP_{2m,4} \setminus S\Sigma_{2m,4}$ for $m \geq 2$.

Unfortunately,

$$L_{2m}(\underline{x}) = \sum_{i < j} (x_i - x_j)^2 \left(- (x_1 + \dots + x_{2m}) + m(x_i + x_j) \right)^2$$

(see [6, Proposition 3.13]) is sos, and so we need a different example in $SP_{2m,4} \setminus S\Sigma_{2m,4}$. For $2m \geq 4$, let

$$C_{2m}(x_1, \dots, x_{2m}) := L_{2m+1}(x_1, \dots, x_{2m}, 0).$$

Trivially, C_{2m} is a symmetric $2m$ -ary quartic and psd. We shall show it is not sos.

Theorem 2.9. *For $m \geq 2, C_{2m}(x_1, \dots, x_{2m})$ is not sos.*

Proof. If $C_{2m} = \sum_t h_t^2$, then $C_{2m}(\underline{x}) = 0 \Rightarrow$ each $h_t(\underline{x}) = 0$, for any $\underline{x} \in \mathbb{R}^n$.

In particular, $C_{2m}(\underline{x}) = 0$ when \underline{x} has m or $(m+1)$ 1's and m or $(m-1)$ 0's. So, $h_t(\underline{x}) = 0$ for \underline{x} with m or $(m+1)$ 1's and m or $(m-1)$ 0's respectively.

Write

$$h_t(\underline{x}) = \sum_{i=1}^n a_i x_i^2 + \sum_{i < j}^n a_{ij} x_i x_j.$$

Then by Lemma 2.7, we get $h_t = 0$. Hence, C_{2m} is not sos. \square

To sum up, the answer to $\mathcal{Q}(S)$ can be summarized by the same chart as for Hilbert's Theorem, given in the Introduction.

Acknowledgements

This paper contains results from the Ph.D. Thesis [6]. Charu Goel is thankful to MFO for inviting her as Leibnitz-Graduate-Student in April 2014. Bruce Reznick wishes to thank the Simons Foundation: A Simons Visiting Professorship supported his visit to the University of Konstanz in April 2014 and the Simons Collaboration Grant 280987 supported his participation at MFO.

References

- [1] J. Bochnak, M. Coste, M.-F. Roy, *Real Algebraic Geometry*, *Ergeb. Math. Grenzgeb.*, vol. 95, Springer, Berlin, 1998.
- [2] M.D. Choi, Positive semidefinite biquadratic forms, *Linear Algebra Appl.* 12 (1975) 95–100.
- [3] M.D. Choi, T.Y. Lam, An old question of Hilbert, in: *Proc. Conf. Quadratic Forms*, Kingston, 1976, *Queen's Papers in Pure and Appl. Math.*, vol. 46, 1977, pp. 385–405.
- [4] M.D. Choi, T.Y. Lam, Extremal positive semidefinite forms, *Math. Ann.* 231 (1) (1977) 1–18.
- [5] M.D. Choi, T.Y. Lam, B. Reznick, Symmetric quartic forms, unpublished, 1980.
- [6] C. Goel, Extension of Hilbert's 1888 theorem to even symmetric forms, Dissertation, University of Konstanz, 2014, <https://kops.uni-konstanz.de/handle/123456789/29352?locale-attribute=en>.
- [7] C. Goel, S. Kuhlmann, B. Reznick, The analogue of Hilbert's 1888 theorem for even symmetric forms, *arXiv:1509.07482*, 2015.
- [8] W.R. Harris, Real even symmetric ternary forms, *J. Algebra* 222 (1) (1999) 204–245.
- [9] D. Hilbert, Über die Darstellung definiter Formen als Summe von Formenquadraten, *Math. Ann.* 32 (1888) 342–350, *Ges. Abh.*, vol. 2, Springer, Berlin, 1933, pp. 154–161, reprinted by Chelsea, New York, 1965.
- [10] S. Kuhlmann, A. Kovacec, C. Riener, Note on extrema of linear combinations of elementary symmetric functions, *Linear Multilinear Algebra* 60 (2012) 219–224.
- [11] A. Lax, P.D. Lax, On sums of squares, *Linear Algebra Appl.* 20 (1978) 71–75.
- [12] T.S. Motzkin, The arithmetic-geometric inequality, in: Oved Shisha (Ed.), *Inequalities*, Academic Press, 1967, pp. 205–224.
- [13] B. Reznick, Forms derived from the arithmetic-geometric inequality, *Math. Ann.* 283 (1989) 431–464.
- [14] R.M. Robinson, Some definite polynomials which are not sums of squares of real polynomials; selected questions of algebra and logic, *Acad. Sci. USSR* (1973) 264–282, *Abstract in Notices Amer. Math. Soc.* 16 (1969) 554.
- [15] K. Schmüdgen, An example of a positive polynomial which is not a sum of squares of polynomials. A positive, but not strongly positive functional, *Math. Nachr.* 88 (1979) 385–390.
- [16] V. Timofte, On the positivity of symmetric polynomial functions. Part I: general results, *J. Math. Anal. Appl.* 284 (2003) 174–190.