

# SOME NEW CANONICAL FORMS FOR POLYNOMIALS

BRUCE REZNICK

ABSTRACT. We give some new canonical representations for forms over  $\mathbb{C}$ . For example, a general binary quartic form can be written as the square of a quadratic form plus the fourth power of a linear form. A general cubic form in  $(x_1, \dots, x_n)$  can be written uniquely as a sum of the cubes of linear forms  $\ell_{ij}(x_i, \dots, x_j)$ ,  $1 \leq i \leq j \leq n$ . A general ternary quartic form is the sum of the square of a quadratic form and three fourth powers of linear forms. The methods are classical and elementary.

## 1. INTRODUCTION AND OVERVIEW

**1.1. Introduction.** Let  $H_d(\mathbb{C}^n)$  denote the  $N(n, d) = \binom{n+d-1}{d}$ -dimensional vector space of complex forms of degree  $d$  in  $n$  variables, or  $n$ -ary  $d$ -ic forms. One of the major accomplishments of 19th century algebra was the discovery of canonical forms for certain classes of  $n$ -ary  $d$ -ics, especially as the sum of  $d$ -th power of linear forms. By a *canonical form* we mean a polynomial  $F(t; x)$  in two sets of variables,  $t \in \mathbb{C}^{N(n,d)}$  and  $x \in \mathbb{C}^n$ , with the property that for general  $p \in H_d(\mathbb{C}^n)$ , there exists  $t$  so that  $p(x) = F(t; x)$ . Put another way, the set  $\{F(t; x) : t \in \mathbb{C}^{N(n,d)}\}$  is a Zariski open set in  $H_d(\mathbb{C}^n)$ .

In this paper, we present some new canonical forms, whose main novelty is that they involve intermediate powers of forms of higher degree, or forms with a restricted set of monomials. (These variations have been suggested by Hilbert's study of ternary quartics [16], which led to his 17th problem, as well as by a remarkable theorem of B. Reichstein [31] on cubic forms.) These expressions, are less susceptible to apolarity arguments than the traditional canonical forms, and lead naturally to (mostly open) enumeration questions.

To take a simple, yet familiar example,

$$(1.1) \quad F(t_1, t_2, t_3; x, y) = (t_1x + t_2y)^2 + (t_3y)^2$$

is a canonical form for binary quadratic forms. By the usual completion of squares,  $p(x, y) = ax^2 + 2bxy + cy^2$  can be put into (1.1) for  $t_1 = \sqrt{a}$ ,  $t_2 = b/t_1$  and  $t_3^2 = c - t_2^2$ . Many of the examples in this paper can be viewed as imperfect attempts to generalize (1.1).

In 1851, Sylvester [39, 40] presented a family of canonical forms for binary forms in all degrees.

---

*Date:* June 17, 2013.

2000 *Mathematics Subject Classification.* Primary: 11E76, 14N15; Secondary: 11E25, 11P05, 15A72.

**Theorem 1.1** (Sylvester’s Theorem).

(i) A general binary form  $p$  of odd degree  $2s - 1$  can be written as

$$(1.2) \quad p(x, y) = \sum_{j=1}^s (\alpha_j x + \beta_j y)^{2s-1}.$$

(ii) A general binary form  $p$  of even degree  $2s$  can be written as

$$(1.3) \quad p(x, y) = \lambda x^{2s} + \sum_{j=1}^s (\alpha_j x + \beta_j y)^{2s}.$$

for some  $\lambda \in \mathbb{C}$ .

The somewhat unsatisfactory nature of the asymmetric summand in (1.3) has been the inspiration for other canonical forms for binary forms of even degree.

Another familiar canonical form is the generalization of (1.1) into the upper-triangular expression for quadratic forms, found by repeated completion of the square:

**Theorem 1.2.** A general quadratic form  $p \in H_2(\mathbb{C}^n)$  can be written as:

$$(1.4) \quad p(x_1, \dots, x_n) = \sum_{k=1}^n (t_{k,k}x_k + t_{k,k+1}x_{k+1} + \dots + t_{k,n}x_n)^2, \quad t_{k,\ell} \in \mathbb{C}.$$

The expression in (1.4) is unique, up to the signs of the linear forms.

There are two ways to verify that a candidate expression  $F(t; x)$  is, in fact, a canonical form. One is the classical non-constructive method based on the existence of a point at which the Jacobian matrix has full rank. (See Corollary 2.3, and see Theorem 3.2 for the apolar version.) Lasker [24] attributes the underlying idea to Kronecker and Lüroth – see [46, p.208].

Ideally, however, a canonical form can be derived constructively, and the number of different representations can thereby be determined. The convention in this paper will be that two representations are the same if they are equal, up to a permutation of like summands and with the identification of  $f^k$  and  $(\zeta f)^k$  when  $\zeta^k = 1$ . The representation in (1.2) is unique in this sense, even though there are  $s! \cdot (2s - 1)^s$  different  $2s$ -tuples  $(\alpha_1, \beta_1, \dots, \alpha_s, \beta_s)$  for which (1.2) is valid.

In addition to Theorem 1.1, another motivational example for this paper is a remarkable canonical form for cubic forms found by Reichstein [31] in 1987, which can be thought of as a “completion of the cube”.

**Theorem 1.3** (Reichstein). A general cubic  $p \in H_3(\mathbb{C}^n)$  can be written uniquely as

$$(1.5) \quad p(x_1, \dots, x_n) = \sum_{k=1}^n \ell_k^3(x_1, \dots, x_n) + q(x_3, \dots, x_n),$$

where  $\ell_k \in H_1(\mathbb{C}^n)$  and  $q \in H_3(\mathbb{C}^{n-2})$ .

This is a canonical form, provided  $q$  is viewed as a  $t$ -linear combination of the monomials in  $(x_3, \dots, x_n)$ ; since  $N(n, 3) = n^2 + N(n-2, 3)$ , the constant count is right. Iteration (see (6.1)) gives  $p$  as a sum of roughly  $n^2/4$  cubes. The minimum from constant-counting, which is justified by the Alexander-Hirschowitz Theorem [1], is roughly  $n^2/6$ . We give Reichstein's constructive proof of Theorem 1.3 in section six.

Here are some representative examples of the new canonical forms in this paper.

**Theorem 1.4.** *A general cubic form  $p \in H_3(\mathbb{C}^n)$  has a unique representation*

$$(1.6) \quad p(x_1, \dots, x_n) = \sum_{1 \leq i < j \leq n} (t_{\{i,j\},i}x_i + \dots + t_{\{i,j\},j}x_j)^3,$$

where  $t_{\{i,j\},k} \in \mathbb{C}$ .

**Theorem 1.5.** *A general binary sextic  $p \in H_6(\mathbb{C}^2)$  can be written as  $p(x, y) = f^2(x, y) + g^3(x, y)$ , where  $f \in H_3(\mathbb{C}^2)$  is a cubic form and  $g \in H_2(\mathbb{C}^2)$  is a quadratic form.*

Theorem 1.4 has a constructive proof. Theorem 1.5 is in fact, a very special case of much deeper recent results of Várilly-Alvarado. (See [43], especially Theorem 1.2 and Remark 4.5, and Section 1.2 of [44].) We include it because our proof, in the next section, is very short.

Theorems 1.1 and 1.5 are both special cases of a more general class of canonical forms for  $H_d(\mathbb{C}^2)$ , which is a corollary of [8, Theorem 4.4] (see Theorem 3.4), but not worked out explicitly there.

**Theorem 1.6.** *Suppose  $d \geq 1$ ,  $\{\ell_j : 1 \leq j \leq m\}$  is a fixed set of pairwise non-proportional linear forms, and suppose  $e_k \mid d$ ,  $d > e_1 \geq \dots \geq e_r$ ,  $1 \leq k \leq r$ , and*

$$(1.7) \quad m + \sum_{k=1}^r (e_k + 1) = d + 1.$$

Then a general binary  $d$ -ic form  $p \in H_d(\mathbb{C}^2)$  can be written as

$$(1.8) \quad p(x, y) = \sum_{j=1}^m t_j \ell_j^d(x, y) + \sum_{k=1}^r f_k^{d/e_k}(x, y),$$

where  $t_j \in \mathbb{C}$  and  $\deg f_k = e_k$ .

The condition  $e_k < d$  excludes the vacuous case  $m = 0, r = 1, e_1 = d$ . If each  $e_k = 1$  and  $r = \lfloor \frac{d+1}{2} \rfloor$ , then  $m = d + 1 - 2 \lfloor \frac{d+1}{2} \rfloor \in \{0, 1\}$  and Theorem 1.6 becomes Theorem 1.1; Theorem 1.5 is Theorem 1.6 in the special case  $d = 6, m = 0, r = 2, e_1 = 3, e_2 = 2$ . As an example of a canonical form which is unlikely to find a constructive proof: for a general  $p \in H_{84}(\mathbb{C}^2)$ , there exist  $f \in H_{42}(\mathbb{C}^2), g \in H_{28}(\mathbb{C}^2)$  and  $h \in H_{12}(\mathbb{C}^2)$  so that  $p = f^2 + g^3 + h^7$ .

By taking  $d = 2s$ ,  $e_1 = 2, e_2 = \dots = e_{s-1} = 1$  and  $m = 0$ , in Theorem 1.6, we obtain an alternative to the dangling term “ $\lambda x^{2s}$ ” in (1.3).

**Corollary 1.7.** *A general binary form  $p$  of even degree  $2s$  can be written as*

$$(1.9) \quad p(x, y) = (\alpha_0 x^2 + \beta_0 xy + \gamma_0 y^2)^s + \sum_{j=1}^{s-1} (\alpha_j x + \beta_j y)^{2s}.$$

A different generalization of Theorem 1.1 focuses on the number of summands.

**Theorem 1.8.** *A general binary form of degree  $uv$  can be written as a sum of  $\lceil \frac{uv+1}{u+1} \rceil$   $v$ -th powers of binary forms of degree  $u$ .*

Cayley proved that, after an invertible linear change of variables  $(x, y) \mapsto (X, Y)$ , a general binary quartic can be written as  $X^4 + 6\lambda X^2 Y^2 + Y^4$ . There are two natural ways to generalize this to higher even degree, and almost 100 years ago, Wakeford [45, 46] did both.

**Theorem 1.9** (Wakeford's Theorem). *After an invertible linear change of variables, a general  $p \in H_d(\mathbb{C}^n)$  can be written so that the coefficient of each  $x_i^d$  is 1 and the coefficient of each  $x_i^{d-1} x_j$  is 0.*

There are  $N(n, d) - n^2$  unmentioned monomials above, and when combined with the  $n^2$  coefficients in the change of variables, the constant count is correct for a canonical form. Wakeford was also interested in knowing *which* sets of  $n(n-1)$  monomials can be eliminated by a change of variables, and we are able to settle this for binary forms in Theorem 2.4. (Theorem 1.9 was independently discovered by Guazzone [14] in 1975, as an attempt to generalize the canonical form  $X^3 + Y^3 + Z^3 + 6\lambda XYZ$  for  $H_3(\mathbb{C}^3)$ . Babbage [2] subsequently observed that this can be proved by the Lasker-Wakeford Theorem, without noting that Wakeford had already done so in [46].)

The second generalization of  $X^4 + 6\lambda X^2 Y^2 + Y^4$  will not be pursued here; see [8, Corollary 4.11]. A canonical form for binary forms of even degree  $2s$  is given by

$$(1.10) \quad \sum_{k=1}^s \ell_k^{2s}(x, y) + \lambda \prod_{k=1}^s \ell_k^2(x, y), \quad \ell_k(x, y) = \alpha_k x + \beta_k y.$$

This construction is due to Sylvester [40] for  $2s = 4, 8$ . His methods failed for  $2s = 6$ , but Wakeford was able to prove it in [45]. The full version of (1.10) is proved in [46, p.408], where Wakeford notes that “the number of ways this reduction can be performed is interesting”, citing “3,8,5” for  $2s = 4, 6, 8$ .

The non-trivial study of canonical forms was initiated by Clebsch's 1861 discovery ([5], see e.g. [12, pp.50-51] and [32, pp.59-60]) that, despite the fact that  $N(3, 4) = 5 \times N(3, 1)$ , a general ternary quartic cannot be written as a sum of five fourth powers of linear forms. This was early evidence that constant-counting can fail. But  $N(3, 4)$  is also equal to  $1 \times N(3, 2) + 3 \times N(3, 1)$ , and ternary quartics *do* satisfy an alternative canonical form as a mixed sum of powers.

**Theorem 1.10.** *A general ternary quartic  $p \in H_4(\mathbb{C}^3)$  can be written as*

$$(1.11) \quad p(x_1, x_2, x_3) = q^2(x_1, x_2, x_3) + \sum_{k=1}^3 \ell_k^4(x_1, x_2, x_3),$$

where  $q \in H_2(\mathbb{C}^3)$  and  $\ell_k \in H_1(\mathbb{C}^3)$ .

As an alternative generalization of canonical forms, one might also consider polynomial maps  $F : S \mapsto H_d(\mathbb{C}^n)$ , where  $S$  is an  $N$ -dimensional subspace of some  $\mathbb{C}^M$ . In the simplest case, for binary quadratic forms, observe that the coefficient of  $x^2$  in

$$(1.12) \quad (t_1x + t_2y)^2 + (it_1x + t_3y)^2,$$

is 0, so (1.12) is not canonical. This is essentially the only kind of exception.

**Theorem 1.11.** *Suppose  $(c_1, c_2, c_3, c_4) \in \mathbb{C}^4$ , and it is not true that  $c_3 = \epsilon c_1$  and  $c_4 = \epsilon c_2$  for  $\epsilon \in \{\pm i\}$ . Then for general  $p \in H_2(\mathbb{C}^2)$ , there exists  $(t_1, t_2, t_3, t_4) \in \mathbb{C}^4$  satisfying  $\sum_{j=1}^4 c_j t_j = 0$  and such that*

$$(1.13) \quad p(x, y) = (t_1x + t_2y)^2 + (t_3x + t_4y)^2.$$

*In the exceptional case, there exists  $(x_0, y_0)$  so that for all feasible choices of  $t_j$ ,  $p(x_0, y_0) = 0$ .*

Another alternative version of (1.3) is the following conjecture, which can be verified up to degree 8.

**Conjecture 1.12.** *A general binary form  $p$  of even degree  $2s$  can be written as*

$$(1.14) \quad p(x, y) = \sum_{j=1}^{s+1} (\alpha_j x + \beta_j y)^{2s}, \quad \text{where } \sum_{j=1}^{s+1} (\alpha_j + \beta_j) = 0.$$

**1.2. Outline.** Here is an outline of the paper. In Section 2, we introduce notation and definitions. The definition of canonical form is the classical one and roughly parallels that in Ehrenborg-Rota [8], an important updating of this subject about 20 years ago. Our point of view is considerably more elementary in many respects than [8], but uses the traditional criterion: A polynomial map  $F : \mathbb{C}^N \mapsto H_d(\mathbb{C}^n)$  is a *canonical form* if a general  $p \in H_d(\mathbb{C}^n)$  is in the range; this occurs if and only if there is at least one point  $u \in \mathbb{C}^N$  so that  $\{\frac{\partial F}{\partial t_j}(u)\}$  spans  $H_d(\mathbb{C}^n)$ . (See Corollary 2.3.) This leads to immediate non-constructive proofs of Theorems 1.2, 1.5, 1.9 and 1.10, and a somewhat more complicated proof of Theorem 2.4, which answers Wakeford's question about missing monomials for binary forms.

In Section 3, we discuss classical apolarity and its implications for canonical forms. (Apolarity methods become more complicated when a component of a canonical form comes from a restricted set of monomials.) A generalization of the classical Fundamental Theorem of Apolarity from [34] allows us to identify a class of bases for  $H_d(\mathbb{C}^n)$  which give a non-constructive proof of Theorem 1.6, and hence Theorem 1.1. A similar argument yields the proof of Theorem 1.8. We also present Sylvester's Algorithm,

Theorem 3.8, allowing for a constructive proof of Theorem 1.1. We conclude with a brief summary of connections with the theorems of Alexander-Hirschowitz and recent work on the rank of forms.

In Section 4 we discuss some special cases of Theorem 1.6. Sylvester's Algorithm is used in constructive proof of Theorem 1.6 when  $e_k \equiv 1$ , in which case the representation is unique. We give some other constructive proofs for  $d \leq 4$ , and present numerical evidence regarding the number of representations in Corollary 1.7 and a few other cases. Using elementary number theory, we show that, for each  $r$ , there are only finitely many canonical forms (1.8) with  $m = 0$ , and, up to degree  $N$ , there are  $N + \mathcal{O}(N^{1/2})$  such canonical forms in which the  $e_k$ 's are equal.

Section 5 discusses some familiar results on sums of two squares of binary forms and canonical representations of quadratic forms as a sum of squares of linear forms. This includes a constructive proof of Theorem 1.2, which provides the groundwork for the proof of Theorem 1.4. We also give a short proof of a canonical form which illustrates the classical result that a general ternary quartic is the sum of three squares of quadratic forms.

In Section 6, we turn to forms in more than two variables and low degree, give constructive proofs of Theorems 1.3 and 1.4, as well as the non-canonical Theorem 6.2, which shows that *every* cubic in  $H_3(\mathbb{C}^n)$  is a sum of at most  $\frac{n(n+1)}{2}$  cubes of linear forms. Theorem 1.3 can be "lifted" to an ungainly canonical form for quartics as a sum of fourth powers (see Corollary 6.3), but not further to quintics. Number theoretic considerations rule out a Reichstein-type canonical form for quartics in 12 variables; see Theorem 6.4 for other instances of this phenomenon.

In Section 7, we offer a preliminary discussion of canonical forms in which the domain of a polynomial map  $F : \mathbb{C}^M \mapsto H_d(\mathbb{C}^n)$  is restricted to an  $N$ -dimensional subspace of  $\mathbb{C}^M$ , of which Theorem 1.11 and Conjecture 1.12 are examples.

The greatest debt of the author is due Richard Ehrenborg and Gian-Carlo Rota for writing [8]. Thanks to Dave Anderson and Julianna Tymoczko for organizing the Special Session on Geometric Commutative Algebra and Applications at the March 2011 AMS Sectional Meeting in Iowa City, to Lek-Heng Lim for organizing the Minisymposium on Tensor Rank at the October 2011 SIAM Conference on Applied Algebraic Geometry in Raleigh and to Eugene Mukhin for organizing ALGECOM5 in Indianapolis in October 2011. Invitations to speak at these conferences provided an opportunity to present preliminary versions of this material.

I am indebted to Tony Várilly-Alvarado for extremely helpful correspondence. I also want to thank T. Y. Lam, Hal Schenck, Bernd Sturmfels and Doron Zeilberger for their assistance. Special thanks go to the Center for Advanced Study at UIUC, where the author was an Associate in the Fall 2011 semester, and thereby free of teaching responsibilities. Finally, I want to thank the referee for a careful reading of the manuscript and for making many useful suggestions.

## 2. BASIC DEFINITIONS, AND PROOFS OF THEOREMS 1.2, 1.5, 1.9 AND 1.10

Let  $\mathcal{I}(n, d)$  denote the index set of monomials in  $H_d(\mathbb{C}^n)$ :

$$(2.1) \quad \mathcal{I}(n, d) = \left\{ (i_1, \dots, i_n) : 0 \leq i_k \in \mathbb{Z}, \sum_k i_k = d \right\}.$$

Let  $x^i = x_1^{i_1} \cdots x_n^{i_n}$  and  $c(i) = \frac{d!}{\prod i_k!}$  denote the multinomial coefficient. If  $p \in H_d(\mathbb{C}^n)$ , then we write

$$(2.2) \quad p(x_1, \dots, x_n) = \sum_{i \in \mathcal{I}(n, d)} c(i) a(p; i) x^i, \quad a(p; i) \in \mathbb{C}.$$

We say that two forms are *distinct* if they are non-proportional, and a set of forms is *honest* if the forms are pairwise distinct. For later reference, recall Biermann's Theorem; see [32, p.31].

**Theorem 2.1** (Biermann's Theorem). *If  $p \in H_d(\mathbb{C}^n)$  and  $p \neq 0$ , then there exists  $i \in \mathcal{I}(n, d)$  so that  $p(i) \neq 0$ .*

The easy verification of whether a formula is a canonical form for  $H_d(\mathbb{C}^n)$  relies on a crucial alternative. A self-contained accessible proof is in [8, Theorem 2.4], for which Ehrenborg and Rota thank M. Artin and A. Mattuck. For further discussion of the underlying algebraic geometry, see Section 9.5 in Cox, Little and O'Shea [7].

**Theorem 2.2.** *Suppose  $M \geq N$  and  $F : \mathbb{C}^M \rightarrow \mathbb{C}^N$  is a polynomial map; that is,*

$$F(t_1, \dots, t_M) = (f_1(t_1, \dots, t_M), \dots, f_N(t_1, \dots, t_M))$$

where each  $f_j \in \mathbb{C}[t_1, \dots, t_M]$ . Then either (i) or (ii) holds:

(i) *The  $N$  polynomials  $\{f_j : 1 \leq j \leq N\}$  are algebraically dependent and  $F(\mathbb{C}^M)$  lies in some non-trivial variety  $\{P = 0\}$  in  $\mathbb{C}^N$ .*

(ii) *The  $N$  polynomials  $\{f_j : 1 \leq j \leq N\}$  are algebraically independent and  $F(\mathbb{C}^M)$  is dense in  $\mathbb{C}^N$ .*

*The second case occurs if and only there is a point  $u \in \mathbb{C}^M$  at which the Jacobian matrix  $\left[ \frac{\partial f_i}{\partial t_j}(u) \right]$  has full rank.*

When  $M = N = N(n, d)$ , we may interpret such an  $F$  as a map from  $\mathbb{C}^N$  to  $H_d(\mathbb{C}^n)$  by indexing  $\mathcal{I}(n, d)$  as  $\{i(k) : 1 \leq k \leq N\}$  and making the interpretation in an abuse of notation that

$$(2.3) \quad F(t; x) = \sum_{k=1}^N c(i(k)) f_k(t_1, \dots, t_N) x^{i(k)}.$$

**Definition.** A *canonical form* for  $H_d(\mathbb{C}^n)$  is any polynomial map  $F : \mathbb{C}^{N(n, d)} \mapsto H_d(\mathbb{C}^n)$  in which  $F$  satisfies Theorem 2.2(ii).

That is,  $F$  is a canonical form if and only if  $N = N(n, d)$  and for a general  $p \in H_d(\mathbb{C}^n)$ , there exists  $t \in \mathbb{C}^N$  so that  $p(x) = F(t; x)$ . The significance of this

choice of  $N$  is that it is the smallest possible value. In the rare cases where  $F$  is surjective, we say that the canonical form is *universal*.

By translating the definitions and using (2.1) and (2.3), we obtain an immediate corollary of Theorem 2.2:

**Corollary 2.3.** *The polynomial map  $F : \mathbb{C}^N \mapsto H_d(\mathbb{C}^n)$  is a canonical form if and only if there exists  $u \in \mathbb{C}^n$  so that  $\{\frac{\partial F}{\partial t_j}(u)\}$  spans  $H_d(\mathbb{C}^n)$ .*

We shall let  $J := J(F; u)$  denote the span of the forms  $\{\frac{\partial F}{\partial t_j}(u)\}$ . In any particular case, the determination of whether  $J = H_d(\mathbb{C}^n)$  amounts to the computation of the determinant of an  $N(n, d) \times N(n, d)$  matrix. As much as possible in this paper, we give proofs which can be checked by hand, by making a judicious choice of  $u$  and ordering of the monomials in  $H_d(\mathbb{C}^n)$ , showing sequentially that they all lie in  $J$ .

Classically, the use of the term “canonical form” has been limited to cases in which  $F(t; x)$  has a natural interpretation as a combination of forms in  $H_d(\mathbb{C}^n)$ , such as a sum of powers of linear forms, or as a result of a linear change of variables. It seems odd that canonical forms are perceived as rare, since a “general” polynomial map from  $\mathbb{C}^N \mapsto H_d(\mathbb{C}^n)$  is a canonical form. (This is an observation which goes back at least to [38].) For example, if  $\{f_j(x)\}$  is a basis for  $H_d(\mathbb{C}^n)$ , then

$$(2.4) \quad F(t; x) = \sum_{j=1}^N t_j f_j(x)$$

should be (but usually isn't) considered a canonical form. In particular, (2.2) with  $f_j(x) = c(i_j)x^{i_j}$  is itself a canonical form.

The following computation will occur repeatedly. If  $es = d$ , then

$$(2.5) \quad g = \sum_{i_j \in \mathcal{I}(n, e)} t_j x^{i_j} \implies \frac{\partial g^s}{\partial t_j} = s x^{i_j} g^{s-1}.$$

If  $g$  is specialized to be a monomial, then all these partials will also be monomials.

*Non-constructive proof of Theorem 1.2.* Given (1.4), let

$$\ell_k(x) = \sum_{m=k}^n t_{k,m} x_m, \quad F(x) = \sum_{k=1}^n \ell_k^2(x).$$

Then  $\frac{\partial F}{\partial t_{k,m}} = 2x_m \ell_k$ . Set  $t_{k,m} = \delta_{k,m}$ , so that  $\ell_k = x_k$  and  $\frac{\partial F}{\partial t_{k,m}} = 2x_k x_m$ . Since  $1 \leq k \leq m \leq n$ , all monomials from  $H_2(\mathbb{C}^n)$  appear in  $J$ .  $\square$

*Non-constructive proof of Theorem 1.5.* Suppose

$$(2.6) \quad \begin{aligned} p(x, y) &= f^2(x, y) + g^3(x, y) : \\ f(x, y) &= t_1 x^3 + t_2 x^2 y + t_3 x y^2 + t_4 y^3, \quad g(x, y) = t_5 x^2 + t_6 x y + t_7 y^2. \end{aligned}$$

Then by (2.5), the partials with respect to the  $t_j$ 's are:

$$2x^3 f, 2x^2 y f, 2x y^2 f, 2y^3 f; \quad 3x^2 g^2, 3x y g^2, 3y^2 g^2.$$

Upon specializing at  $f = x^3, g = y^2$ , these become:

$$2x^6, 2x^5y, 2x^4y^2, 2x^3y^3; \quad 3x^2y^4, 3xy^5, 3y^6.$$

It is then evident that  $J = H_6(\mathbb{C}^2)$ .  $\square$

*Non-constructive proof of Theorem 1.9.* Let  $\mathcal{L} \subset \mathcal{I}(n, d)$  consist of all  $n$ -tuples except the permutations of  $(d, 0, \dots, 0)$  and  $(d-1, 1, \dots, 0)$  and let  $X_i = \sum_{j=1}^n \alpha_{ij}x_j$ . The assertion is that, with the  $(N(n, d) - n - \binom{n}{2}) + n^2 = N(n, d)$  parameters  $t_\ell$  and  $\alpha_{ij}$ ,

$$(2.7) \quad \sum_{i=1}^n X_i^d + \sum_{\ell \in \mathcal{L}} t_\ell X_1^{\ell_1} \cdots X_n^{\ell_n}.$$

is a canonical form. Evaluate the partials at the point where  $X_i = x_i$  and  $t_\ell = 0$ : they are  $dx_j x_i^{d-1}$  (for  $\alpha_{ij}$ ) and  $x^\ell$  (for  $t_\ell$ ). Taking  $1 \leq i, j \leq n$  and  $\ell \in \mathcal{L}$ , we see that  $J$  contains all monomials in  $H_d(\mathbb{C}^n)$ .  $\square$

As a special case (used later in Theorem 4.6), we obtain the familiar result that after appropriate linear changes of variable, a general binary quartic may be written as  $x^4 + 6\lambda x^2 y^2 + y^4$ . It is classically known (see [9, §211]) the choice of  $\lambda$  is not unique: in fact, after appropriate linear changes of variable,  $x^4 + 6\lambda x^2 y^2 + y^4$  can be written as  $x^4 + 6\mu x^2 y^2 + y^4$  for  $\mu \in \{\pm\lambda, \pm\frac{1-\lambda}{1+3\lambda}, \pm\frac{1+\lambda}{1-3\lambda}\}$ .

Wakeford asserts that Theorem 1.9 is also true with  $x_i^{d-1}x_j$  replaced by  $x_i^{d-r}x_j^r$  (evidently when  $r \neq \frac{d}{2}$ ), but his proof seems sketchy. He also gives necessary conditions for sets of  $n(n-1)$  monomials which may be omitted, and these are hard to follow as well. Below, we answer his question in the binary case: in the only two excluded cases below, (2.8) has a square factor, and so cannot be canonical.

**Theorem 2.4.** *Let  $\mathcal{B} = (m_1, m_2, n_1, n_2)$  be four distinct integers in  $\{0, \dots, d\}$  so that  $\{m_1, m_2\} \neq \{0, 1\}, \{d-1, d\}$ . Then, after an invertible linear change of variable, a general binary form  $p$  of degree  $d$  can be written as*

$$(2.8) \quad p(x, y) = x^{d-n_1}y^{n_1} + x^{d-n_2}y^{n_2} + \sum_{k \notin \mathcal{B}} t_k x^{d-k}y^k$$

for some  $\{t_k\} \subset \mathbb{C}$ .

*Proof.* Writing  $(x, y) \mapsto (\alpha_1 x + \alpha_2 y, \alpha_3 x + \alpha_4 y) := (X, Y)$ , we have

$$(2.9) \quad F = X^{d-n_1}Y^{n_1} + X^{d-n_2}Y^{n_2} + \sum_{k \notin \mathcal{B}} t_k X^{d-k}Y^k.$$

Evaluate the partials of (2.9) at  $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (1, 0, 0, 1)$  (so  $X = x, Y = y$ ) and  $t_k = 1$  (note the difference with the previous proof, in which  $t_k = 0$ ). The  $d-3$  partials with respect to the  $t_k$ 's are simply  $x^{d-k}y^k$ ,  $k \notin \mathcal{B}$ , so these are in  $J$ . Further,

$$(2.10) \quad \frac{\partial F}{\partial \alpha_1} = \sum_{j \neq m_1, m_2} (d-j)x^{d-j}y^j, \quad \frac{\partial F}{\partial \alpha_4} = \sum_{j \neq m_1, m_2} jx^{d-j}y^j.$$

Since most monomials used in (2.10) are already in  $J$ , it follows that  $J$  also contains

$$(2.11) \quad (d - n_1)x^{d-n_1}y^{n_1} + (d - n_2)x^{d-n_2}y^{n_2}, \quad n_1x^{d-n_1}y^{n_1} + n_2x^{d-n_2}y^{n_2},$$

and since  $(d - n_1)n_2 \neq (d - n_2)n_1$ , (2.11) implies that  $x^{d-n_j}y^{n_j} \in J$  for  $j = 1, 2$ . To this point, we have shown that  $J$  contains all monomials from  $H_d(\mathbb{C}^2)$  except for  $x^{d-m_j}y^{m_j}$ , where  $m_1 < m_2$ . The two remaining partial derivatives are

$$(2.12) \quad \frac{\partial F}{\partial \alpha_2} = \sum_{j \neq m_1, m_2} (d - j)x^{d-j-1}y^{j+1}, \quad \frac{\partial F}{\partial \alpha_3} = \sum_{j \neq m_1, m_2} jx^{d-j+1}y^{j-1},$$

and so  $J$  contains as well the forms in (2.12) of the shape  $c_1x^{d-m_1}y^{m_1} + c_2x^{d-m_2}y^{m_2}$ . We need to distinguish a number of cases. If  $m_1 = 0, m_2 = d$ , then these forms are  $y^d, x^d$ . If  $m_1 = 0$  and  $2 \leq m_2 \leq d - 1$ , then these forms are  $(d - m_2)x^{d-m_2}y^{m_2}$  and  $x^d + (m_2 + 1)x^{d-m_2}y^{m_2}$ , and similarly when  $1 \leq m_1 \leq d - 2$  and  $m_2 = d$ . (Recall that we have excluded the cases  $(m_1, m_2) = (0, 1)$  and  $(d - 1, d)$ . In the remaining cases,  $1 \leq m_1 < m_2 \leq d - 1$ . If  $m_2 = m_1 + 1$ , then these forms are  $(d - (m_1 - 1))x^{d-m_1}y^{m_1}$  and  $(m_2 + 1)x^{d-m_2}y^{m_2}$ . Finally, if  $m_2 > m_1 + 1$ , then all four terms appear, and the forms are

$$(2.13) \quad \begin{aligned} &(d - m_1 + 1)x^{d-m_1}y^{m_1} + (d - m_2 + 1)x^{d-m_2}y^{m_2}, \\ &(m_1 + 1)x^{d-m_1}y^{m_1} + (m_2 + 1)x^{d-m_2}y^{m_2}. \end{aligned}$$

In each of the cases, linear combinations of the forms produce the missing monomials, so  $J = H_d(\mathbb{C}^2)$ .  $\square$

*Remark.* By writing  $p(x, y) = \prod_k (x + \alpha_k y)$ , it follows from Theorem 1.9 that, for a general set of  $d$  complex numbers  $\alpha_k$ , there exists a Möbius transformation  $T$  so that

$$(2.14) \quad \sum_{k=1}^d T(\alpha_k) = 0, \quad \sum_{k=1}^d T\left(\frac{1}{\alpha_k}\right) = 0, \quad \prod_{k=1}^d T(\alpha_k) = 1.$$

*Non-constructive proof of Theorem 1.10.* Write (1.11) as  $F(x; t)$ , where

$$\begin{aligned} q(x_1, x_2, x_3) &= t_1x_1^2 + t_2x_2^2 + t_3x_3^2 + t_4x_1x_2 + t_5x_1x_3 + t_6x_2x_3, \\ \ell_k(x_1, x_2, x_3) &= t_{k1}x_1 + t_{k2}x_2 + t_{k3}x_3. \end{aligned}$$

Evaluate the partials at:  $q = x_1x_2 + x_1x_3 + x_2x_3$  and  $(\ell_1, \ell_2, \ell_3) = (x_1, x_2, x_3)$ . Then  $\frac{\partial F}{\partial t_{k\ell}} = 4x_\ell x_k^3$ , so  $x_i^4, x_i^3x_j \in J$ ; since  $\frac{\partial F}{\partial t_1} = 2x_1^2q = 2x_1^2(x_1x_2 + x_1x_3 + x_2x_3)$ , it follows that  $x_1^2x_2x_3 \in J$ , similarly, by considering  $\frac{\partial F}{\partial t_2}$  and  $\frac{\partial F}{\partial t_3}$ , it follows that  $x_1x_2^2x_3, x_1x_2x_3^2$  are in  $J$ . Finally,  $\frac{\partial F}{\partial t_4} = 2x_1x_2q = 2x_1x_2(x_1x_2 + x_1x_3 + x_2x_3)$ , and so now  $x_1^2x_2^2 \in J$ . Similarly, by considering  $\frac{\partial F}{\partial t_5}$  and  $\frac{\partial F}{\partial t_6}$ , it follows that  $x_1^2x_3^2, x_2^2x_3^2$  are also in  $J$ , and this accounts for all monomials in  $H_4(\mathbb{C}^3)$ .  $\square$

Other applications of Corollary 2.3 to canonical forms can be found in [8], including interpretations of the older results in [38] and [42, pp.265-269].

## 3. APOLARITY AND PROOFS OF THEOREMS 1.1, 1.6 AND 1.8

Using the notation of (2.1) and (2.2), for  $p, q \in H_d(\mathbb{C}^n)$ , define the following bilinear form:

$$(3.1) \quad [p, q] = \sum_{i \in \mathcal{I}(n, d)} c(i) a(p; i) a(q; i).$$

Recall two basic notations. For  $\alpha \in \mathbb{C}^n$ , define  $(\alpha \cdot)^d \in H_d(\mathbb{C}^n)$  by

$$(3.2) \quad (\alpha \cdot)^d(x) = (\alpha \cdot x)^d = \left( \sum_{j=1}^n \alpha_j x_j \right)^d = \sum_{i \in \mathcal{I}(n, d)} c(i) \alpha^i x^i.$$

Define the differential operator  $f(D)$  for  $f \in H_e(\mathbb{C}^n)$  in the usual way by

$$(3.3) \quad f(D) = \sum_{i \in \mathcal{I}(n, e)} c(i) a(f; i) \left( \frac{\partial}{\partial x_1} \right)^{i_1} \cdots \left( \frac{\partial}{\partial x_n} \right)^{i_n}.$$

It follows immediately that for  $\alpha \in \mathbb{C}^n$ ,

$$(3.4) \quad [p, (\alpha \cdot)^d] = \sum_{i \in \mathcal{I}(n, d)} c(i) a(p; i) \alpha^i = p(\alpha).$$

If  $i \neq j \in \mathcal{I}(n, d)$ , then  $i_k > j_k$  for some  $k$ , so  $D^i x^j = 0$ ; otherwise  $D^i x^i = \prod_k (i_k)! = d! / c(i)$ . Suppose  $p, q \in H_d(\mathbb{C}^n)$ . Bilinearity and (3.3) imply the classical result that

$$(3.5) \quad \begin{aligned} p(D)q &= \sum_{i \in \mathcal{I}(n, d)} c(i) a(p; i) D^i \left( \sum_{j \in \mathcal{I}(n, d)} c(j) a(q; j) x^j \right) = \\ &= \sum_{i \in \mathcal{I}(n, d)} \sum_{j \in \mathcal{I}(n, d)} c(i) c(j) a(p; i) a(q; j) D^i x^j = \sum_{i \in \mathcal{I}(n, d)} c(i) c(i) a(p; i) a(q; i) D^i x^i \\ &= \sum_{i \in \mathcal{I}(n, d)} c(i)^2 a(p; i) a(q; i) \frac{d!}{c(i)} = d! [p, q] = d! [q, p] = q(D)p. \end{aligned}$$

**Definition.** If  $p \in H_d(\mathbb{C}^n)$  and  $q \in H_e(\mathbb{C}^n)$ , then  $p$  and  $q$  are *apolar* if  $p(D)q = q(D)p = 0$ .

Note that if  $d = e$ , then  $p$  and  $q$  are apolar if and only if  $[p, q] = 0$  and if  $d > e$ , say, then the equation  $p(D)q = 0$  is automatic, so only  $q(D)p = 0$  need be checked. By (3.4),  $p$  is apolar to  $(\alpha \cdot)^d$  if and only if  $p(\alpha) = 0$ .

The following lemma is both essential and trivial.

**Lemma 3.1.** *Suppose  $X = \text{span}(\{h_j\}) \subseteq H_d(\mathbb{C}^n)$ . Then  $X = H_d(\mathbb{C}^n)$  if and only if there is no  $0 \neq p \in H_d(\mathbb{C}^n)$  which is apolar to each of the  $h_j$ 's.*

From this point of view, Theorem 3.2 is a direct consequence of Corollary 2.3:

**Theorem 3.2** (Lasker-Wakeford). *If  $F : \mathbb{C}^N \rightarrow H_d(\mathbb{C}^n)$ , then  $F$  is a canonical form if and only if there is a point  $u$  so that there is no non-zero form  $q \in H_d(\mathbb{C}^n)$  which is apolar to all  $N$  forms  $\left\{ \frac{\partial F}{\partial t_k}(u) \right\}$ .*

The attribution “Lasker-Wakeford” (for [24, 46]) is taken from [42]: H. W. Turnbull (1885-1961) was one of the last practicing invariant theorists who had been trained in the pre-Hilbert approach, see [10, pp.231-232]. (His text [42] is a Rosetta Stone for understanding the 19th century approach to algebra in more modern terminology.) Turnbull referred to Theorem 3.2 as “paradoxical and very curious”. E. Lasker (1868-1941) received his Ph.D. under M. Noether at Göttingen in 1902. He is probably better known for being the world chess champion for 27 years (1894-1921), spanning the life of E. K. Wakeford (1894-1916). J. H. Grace, Wakeford’s professor at Oxford, edited the second half of his thesis into the article [46] and also wrote a memorial article [13] for him in 1918:

“He [EKW] was slightly wounded early in 1916, and soon after coming home was busy again with Canonical Forms.... [H]e discovered a paper of Hilbert’s which contained the very theorem he had long been in want of – first vaguely, and later quite definitely. This was in March; April found him, full of the most joyous and reverential admiration for the great German master, working away in fearful haste to finish the dissertation ... He returned to the front in June and was killed in July.... He only needed a chance, and he never got it.”

The following properties are easily established; see, e.g., [32, 34] for proofs.

**Theorem 3.3.**

(i) If  $e \leq d$  and  $f \in H_e(\mathbb{C}^n)$ ,  $g \in H_{d-e}(\mathbb{C}^n)$  and  $p \in H_d(\mathbb{C}^n)$ , then

$$(3.6) \quad d![fg, p] = (fg)(D)p = f(D)g(D)p = e![f, g(D)p].$$

Thus,  $p$  is apolar to every multiple of  $g$  in  $H_d(\mathbb{C}^n)$  if and only if  $p$  and  $g$  are apolar. (ii) If  $p \in H_d(\mathbb{C}^n)$ , then  $\frac{1}{d} \frac{\partial p}{\partial x_j}(\alpha) = [p, x_j(\alpha)^{d-1}]$ . Thus,  $p$  is apolar to  $(\alpha \cdot)^{d-1}$  if and only if  $p$  is singular at  $\alpha$ . More generally,  $p$  is apolar to  $(\alpha \cdot)^{d-e}$  if and only if  $p$  vanishes to  $e$ -th order at  $\alpha$ .

(iii) If  $e \leq d$  and  $g \in H_{d-e}(\mathbb{C}^n)$ , then  $g(D)(\alpha \cdot)^d = \frac{d!}{e!} g(\alpha)(\alpha \cdot)^e$ .

Suppose  $F(t; x)$  contains  $h^s$  as a summand, where  $h(x) = \sum_{\ell \in \mathcal{I}(n, e)} t_\ell x^\ell$ , and suppose that no  $t_\ell$  occurs elsewhere in  $F(t; x)$ . If  $p$  is apolar to each partial of  $F$ , then it will be apolar to  $\frac{\partial F}{\partial t_\ell} = s x^\ell h^{s-1}$  by (2.5). Since this is true for every  $\ell \in \mathcal{I}(n, e)$ , it follows from (i) that  $p$  is apolar to  $h^{s-1}$ . It is critical to note that this observation requires that each of the monomials of degree  $e$  appear in  $h$ , and does not apply if  $h$  is defined as a sum from a restricted set of monomials.

We are now able to give a short proof of the “Second main theorem on apolarity” from [8], which was not concerned with preserving the constant-count.

**Theorem 3.4.** Suppose  $j_\ell = (j_{\ell,1}, \dots, j_{\ell,m})$ ,  $1 \leq \ell \leq r$ , are  $m$ -tuples of non-negative integers, and suppose positive integers  $d_k$ ,  $1 \leq k \leq m$ , and  $d$  are chosen so that

$$(3.7) \quad u_\ell := d - \sum_{k=1}^m j_{\ell,k} d_k \geq 0$$

for each  $\ell$ . Fix forms  $q_\ell \in H_{u_\ell}(\mathbb{C}^n)$  and for  $f_k \in H_{d_k}(\mathbb{C}^n)$ , define

$$(3.8) \quad F(f_1, \dots, f_m) = \sum_{\ell=1}^r q_\ell(x) f_1^{j_{\ell,1}} \cdots f_m^{j_{\ell,m}}.$$

Let  $F_j := \frac{\partial F}{\partial f_j}$ . Then a general  $p \in H_d(\mathbb{C}^n)$  can be written as (3.8) if and only if there exists a specific  $\bar{f} = (\bar{f}_k)$  so that no non-zero  $p \in H_d(\mathbb{C}^n)$  is apolar to each  $F_j(\bar{f})$ ,  $1 \leq j \leq m$ . If, in addition,

$$(3.9) \quad \sum_{k=1}^m N(n, d_k) = N(n, d),$$

then (3.8) is a canonical form.

*Proof.* Let

$$(3.10) \quad f_j(x) = \sum_{i_v \in \mathcal{I}(n, d_j)} t_{j,v} x^{i_v}.$$

By Theorem 2.2, (3.7) and Lemma 3.1, (3.8) represents general  $p \in H_d(\mathbb{C}^n)$  if and only if there is some  $\bar{f}$  so that there is no non-zero form in  $p \in H_d(\mathbb{C}^n)$  which is apolar to each  $\frac{\partial F}{\partial t_{j,v}}(\bar{f}) = d_k x^{i_v} F_j(\bar{f})$ , or by Theorem 3.3(i), to each  $F_j(\bar{f})$ . The constant count is checked by (3.9).  $\square$

By Theorem 3.3(ii) and Theorem 3.4,  $F = \sum_{k=1}^r (\alpha_k \cdot)^d$  is a canonical form if and only if there exist  $r$  points  $\bar{\alpha}_k \in \mathbb{C}^n$  at which no non-zero form  $p \in H_d(\mathbb{C}^n)$  is singular. This result is classical, and goes back to Clebsch [5]; see also [8, Theorem 4.2]. A particularly deep result of Alexander and Hirschowitz [1] from the early 1990s states that a general form in  $H_d(\mathbb{C}^n)$ ,  $d \geq 3$ , may be written as a sum of  $\lceil \frac{1}{n} N(n, d) \rceil$   $d$ -th powers of linear forms, except when  $(n, d) = (5, 3), (3, 4), (4, 4), (5, 4)$ , when an extra summand is needed. (For much more on this, see [12, Lecture 7], [17, Corollary 1.62], [22, Chapter 15] and [30, Theorem 0.2]; for a brief exposition of the proof, see [22, Chapter 15].) These references also discuss the exceptional examples, which were all known in the 19th century. The expression of forms as a sum of powers of forms is currently a very active area of interest; see the references above as well as [3], [11] and [23].

The Fundamental Theorem of Apolarity (see [34] for a history) states that if  $f$  is irreducible and  $p \in H_d(\mathbb{C}^n)$ , then  $f$  and  $p$  are apolar if and only if  $p$  can be written as a sum of terms of the form  $(\alpha_j \cdot)^d$ , where  $f(\alpha_j) = 0$ . This was generalized in [34].

**Theorem 3.5.** [34, Theorem 4.1] *Suppose  $q \in H_e(\mathbb{C}^n)$  factors as  $\prod_{j=1}^r q_j^{m_j}$  into a product of powers of distinct irreducible factors and suppose  $p \in H_d(\mathbb{C}^n)$ . Then  $q(D)p = 0$  if and only if there exist  $\alpha_{jk} \subset \{q_j(\alpha) = 0\}$ , and  $\phi_{jk} \in H_{m_j-1}(\mathbb{C}^n)$  such that*

$$p = \sum_{j=1}^r \left( \sum_{k=1}^{n_j} \phi_{jk}(\alpha_{jk} \cdot)^{d-(m_j-1)} \right).$$

The application of apolarity to binary forms is particularly simple, because zeros correspond to factors. If  $e = d + 1$ , then  $q(D)p = 0$  for every  $p \in H_d(\mathbb{C}^n)$ , and we obtain the following result, also found in [8, Theorem 4.5].

**Corollary 3.6.** *Suppose  $\{\alpha_j x + \beta_j y : 1 \leq j \leq r\}$  is honest and suppose  $\sum_{j=1}^r m_j = d + 1$ . Then the following set is a basis for  $H_d(\mathbb{C}^2)$ :*

$$(3.11) \quad \mathcal{S} = \{x^k y^{m_j-1-k} (\beta_j x - \alpha_j y)^{d-m_j+1} : 0 \leq k \leq m_j - 1, \quad 1 \leq m_j \leq r\}.$$

*Proof.* If  $p$  is apolar to each term in (3.11), then  $(\alpha_j x + \beta_j y)^{m_j} \mid p$  by Theorem 3.3(ii). Thus  $p = 0$  by degree considerations, and  $\mathcal{S}$  has  $d + 1$  elements, so it is a basis.  $\square$

If each  $m_j = 1$ , then Corollary 3.6 states that an honest set  $\mathcal{S} = \{(\alpha_j x + \beta_j y)^d\}$  of  $d+1$  forms is a basis for  $H_d(\mathbb{C}^2)$ . This is easily proved directly, since the representation of  $\mathcal{S}$  with respect to the basis  $\{\binom{d}{j} x^{d-j} y^j\}$ ,  $[\alpha_j^{d-k} \beta_j^k]$ , has Vandermonde determinant

$$(3.12) \quad \prod_{1 \leq i < j \leq n} (\alpha_i \beta_j - \alpha_j \beta_i).$$

Each product in (3.12) is non-zero because  $\{(\alpha_j x + \beta_j y)^d\}$  is honest. One implication of this independence is found in [36, Corollary 4.3].

**Lemma 3.7.** *If  $p(x, y) \in H_d(\mathbb{C}^2)$  has two honest representations*

$$(3.13) \quad p(x, y) = \sum_{i=1}^m (\alpha_i x + \beta_i y)^d = \sum_{j=1}^n (\gamma_j x + \delta_j y)^d$$

*and  $m + n \leq d + 1$ , then the representations are permutations of each other.*

*Proof.* If (3.13) holds, then  $\{(\alpha_i x + \beta_i y)^d, (\gamma_j x + \delta_j y)^d\}$  is linearly dependent, which is impossible unless the dependence is trivial.  $\square$

It follows immediately from Lemma 3.7 that the representations (1.2) and (1.3), if they exist for  $p$ , are unique. When  $n \geq 3$ , the linear dependence of a set  $\{(\alpha_j \cdot)^d\}$  depends on the geometry of the points as well as the number (see the discussion of Serret's Theorem in [32, p.29].) Even for powers of binary forms of degree  $e \geq 2$ , there are singular cases. It is not hard to show that a *general* set of  $(2k + 1)$   $k$ -th powers of quadratic forms is linearly independent; however, for example,  $(x^2 - y^2)^2 + (2xy)^2 = (x^2 + y^2)^2$ . For much more on this, see [37].

*Non-constructive proof of Theorem 1.6.* For  $1 \leq k \leq r$ , write

$$f_k(x, y) = \sum_{\ell=0}^{e_k} t_{k,\ell} x^{e_k-\ell} y^\ell.$$

By Corollary 2.3 and (2.5), (1.8) is a canonical form in the variables  $\{t_j, t_{k,\ell}\}$  provided there is a point at which the partials

$$\{\ell_j^d, 1 \leq j \leq m\} \cup \{x^{e_k-\ell} y^\ell f_k^{d/e_k-1}, 1 \leq \ell \leq e_k, \quad 1 \leq k \leq r\}$$

span  $H_d(\mathbb{C}^2)$ . Let  $f_k = \tilde{\ell}_k^{e_k}$ , where  $\{\ell_1, \dots, \ell_m, \tilde{\ell}_1, \dots, \tilde{\ell}_r\}$  is chosen to be honest. Then by (1.7), the desired assertion follows immediately from Corollary 3.6.  $\square$

*Non-constructive proof of Theorem 1.8.* Write  $uv + 1 = r(u + 1) + s$ . If  $s = 0$ , then Theorem 1.8 is simply a special case of Theorem 1.6 with  $m = 0$ ,  $d = uv$  and  $e_k \equiv u$ . Otherwise,  $1 \leq s \leq u$ , so that  $r + 1 = \lceil \frac{uv+1}{u+1} \rceil$ . Let

$$F(\{\alpha_{ij}\}) = \sum_{i=1}^{r+1} f_i^v(x, y), \quad f_i(x, y) = \sum_{j=0}^u \alpha_{ij} x^{u-j} y^j.$$

This is *not* a canonical form, as there are too many constants. As before,  $\frac{\partial F}{\partial \alpha_{ij}} = vx^{u-j} y^j f_i^{v-1}$ . We now specialize to  $f_i(x, y) = (ix - y)^u$  and use the apolarity argument to show that  $J = H_{uv}(\mathbb{C}^2)$ . Suppose  $q \in H_{uv}(\mathbb{C}^2)$  is apolar to each partial. Then by Theorem 3.3, it is apolar to  $f_i^{v-1} = (ix - y)^{uv-u}$ , and so  $q$  vanishes to  $u$ -th order at  $(i, -1)$  for  $1 \leq i \leq r + 1$ . It follows that  $q$  is a multiple of  $\prod_{i=1}^{r+1} (x + iy)^{u+1}$ , and so  $q = 0$  by degree considerations.

It is an exercise to show that  $F$  can be converted to an canonical form by requiring, say, that  $f_{r+1}$  only contain monomials  $x^{u-j} y^j$  for  $0 \leq j \leq s - 1$ .  $\square$

We present now Sylvester's Algorithm. For modern discussions of this, along with Gundelfinger's generalization [15], which is not included here, see [21, §5], [18],[19], [20], [34] and [36].

**Theorem 3.8** (Sylvester's Algorithm). *Let*

$$p(x, y) = \sum_{j=0}^d \binom{d}{j} a_j x^{d-j} y^j$$

*be a given binary form and suppose  $\{\alpha_j x + \beta_j y\}$  is honest. Let*

$$h(x, y) = \sum_{t=0}^r c_t x^{r-t} y^t = \prod_{j=1}^r (\beta_j x - \alpha_j y).$$

*Then there exist  $\lambda_k \in \mathbb{C}$  so that*

$$p(x, y) = \sum_{k=1}^r \lambda_k (\alpha_k x + \beta_k y)^d$$

*if and only if*

$$(3.14) \quad \begin{pmatrix} a_0 & a_1 & \cdots & a_r \\ a_1 & a_2 & \cdots & a_{r+1} \\ \vdots & \vdots & \ddots & \vdots \\ a_{d-r} & a_{d-r+1} & \cdots & a_d \end{pmatrix} \cdot \begin{pmatrix} c_0 \\ c_1 \\ \vdots \\ c_r \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Theorem 3.8 can be put in the context of our previous discussion. Let  $A_r(p)$  denote the  $(d - r + 1) \times (r + 1)$  Hankel matrix on the left-hand side of (3.14). If  $h(D) = \prod_{j=1}^r (\beta_j \frac{\partial}{\partial x} - \alpha_j \frac{\partial}{\partial y})$ , then a direct computation shows that

$$(3.15) \quad h(D)p = \sum_{m=0}^{d-r} \frac{d!}{(d-r-m)!m!} \left( \sum_{i=0}^{d-r} a_{i+m} c_i \right) x^{d-r-m} y^m.$$

It follows from (3.15) that the coefficients of  $h(D)p$  are thus, up to multiple, the rows of the matrix product, so (3.14) is equivalent to  $h(D)p = 0$ . In this way, Theorem 3.8 follows from Theorem 3.5. Sylvester's algorithm can also be visualized as seeking constant-coefficient linear recurrences satisfied by  $\{a_k\}$  and looking for the shortest one whose characteristic equation has distinct roots; this is the proof given in [36]. In this case, Gundelfinger's results handle the case when the roots are not distinct.

*Constructive proof of Theorem 1.1.* Suppose  $d = 2s - 1$  is odd. The matrix  $A_s(p)$  is  $s \times (s + 1)$  and has a non-trivial null-vector. The corresponding  $h$  (which can be given in terms of the coefficients of  $p$ ) has distinct factors unless its discriminant vanishes. Thus for general  $p \in H_{2s-1}(\mathbb{C}^2)$ , Theorem 3.8 gives  $p$  as a sum of  $s$   $(2s - 1)$ -st powers of linear forms.

If  $d = 2s$ , the matrix  $A_s(p)$  is square, and if  $p$  is a sum of  $s$   $2s$ -th powers, then  $\det A_s(p) = 0$ . Conversely, if  $\det A_s(p) = 0$  and the corresponding  $h$  has distinct factors (which is generally true), then  $p$  is a sum of  $s$   $2s$ -th powers. If  $M_1$  and  $M_2$  are two square matrices and  $\text{rank}(M_2) = k$ , then  $\det(M_1 + \lambda M_2)$  is a polynomial in  $\lambda$  of degree  $k$ . In particular, if  $q = (\alpha x + \beta y)^{2s}$ , then  $\text{rank}(H_s(q)) = 1$ . Thus, in general, there is a unique value of  $\lambda$  and some matrix  $M$  so that  $0 = \det A_s(p - \lambda(\alpha x + \beta y)^{2s}) = \det A_s(p) - \lambda \det M$ . (When  $\alpha x + \beta y = x$ ,  $M$  is the  $(1,1)$ -cofactor of  $A_s(p)$ .) In the special case  $\alpha x + \beta y = x$ , this proves Theorem 1.1(ii). The same argument shows that for general  $q \in H_{2s}(\mathbb{C}^2)$ , there exist  $s + 1$  values of  $\lambda$  so that  $p - \lambda q$  is a sum of  $s$   $2s$ -th powers.  $\square$

In 1869, Sylvester [41] recalled his discovery of this algorithm and its consequences.

“I discovered and developed the whole theory of canonical binary forms for odd degrees, and, as far as yet made out, for even degrees too, at one evening sitting, with a decanter of port wine to sustain nature's flagging energies, in a back office in Lincoln's Inn Fields. The work was done, and well done, but at the usual cost of racking thought — a brain on fire, and feet feeling, or feelingless, as if plunged in an ice-pail. *That night we slept no more.*”

*Example 3.1.* This example of Sylvester's algorithm will be used in Example 4.1. Let  $p(x, y) = 2x^3 + 3x^2y - 21xy^2 - 41y^3 = \binom{3}{0} \cdot 2 x^3 + \binom{3}{1} \cdot 1 x^2y + \binom{3}{2} \cdot (-7) xy^2 + \binom{3}{3} \cdot (-41) y^3$

Since

$$\begin{pmatrix} 2 & 1 & -7 \\ 1 & -7 & -41 \end{pmatrix} \cdot \begin{pmatrix} 6 \\ -5 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix},$$

we have  $h(x, y) = 6x^2 - 5xy + y^2 = (2x - y)(3x - y)$ . It now follows that  $p(x, y) = \lambda_1(x + 2y)^3 + \lambda_2(x + 3y)^3$ , and a simple computation shows that  $\lambda_1 = 5, \lambda_2 = -3$ .

Lemma 3.1, when applied to Theorem 2.1, yields the following corollary.

**Corollary 3.9.** *A basis for  $H_d(\mathbb{C}^n)$  is given by  $\{(i \cdot)^d : i \in \mathcal{I}(n, d)\}$ .*

This in turn gives a very weak version of the Alexander-Hirschowitz Theorem,

**Corollary 3.10.** *A general form in  $H_d(\mathbb{C}^n)$  is a sum of  $N(n, d-1) = \frac{nd}{n+d-1} \cdot \frac{1}{n} N(n, d)$   $d$ -th powers of linear forms.*

*Proof.* Consider the sum

$$\sum_{\ell=1}^{N(n, d-1)} (t_{\ell, 1}x_1 + \cdots + t_{\ell, n}x_n)^d,$$

and apply Corollary 2.3 with  $t_\ell$  specialized to  $i_\ell \in \mathcal{I}(n, d-1)$ . Then  $J$  contains  $x_k(i_\ell \cdot)^{d-1}$  for each  $k, \ell$  and hence  $x_k H_{d-1}(\mathbb{C}^n) \subseteq J$  for each  $k$ , so  $J = H_d(\mathbb{C}^n)$ .  $\square$

#### 4. EXAMPLES OF BINARY CANONICAL FORMS AND THE PROOF THEOREM 1.7

This section is devoted to special cases of Theorem 1.6. First, in the special case  $e_k = 1$ , we give a constructive proof showing uniqueness, which gives a kind of interpolation between Sylvester's Theorem and the representations of  $H_d(\mathbb{C}^2)$  by (2.4) with a fixed basis consisting of  $d$ -th powers, as in Corollary 3.6.

**Corollary 4.1.** *Suppose  $d \geq 1$ , and  $\{\ell_j(x, y) = \alpha_j x + \beta_j y\}$  is a fixed honest set of  $m = d + 1 - 2r$  linear forms. Then a general binary  $d$ -ic form  $p \in H_d(\mathbb{C}^2)$  can be written uniquely as*

$$(4.1) \quad p(x, y) = \sum_{j=1}^m t_j \ell_j(x, y)^d + \sum_{k=1}^r (t_{k1}x + t_{k2}y)^d.$$

for suitable  $t_{k1}, t_{k2} \in \mathbb{C}$ .

*Proof.* Let

$$f(x, y) = \prod_{j=1}^m (\beta_j x - \alpha_j y).$$

Then  $f(D)p$  has degree  $d - m = 2r - 1$  and by Theorem 3.8 generally has a unique representation as a sum of  $r$   $2r - 1$ -st powers of linear forms, say

$$(4.2) \quad f(D)p = \sum_{k=1}^r (u_{k1}x + u_{k2}y)^{2r-1}.$$

Further, it is generally true that  $f(u_{k1}, u_{k2}) \neq 0$ . Let

$$(4.3) \quad q(x, y) = \frac{(2r-1)!}{d!} \sum_{k=1}^r \frac{(u_{k1}x + u_{k2}y)^d}{f(u_{k1}, u_{k2})}.$$

It follows from Theorem 3.3(iii), (4.2) and (4.3) that  $f(D)p = f(D)q$ . Since  $f$  has distinct factors, it then follows from Theorem 3.8 that there exist  $t_j \in \mathbb{C}$  so that

$$p(x, y) - q(x, y) = \sum_{j=1}^m t_j (\alpha_j x + \beta_j y)^d.$$

Conversely, suppose  $p$  has two different representations:

$$(4.4) \quad \sum_{j=1}^m t_j \ell_j^d(x, y) + \sum_{k=1}^r (t_{k1}x + t_{k2}y)^d = \sum_{j=1}^m \tilde{t}_j \ell_j^d(x, y) + \sum_{k=1}^r (\tilde{t}_{k1}x + \tilde{t}_{k2}y)^d.$$

By combining the first sum on each side, (4.4) becomes a linear dependence with  $m + 2r = d + 1$  summands, which by Lemma 3.7 must be trivial; thus, the representations in (4.4) are essentially the same.  $\square$

*Example 4.1.* Let  $\ell_1(x, y) = x + y$  and  $\ell_2(x, y) = -x + 3y$  and let

$$p(x, y) = -x^5 + 15x^4y - 170x^3y^2 + 390x^2y^3 - 505x^2y^3 + 483y^5.$$

In an application of the last proof,  $f(x, y) = (x - y)(3x + y) = 3x^2 - 2xy - y^2$ , and

$$3 \frac{\partial^2 p}{\partial x^2} - 2 \frac{\partial^2 p}{\partial x \partial y} - \frac{\partial^2 p}{\partial y^2} = 160x^3 + 240x^2y - 1680xy^2 - 3280y^3.$$

Example 3.1 implies that this expression equals  $400(x + 2y)^3 - 240(x + 3y)^3$ . Since  $f(1, 2) = -5$  and  $f(1, 3) = -12$ , it follows that

$$\begin{aligned} p(x, y) = & \\ & \frac{3! \cdot 400}{5! \cdot (-5)} (x + 2y)^5 + \frac{3! \cdot (-240)}{5! \cdot (-12)} (x + 3y)^5 + t_1 (x + y)^5 + t_2 (-x + 3y)^5 = \\ & -4(x + 2y)^5 + (x + 3y)^5 + t_1 (x + y)^5 + t_2 (-x + 3y)^5 \end{aligned}$$

and it can be readily be computed that  $t_1 = \frac{7}{2}$  and  $t_2 = \frac{3}{2}$ .

If each  $e_k = 2$  in Theorem 1.6 and  $m$  is as small as possible, then we obtain an analogue of Sylvester's Theorem for forms of even degree.

**Corollary 4.2.**

(i) *A general binary form of degree  $d = 6s$  can be written as*

$$(4.5) \quad \lambda x^{6s} + \sum_{j=1}^{2s} (\alpha_j x^2 + \beta_j xy + \gamma_j y^2)^{3s}$$

for some  $\lambda \in \mathbb{C}$ .

(ii) A general binary form of degree  $d = 6s + 2$  can be written as

$$(4.6) \quad \sum_{j=1}^{2s+1} (\alpha_j x^2 + \beta_j xy + \gamma_j y^2)^{3s+1}.$$

(iii) A general binary form of degree  $d = 6s + 4$  can be written as

$$(4.7) \quad \lambda_1 x^{6s+4} + \lambda_2 y^{6s+4} + \sum_{j=1}^{2s+1} (\alpha_j x^2 + \beta_j xy + \gamma_j y^2)^{3s+2}$$

for some  $\lambda_i \in \mathbb{C}$ .

We have not been able to find an analogue to Sylvester's Algorithm for determining the representations (4.5), (4.6), (4.7) in Corollary 4.2. In the linear case,  $(\alpha x + \beta y)^d$  is killed by  $\beta \frac{\partial}{\partial x} - \alpha \frac{\partial}{\partial y}$ , and two operators of this shape commute. Although each  $(\alpha x^2 + 2\beta xy + \gamma y^2)^d$  is killed by the non-constant-coefficient  $(\beta x + \gamma y) \frac{\partial}{\partial x} - (\alpha x + \beta y) \frac{\partial}{\partial y}$ , two operators of this kind do not usually commute. The smallest constant-coefficient differential operator which kills  $(\alpha x^2 + 2\beta xy + \gamma y^2)^d$  has degree  $d + 1$ ; the product of any two of these would kill every form of degree  $2d$  and so provide no information.

Let us say that (1.8) is a *neat* canonical form if  $m = 0$ , and of *Sylvester-type* if it is neat and if  $e_k = e$  for  $1 \leq k \leq r$ . Counting the numbers of neat and Sylvester-type canonical forms leads to some number theory. The first lemma is standard.

**Lemma 4.3.** *Given  $0 < \frac{p}{q} \in \mathbb{Q}$  and  $0 < n \in \mathbb{N}$ , there exist only finitely many choices of  $m_j \in \mathbb{Z}$ ,  $0 < m_1 \leq m_2 \leq \dots \leq m_n$ , such that  $\frac{p}{q} = \sum_{j=1}^n \frac{1}{m_j}$ .*

*Proof.* If  $n = 2$ , then  $\frac{p}{q} > \frac{1}{m_1} \geq \frac{p}{2q}$  implies that there are finitely many integral choices for  $m_1$ , each of which determines  $m_2 = (\frac{p}{q} - \frac{1}{m_1})^{-1}$ . Supposing the lemma valid for  $n - 1$ , we have  $\frac{p}{q} > \frac{1}{m_1} \geq \frac{p}{nq}$ , and each choice of  $m_1$  implies the equation  $\frac{p}{q} - \frac{1}{m_1} = \sum_{j=2}^n \frac{1}{m_j}$ . This has finitely many solutions by the induction hypothesis.  $\square$

**Theorem 4.4.** *For fixed value of  $r$ , there are only finitely many neat canonical forms (1.8) with  $r$  summands.*

*Proof.* Suppose  $m = 0$  in Theorem 1.6. Write  $d = e_k m_k$ , then by (1.7),

$$(4.8) \quad d + 1 = \sum_{k=1}^r \left( \frac{d}{m_k} + 1 \right) \implies 1 = \sum_{k=1}^r \frac{1}{m_k} + \frac{r-1}{d} = \sum_{k=1}^r \frac{1}{m_k} + \sum_{\ell=1}^{r-1} \frac{1}{d}.$$

Now apply Lemma 4.3 with  $\frac{p}{q} = 1$  and  $n = 2r - 1$ : there are only finitely many expressions of 1 as a sum of  $2r - 1$  unit fractions, of which only a subset satisfy the additional restrictions of (4.8).  $\square$

It is not hard to work out that for  $r = 2$ , there are three neat canonical forms:  $(d, e_1, e_2) = (3, 1, 1)$ ,  $(4, 2, 1)$  and  $(6, 3, 2)$ . The first is Theorem 1.1(i) with  $d = 3$ , the

second is Corollary 1.7 with  $d = 4$  (see Theorem 4.6 below), and the third is Theorem 1.5. When  $r = 3$ , there are twenty-two neat canonical forms.

Let  $s(d)$  denote the number of neat Sylvester-type canonical forms of degree  $d$ . Suppose  $e_k = e$  for all  $k$  in one of these. Then  $e \mid d$  and, by (1.7),  $r(e+1) = d+1$ , so  $(e+1) \mid (d+1)$ . Since  $d \equiv 0 \pmod{e}$  and  $d \equiv -1 \pmod{e+1}$ , it follows from the Chinese Remainder Theorem that  $d \equiv e \pmod{e(e+1)}$ ; that is,  $d = e + ue(e+1)$ ,  $u \geq 1$ , so that  $e < \sqrt{d}$ .

**Theorem 4.5.** *Let  $S(N) := \sum_{d=1}^N s(d)$ . Then  $S(N) = N + \mathcal{O}(N^{1/2})$  and  $\sup_d s(d) = \infty$ .*

*Proof.* The generating function for the sequence  $(s(d))$  is

$$(4.9) \quad \sum_{n=1}^{\infty} s(d)x^d = \sum_{e=1}^{\infty} \sum_{u=1}^{\infty} x^{e+ue(e+1)} = \sum_{e=1}^{\infty} \frac{x^{e^2+2e}}{1-x^{e^2+e}} = \sum_{N=e}^{\infty} \left\lfloor \frac{N-e}{e^2+e} \right\rfloor X^N.$$

Let  $T = \lfloor N^{1/2} \rfloor$ . It follows from (4.9) that

$$(4.10) \quad S(N) = \sum_{n=1}^N s_n = \sum_{e=1}^{\infty} \left\lfloor \frac{N-e}{e^2+e} \right\rfloor = \sum_{e=1}^T \left\lfloor \frac{N-e}{e^2+e} \right\rfloor.$$

Thus, using the telescoping sum for  $\sum \frac{1}{e(e+1)}$ , (4.10) implies that

$$(4.11) \quad \begin{aligned} S(N) &\leq \sum_{e=1}^T \frac{N-e}{e^2+e} = N \sum_{e=1}^T \frac{1}{e^2+e} - \sum_{e=1}^T \frac{1}{e+1} \\ &\leq N(1 - \frac{1}{T+1}) - \log T + \mathcal{O}(1) = N - N^{1/2} + \mathcal{O}(\log N). \end{aligned}$$

The lower bound is the same, minus  $T$ , so (4.11) implies that  $S(N) = N + \mathcal{O}(N^{1/2})$ .

Now,  $s(d)$  counts the number of  $e < d$  so that  $e$  divides  $d$  and  $e+1$  divides  $d+1$ . If  $d = 2^r - 1$ , then  $e+1 \mid 2^r$  implies that  $e+1 = 2^t$  for some  $t < r$ . But  $2^t - 1 \mid 2^r - 1$  if and only if  $t \mid r$ , hence  $s(2^r - 1) = d(r) - 1$ , where  $d(n)$  denotes the divisor function. In particular,  $s(2^{2^t} - 1) = t$ , so the sequence  $(s(d))$  is unbounded. More generally, if  $e \mid d$  and  $e+1 \mid d+1$ , then  $e \mid d^2 + 2d$  and  $e+1 \mid d^2 + 2d + 1$ , and since  $e = d$  contributes to the count in  $s(d^2 + 2d)$  but not in  $s(d)$ ,  $s(d^2 + 2d) \geq s(d) + 1$ .  $\square$

Half of the neat Sylvester forms come from Theorem 1.1(i), another sixth come from Corollary 4.2(ii), etc. The smallest  $d$  for which  $s(d) = 2$  is  $d = 15$ :  $(e, r) = (1, 8), (3, 4)$ , so a general binary form of degree 15 is a sum of eight linear forms to the 15th power, or four cubics to the 5th power. Mathematica computations show that the smallest  $d$  for which  $s(d) = 3$  is  $d = 99$ :  $(e, r) = (1, 50), (3, 25), (9, 10)$ . For  $d < 10^7$ , the largest value of  $s(d)$  is  $s(7316000) = 12$ . Note that  $2^{2^{13}} - 1 = 2^{4096} - 1 \approx 1.04 \times 10^{1233}$ , so the examples given in the proof are not likely to describe the fastest growth. We conjecture as well that  $\{s(d)\}$  has an underlying distribution.

If the degree  $d$  is prime, then Theorem 4.1 accounts for all canonical forms in Theorem 1.6. The smallest  $d$  which is not covered by Theorem 4.1 is then  $d = 4$ ,

and there are two such cases, one of which is neat:  $e_1 = 2, e_2 = 1, m = 0$  and  $e_1 = 2, m = 2$ . Both can be discussed constructively.

**Theorem 4.6.** *A general binary quartic  $p \in H_4(\mathbb{C}^2)$  can be written as*

$$(4.12) \quad p(x, y) = (t_1x^2 + t_2xy + t_3y^2)^2 + (t_4x + t_5y)^4$$

*in six different ways. Further, the set of possible values for  $\{\frac{t_5}{t_4}\}$  is the image of the set  $\{0, \infty, 1, -1, i, -i\}$  under a Möbius transformation.*

*Proof.* By Theorem 2.4, if  $p$  is a general binary quartic, then there exist  $c_i, \lambda$  so that  $p(c_1x + c_2y, c_3x + c_4y) = p_\lambda(x, y) := x^4 + 6\lambda x^2y^2 + y^4$ . If (4.12) holds for  $p_\lambda$ , then

$$(4.13) \quad \begin{aligned} 1 &= t_1^2 + t_4^4, & 0 &= 2t_1t_2 + 4t_4^3t_5, & 6\lambda &= 2t_1t_3 + t_2^2 + 6t_4^2t_5^2, \\ 0 &= 2t_2t_3 + 4t_4t_5^3, & 1 &= t_3^2 + t_5^4. \end{aligned}$$

First suppose that  $t_4 = 0$ . Then (4.13) implies that  $1 = t_1^2$  and  $0 = 2t_1t_2$ , so  $t_1 = 1$  (without loss of generality) and  $t_2 = 0$ . The remaining equations imply that  $t_3 = 3\lambda$  and  $t_5^4 = 1 - 9\lambda^2$ . A similar argument works if  $t_5 = 0$ , giving two representations:

$$(4.14) \quad p_\lambda(x, y) = (x^2 + 3\lambda y^2)^2 + (1 - 9\lambda^2)y^4 = (3\lambda x^2 + y^2)^2 + (1 - 9\lambda^2)x^4.$$

Now suppose  $t_4t_5 \neq 0$ , so  $t_1t_2t_3 \neq 0$  and so

$$\frac{t_3}{t_1} = \frac{-2t_2t_3}{-2t_1t_2} = \frac{4t_4t_5^3}{4t_4^3t_5} = \frac{t_5^2}{t_4^2} \implies \frac{1 - t_3^2}{1 - t_1^2} = \frac{t_5^4}{t_4^4} = \frac{t_3^2}{t_1^2} \implies t_1^2 = t_3^2.$$

It follows that  $t_5 = i^k t_4$  and  $t_3 = (-1)^k t_1$ , and (4.13) can be completely solved:

$$t_4^4 = 1 - t_1^2, \quad t_2 = 2i^k(t_1 - t_1^{-1}), \quad 2 + 6(-1)^k\lambda = 4t_1^{-2}.$$

After some massaging of the algebra, this gives four representations:

$$(4.15) \quad \begin{aligned} p_\lambda(x, y) &= \left( \frac{(-1)^k 2}{3\lambda + (-1)^k} \right) (x^2 - i^{3k}(3\lambda - (-1)^k)xy + (-1)^k y^2)^2 \\ &+ \left( \frac{3\lambda - (-1)^k}{3\lambda + (-1)^k} \right) (x + i^k y)^4, \quad k = 0, 1, 2, 3. \end{aligned}$$

In order to find the six representations of  $p$  as (4.12), we start with the six representations of  $p_\lambda$  given in (4.14) and (4.15), in which  $t_4x + t_5y$  is a multiple of one of the six linear forms  $x, y, x + i^k y$ . Apply the the inverse of the map  $(x, y) \mapsto (c_1x + c_2y, c_3x + c_4y)$ , which takes  $t_4x + t_5y$  to a multiple of  $t_4(c_4x - c_2y) + t_5(-c_3x + c_1y)$ :  $\frac{t_5}{t_4} \mapsto G(\frac{t_5}{t_4})$ , where  $G(z) = \frac{c_1z - c_2}{c_4 - c_3z}$ .  $\square$

**Theorem 4.7.** *Given two fixed non-proportional binary linear forms  $\ell_1, \ell_2$ , a general binary quartic in  $H_4(\mathbb{C}^2)$  has two representations as*

$$(4.16) \quad p(x, y) = (t_1x^2 + t_2xy + t_3y^2)^2 + t_4\ell_1(x, y)^4 + t_5\ell_2(x, y)^4.$$

$d$	$e_1, \dots, e_r$	$m$	$F(d; e)$	Source
$d$	$1^{\lfloor \frac{d+1}{2} \rfloor}$	0 or 1	1	Theorem 1.1
$d$	$1^r$	$d + 1 - 2r$	1	Theorem 4.1
4	2,1	0	6	Theorem 4.6
4	2	2	2	Theorem 4.7
6	3,2	0	40	[43, 44]
6	$2,1^2$	0	22	Experiment
6	3,1	1	14	Experiment
6	$2^2$	1	9	Experiment
6	2,1	2	12	Experiment
6	3	3	5	Experiment
6	2	4	5	Experiment
8	$2,1^3$	0	62	Experiment
10	$2,1^4$	0	147	Experiment
12	$2,1^5$	0	308	Experiment
$2s$	$2,1^{s-1}$	0	$2\binom{s+3}{5} + \binom{s+2}{3}$	Conjecture

TABLE 1. Values of  $F(d; e)$ 

*Proof.* Given  $p, \ell_1, \ell_2$ , make an invertible linear change of variable taking  $(\ell_1, \ell_2) \mapsto (x, y)$ , and suppose  $p(x, y) \mapsto q(x, y) = \sum_i a_i x^{4-i} y^i$ . Then  $q$  has the shape (4.16) if and only if the coefficients of  $x^3y, x^2y^2, xy^3$  in  $(t_1x^2 + t_2xy + t_3y^2)^2$  and  $q$  agree. Thus, we seek to solve the system

$$(4.17) \quad a_1 = 2t_1t_2, \quad a_2 = 2t_1t_3 + t_2^2, \quad a_3 = 2t_2t_3.$$

But (4.17) implies  $a_1t_2^2 - 2a_2t_1t_2 + 2a_3t_1^2 = 0$ , hence in general, there are exactly two values of  $\beta$  so that  $t_2 = \beta t_1$ ; in each case,  $t_1^2 = \frac{a_1}{2\beta}$ . The two choices of sign for  $t_1$  lead to the same square, and  $t_3 = \frac{a_1}{a_3}t_1$ , so (4.17) has these two solutions.  $\square$

In the case of Theorem 1.6 let  $F(d; e_1, \dots, e_r)$  denote the number of different representations that a general  $p \in H_d(\mathbb{C}^2)$  has, by our convention. We present in Table 1 a complete list of proved or conjectural values when  $d \leq 6$ , reflecting numerical experiments on Mathematica. (Recall that if  $d$  is prime, then Theorem 4.1 presents all possible canonical forms of this type.) The conjectural value of  $F(2s; 2, 1^{s-1})$  is suggested by the given data for  $2 \leq s \leq 6$  and OEIS[25, A081282].

Várilly-Alvarado, in [43, 44], constructs explicitly all 240 representations of  $x^6 + y^6$  as  $f^2 + g^3$ ; he considers forms multiplied by roots of unity as different, which explains the appearance of  $\frac{240}{2 \cdot 3}$  in the table above. This is also proved to be the number of representations for a general sextic.

To describe the experiments for  $F(2s; 2, 1^{s-1})$  more precisely, we generate a form

$$p(x, y) = \sum_{k=0}^{2s} \binom{2s}{k} a_k x^{2s-k} y^k,$$

where  $a_k = t + iu$  for random integers  $t, u$  in  $[-100, 100]$ . In case  $m \leq 2$ , we assume a change of variables so that the fixed linear forms are  $x^d$  or  $y^d$ ; for  $m > 2$  we choose additional linear forms with random coefficients. Let  $h(x, y) = Ux^2 + Vxy + Wy^2$  for variables  $(U, V, W)$  and let  $q(x, y) = p(x, y) - h^s(x, y)$ , and apply Sylvester's Algorithm to  $q$ . That is, we construct the  $(s+2) \times s$  matrix  $A_{s-1}(q)$ , with polynomial entries in  $(U, V, W)$  of degree  $s$  and require that it have rank  $< s$ . This is done by counting the number of  $(U, V, W)$  which are common zeros of all  $s \times s$  minors. This number is divided by  $s$  to account for  $h^s = (\zeta_s^k h)^s$ . As a back of the envelope calculation, one might take the first  $s-1$  rows of  $A_{s-1}$  and use the cofactors to compute a non-trivial null-vector. Ignoring possible cancellation, the components would be polynomials of degree  $s(s-1)$  in  $(U, V, W)$ . Taking the dot product with the last three rows of  $A_{s-1}$  gives three polynomials of degree  $s^2$ . Ignoring cancellations and multiplicity, there should be  $(s^2)^3$  common zeros, and dividing by  $s$  gives an upper bound for  $F(2s; 2, 1^{s-1})$  of  $s^5$ . The conjectural value is asymptotically  $\frac{1}{60}s^5$ , which shows the same order of growth.

## 5. QUADRATIC FORMS AND THE PROOF OF THEOREM 1.2

We begin this section with a constructive proof of Theorem 1.2 which will serve as a template for constructive proofs involving cubic forms.

*Constructive Proof of Theorem 1.2.* Suppose  $p \in H_2(\mathbb{C}^n)$ , and specifically,

$$p(x_1, \dots, x_n) = \sum_{i=1}^n a_{ii} x_i^2 + 2 \sum_{1 \leq i < j \leq n} a_{ij} x_i x_j.$$

Then  $\frac{\partial p}{\partial x_1} = 2 \sum_{j=1}^n a_{1j} x_j$ . Since  $a_{11} \neq 0$  in general, we can define  $q(x_1, \dots, x_n) = p(x_1, \dots, x_n) - \frac{1}{a_{11}} (\sum_{j=1}^n a_{1j} x_j)^2$ . Observe that  $\frac{\partial q}{\partial x_1} = 0$ , so  $q = q(x_2, \dots, x_n)$ . Iterating this argument gives the construction. There is only one linear form  $\pm \ell$  so that  $\frac{\partial p}{\partial x_1} = 2\ell \frac{\partial \ell}{\partial x_1}$ , so the representation is unique.  $\square$

Constant-counting for sums of squares is complicated by the action of the orthogonal group on a sum of  $t$  squares. If  $M \in Mat_t(\mathbb{C})$  and  $MM^t = I$ , then

$$\sum_{i=1}^t f_i^2 = \sum_{i=1}^t \left( \sum_{j=1}^t m_{ij} f_j \right)^2.$$

When  $t = 2$ , choose  $\theta \in \mathbb{C}$  and let  $e^{i\theta} = \cos \theta + i \sin \theta := (u, v)$ , so that

$$(5.1) \quad f^2 + g^2 = (uf - vg)^2 + (vf + ug)^2.$$

This means that we may safely remove one monomial from one of the summands.

**Theorem 5.1.** *A general binary form  $p \in H_{2s}(\mathbb{C}^2)$  can be written as*

$$(5.2) \quad \left( \sum_{k=0}^s t_k x^{s-k} y^k \right)^2 + \left( \sum_{k=1}^s t_{s+k} x^{s-k} y^k \right)^2$$

in  $\binom{2s-1}{s}$  different ways.

*Proof.* The non-constructive proof is a simple application of Corollary 2.3. Writing (5.2) as  $f^2 + g^2$  gives the partials with respect to the  $t_j$ 's as

$$\{2x^{s-k}y^k f, 0 \leq k \leq s\} \cup \{2x^{s-k}y^k g, 1 \leq k \leq s\};$$

specializing to  $f = x^s$  and  $g = y^s$  above gives all monomials in  $H_{2s}(\mathbb{C}^2)$ .

The more obvious expression

$$(5.3) \quad p(x, y) = f^2(x, y) + g^2(x, y), \quad g, h \in H_s(\mathbb{C}^2)$$

is not a canonical form, because  $2(s+1) > 2s+1$ . However, every sum of two squares can be formally factored, and these behave nicely with respect to (5.1).

$$f^2 + g^2 = (f + ig)(f - ig) \iff (uf + vg)^2 + (vf - ug)^2 = (e^{i\theta}(f + ig))(e^{-i\theta}(f - ig)).$$

Suppose  $p(1, 0) = a_0 \neq 0$  (true for general  $p$ ) and (5.3) holds, where  $f(1, 0) = \rho$  and  $g(1, 0) = \tau$ . Then  $\rho^2 + \tau^2 = a_0$ , so that  $\frac{\tau}{\rho} \neq \pm i$  and the coefficient of  $x^s$  in  $vf + ug$  will be  $v\rho + u\tau = \sin\theta\rho + \cos\theta\tau$ , which is zero exactly when  $\tan\theta = -\frac{\tau}{\rho}$ . Thus for precisely one value of  $\tan\theta$ , the right-hand side of (5.1) will be in the form (5.2). This determines a pair  $(\pm u, \pm v)$ ; however, the squares in (5.2) will be the same.

In other words, each distinct factorization of  $p$  (up to multiple) as a product of two  $s$ -ic forms, when combined with the orthogonal action of (5.1), yields exactly one representation as (5.2). A general  $p \in H_{2s}(\mathbb{C}^2)$  is a product of  $2s$  distinct linear factors; these can be organized into an unordered pair of products of  $s$  distinct linear factors in  $\frac{1}{2}\binom{2s}{s} = \binom{2s-1}{s}$  ways.  $\square$

The ‘‘lost’’ degree of freedom in a sum of squares never arises in Theorem 1.6 because  $2\left(\frac{d}{2} + 1\right) > d + 1$ . The missing monomial  $x^s$  in the second summand of (5.2) may be replaced by any specified monomial  $x^{s-k_0}y^{k_0}$  by a similar argument.

Another classical result is that a general ternary quartic is a sum of three squares of quadratic forms, generally in 63 different ways up to the action of the orthogonal group (see [29].) Hilbert proved that *every* positive semidefinite  $p \in H_4(\mathbb{R}^3)$  is a sum of three squares from  $H_2(\mathbb{R}^3)$  [16]. He then showed that there exist psd forms in  $H_6(\mathbb{R}^3)$  and  $H_4(\mathbb{R}^4)$  which are not sums of squares in  $H_3(\mathbb{R}^3)$  and  $H_2(\mathbb{R}^4)$ , respectively, which ultimately led to his 17th problem. (See [31] for much more on this subject.)

A constructive discussion of Hilbert’s theorem on  $p \in H_4(\mathbb{R}^3)$  has recently been given in papers by Powers and the author [28], Powers, Scheiderer, Sottile and the author [29], Pfister and Scheiderer [26] and Plaumann, Sturmfels and Vinzant [27]. A non-constructive proof (without the count) can easily be given.

**Theorem 5.2.** *A general ternary quartic  $p \in H_4(\mathbb{C}^3)$  can be written as  $p = q_1^2 + q_2^2 + q_3^2$ , where  $q_j \in H_2(\mathbb{C}^3)$ .*

*Proof.* We take  $q_i$ 's so that the monomial  $x^2$  only appears in  $q_1$  and the monomial  $y^2$  only appears in  $q_1$  and  $q_2$ , and so the number of coefficients in the  $q_j$ 's is  $6 + 5 + 4 = 15$ . Taking the partials where  $(q_1, q_2, q_3) = (x^2, y^2, z^2)$  shows that  $J$  contains  $2x^2\{x^2, y^2, z^2, xy, xz, yz\}$ ,  $2y^2\{y^2, z^2, xy, xz, yz\}$  and  $2z^2\{z^2, xy, xz, yz\}$ , and so is equal to  $H_4(\mathbb{C}^3)$ .  $\square$

Since  $3\binom{m+2}{2} - 3 < \binom{2m+1}{2}$  for  $m \geq 3$ , this result does not generalize to ternary forms of higher even degree.

The situation is somewhat simpler over  $\mathbb{R}$ . A real version of Theorem 5.1 appears in [35]. If  $p$  is real and positive definite and  $p = f^2 + g^2$ , where  $f$  and  $g$  are also real, then the factors of  $p$  consist of  $s$  conjugate pairs. In the factorization  $p = (f + ig)(f - ig)$ , the pairs must be split between the conjugate factors, and if  $p$  has distinct factors, this can be done in  $2^{s-1}$  different ways. A real generalization of Theorem 5.2 appears in [4, Corollary 2.12]. Suppose a real psd form  $p \in H_{2s}(\mathbb{R}^n)$  is a sum of  $t$  squares and  $x^{\beta_i} \in H_s(\mathbb{R}^n)$ ,  $1 \leq i \leq t$ , is given. Then there is a representation  $p = \sum_{j=1}^t g_j^2$ , in which  $x^{\beta_i}$  does not occur in  $g_j$  for  $j > i$ . This argument can also be applied to a *general* sum of  $t$  squares over  $\mathbb{C}$ , but it no longer applies to all forms. For example, if  $xy = (ax + by)^2 + (cx + dy)^2$ , then  $abcd \neq 0$ .

## 6. CUBIC FORMS AND PROOFS OF THEOREMS 1.3 AND 1.4

In this section, we present three representations for forms in  $H_3(\mathbb{C}^n)$  as a sum of cubes of linear forms. The first two are canonical; the third isn't, but it represents *all* cubics, not just general cubics.

We begin with Theorem 1.3, which first appeared [31] in a 1987 paper of Boris Reichstein. At the time of this writing, [31] has had no citations in MathSciNet. (It was discussed in [33] and, from there, in [6]. The former was never submitted for publication and the latter appeared in an unindexed journal.) The original presentation and proof in [31] were given for trilinear forms (see §2); the theorem is applied to cubic forms there mainly in the examples.

By iterating (1.5), we obtain a canonical form for  $p \in H_3(\mathbb{C}^n)$ , see [31, p.98].

**Corollary 6.1.** *A general  $n$ -ary cubic  $p \in H_3(\mathbb{C}^n)$  can be written uniquely as*

$$(6.1) \quad p(x_1, \dots, x_n) = \sum_{m=0}^{\lfloor (n-1)/2 \rfloor} \sum_{k=1}^{n-2m} (t_{m,1+2m}^{\{k\}} x_{1+2m} + \dots + t_{m,n}^{\{k\}} x_n)^3$$

for some  $t_{m,j}^{\{k\}} \in \mathbb{C}$ .

This gives  $p$  as a sum of  $n + (n - 2) + \dots = \lfloor \frac{(n+1)^2}{4} \rfloor$  cubes. Recall that by Alexander-Hirschowitz, for  $n \neq 5$ , a general cubic form in  $n$  variables can be written as a sum of  $\lceil \frac{(n+1)(n+2)}{6} \rceil$  cubes. Thus (6.1) is a canonical form which represents a

general cubic as a sum of about 50% more cubes than the true minimum; this is due to the large number of linear forms with restricted sets of variables.

Reichstein's proof of Theorem 1.3 requires the well-known "generalized eigenvalue problem" for pairs of symmetric matrices, as interpreted for quadratic forms: if a general pair of quadratic forms  $f, g \in H_2(\mathbb{C}^n)$  is given, then there exist  $n$  linearly independent forms  $L_i(x) = \sum_{j=1}^n \alpha_{ij}x_j$  and  $c_i \in \mathbb{C}$  so that

$$(6.2) \quad f = \sum_{i=1}^n L_i^2, \quad g = \sum_{i=1}^n c_i L_i^2.$$

If  $M_f, M_g$  are the matrices associated to  $f, g$ , then the  $c_i$ 's are the  $n$  roots of the determinantal equation  $\det(M_g - \lambda M_f) = 0$ , which are generally distinct, so the  $L_i$ 's are uniquely determined up to multiple. We may also assume that the coefficients  $\alpha_{ij}$  of the linear forms are generally non-zero; cf. Corollary 6.3.

*Proof of Theorem 1.3.* For general  $p \in H_3(\mathbb{C}^n)$ , we simultaneously diagonalize  $f = \frac{\partial p}{\partial x_1}$  and  $g = \frac{\partial p}{\partial x_2}$  as in (6.2). Since mixed partials are equal,

$$(6.3) \quad \frac{\partial f}{\partial x_2} = \frac{\partial g}{\partial x_1} = \sum_{i=1}^n 2\alpha_{i2}L_i = \sum_{i=1}^n 2c_i\alpha_{i1}L_i,$$

and since the  $L_i$ 's are linearly independent, (6.3) implies that  $\alpha_{i2} = c_i\alpha_{i1}$ .

It is generally true that  $\alpha_{i1} \neq 0$ . Let

$$q(x_1, \dots, x_n) = p(x_1, \dots, x_n) - \sum_{i=1}^n \frac{1}{3\alpha_{i1}} L_i^3.$$

It follows that

$$\begin{aligned} \frac{\partial q}{\partial x_1} &= \frac{\partial p}{\partial x_1} - \sum_{i=1}^n \frac{3\alpha_{i1}}{3\alpha_{i1}} L_i^2 = \frac{\partial p}{\partial x_1} - \sum_{i=1}^n L_i^2 = 0, \\ \frac{\partial q}{\partial x_2} &= \frac{\partial p}{\partial x_2} - \sum_{i=1}^n \frac{3\alpha_{i2}}{3\alpha_{i1}} L_i^2 = \frac{\partial p}{\partial x_2} - \sum_{i=1}^n c_i L_i^2 = 0. \end{aligned}$$

Since  $\frac{\partial q}{\partial x_1} = \frac{\partial q}{\partial x_2} = 0$ , we have  $q = q(x_3, \dots, x_n)$ .

For uniqueness, suppose (1.5) holds and  $\ell_k(x_1, \dots, x_n) = \sum_j \beta_{kj}x_j$ . Then

$$\frac{\partial p}{\partial x_1} = \sum_{k=1}^n 3\beta_{k1}\ell_k^2; \quad \frac{\partial p}{\partial x_2} = \sum_{k=1}^n 3\beta_{k2}\ell_k^2.$$

Thus, after a scaling,  $\frac{\partial p}{\partial x_1}$  and  $\frac{\partial p}{\partial x_2}$  have already been simultaneously diagonalized (as in (6.2)), and the  $\ell_k$ 's are, up to multiples, a rearrangement of the  $L_k$ 's.  $\square$

We now give a constructive proof of Theorem 1.4, which gives a different canonical form for  $H_3(\mathbb{C}^n)$  requiring even more cubes.

*Proof of Theorem 1.4.* The constant-counting makes this a potential canonical form: the variables are  $t_{\{i,j\},k}$  with  $1 \leq i \leq j \leq k \leq n$ , and there are  $\binom{n+2}{3} = N(n, 3)$  such triples  $(i, j, k)$ . Given  $p \in H_3(\mathbb{C}^n)$ ,  $\frac{\partial p}{\partial x_n}$  is a quadratic form, so we can generally complete the square by Theorem 1.2:

$$\frac{\partial p}{\partial x_n} = \sum_{j=1}^n (t_{jj}x_j + \cdots + t_{jn}x_n)^2.$$

Then  $t_{jn} \neq 0$  for general  $p$  and if we let

$$q(x_1, \dots, x_n) = p(x_1, \dots, x_n) - \sum_{j=1}^n \frac{1}{3t_{jn}} (t_{jj}x_j + \cdots + t_{jn}x_n)^3,$$

then  $\frac{\partial q}{\partial x_n} = 0$ , so  $q = q(x_1, \dots, x_{n-1})$ . Iterate this construction to get (1.6).

Uniqueness follows by working backwards. If (1.6) holds for a cubic  $p$ , then it gives  $\frac{\partial p}{\partial x_n}$  in its (unique) upper-triangular diagonalization. This can be integrated with respect to  $x_n$  and subtracted from  $p$ , giving a cubic  $q(x_1, \dots, x_{n-1})$ . Again, iterate.  $\square$

It is not hard to give nonconstructive proofs of Theorems 1.3 and 1.4 using Corollary 2.3. These are left for the reader.

We first presented this next construction in [33]; an outline of the proof can be found in [6]. This is not a canonical form, but is included here because it gives an absolute upper bound for the length of cubic forms.

**Theorem 6.2.** *If  $p \in H_3(\mathbb{C}^n)$ , then there exists an invertible linear change of variables  $y_j = \sum \lambda_{jk}x_k$  and  $n$  linear forms  $\ell_j$  so that for some  $q \in H_3(\mathbb{C}^{n-1})$ ,*

$$(6.4) \quad p(x_1, \dots, x_n) = \sum_{j=1}^n \ell_j^3(x_1, \dots, x_n) + q(y_2, \dots, y_n).$$

*Thus every cubic in  $n$  variables is a sum of at most  $\binom{n+1}{2}$  cubes of linear forms.*

*Proof.* Define linear forms  $\ell_{j,m}(y)$  for  $1 \leq j \leq m+1$  by

$$(6.5) \quad \begin{aligned} \ell_{j,m}(y_1, \dots, y_n) &= y_j + \alpha \sum_{j=1}^m y_j, & 1 \leq j \leq m, \\ \ell_{m+1,m}(y_1, \dots, y_n) &= -(1 + m\alpha) \sum_{j=1}^m y_j, & \alpha = \frac{-(m+1) + \sqrt{m+1}}{m(m+1)}. \end{aligned}$$

Then it can be easily checked that

$$(6.6) \quad \sum_{j=1}^{m+1} \ell_{j,m}(y) = 0 \quad \text{and} \quad \sum_{j=1}^{m+1} \ell_{j,m}^2(y) = \sum_{k=1}^m y_k^2.$$

Suppose  $0 \neq p \in H_3(\mathbb{C}^n)$ . Use Biermann's Theorem to find a point  $u$  where  $p(u) \neq 0$ , and after an invertible linear change of variables, taking  $\{x_j\} \mapsto \{u_j\}$ , we may assume that  $p(1, 0, \dots, 0) = 1$  and so

$$(6.7) \quad p = u_1^3 + 3h_1(u_2, \dots, u_n)u_1^2 + 3h_2(u_2, \dots, u_n)u_1 + h_3(u_2, \dots, u_n),$$

where  $\deg h_j = j$ . Now let  $u_1 = y_1 - h_1(u_2, \dots, u_n)$  to clear the quadratic term, so

$$(6.8) \quad p = y_1^3 + 3y_1\tilde{h}_2(u_2, \dots, u_n) + \tilde{h}_3(u_2, \dots, u_n),$$

where again  $\deg \tilde{h}_j = j$ . Diagonalize  $\tilde{h}_2(u_2, \dots, u_n)$  as a quadratic form into  $y_2^2 + \dots + y_r^2$ , where  $r \leq n$ , and make the accompanying change of variables. We now have

$$(6.9) \quad p = y_1^3 + 3y_1(y_2^2 + \dots + y_r^2) + k_3(y_2, \dots, y_n); \quad r \leq n,$$

where  $\deg k_3 = 3$ . Finally, using (6.5) and (6.6), we construct  $g$ , a sum of  $r \leq n$  cubes:

$$(6.10) \quad \begin{aligned} g(y_1, \dots, y_n) &:= \frac{1}{r} \sum_{j=1}^r (y_1 + \sqrt{r} \cdot \ell_{j,r-1}(y_2, \dots, y_r))^3 \\ &= \frac{1}{r} \sum_{j=1}^r y_1^3 + \frac{3}{\sqrt{r}} \sum_{j=1}^r y_1^2 \ell_{j,r-1} + 3 \sum_{j=1}^r y_1 \ell_{j,r-1}^2 + \sqrt{r} \sum_{j=1}^r \ell_{j,r-1}^3 \\ &= y_1^3 + 3y_1(y_2^2 + \dots + y_r^2) + \sqrt{r} \sum_{j=1}^r \ell_{j,r-1}^3(y_2, \dots, y_r). \end{aligned}$$

Then  $q := p - g$  is a cubic form in  $(y_2, \dots, y_n)$  as in (6.4). Iteration of this argument shows that any cubic  $p \in H_3(\mathbb{C}^n)$  is a sum of at most  $\frac{n(n+1)}{2}$  cubes.  $\square$

Theorem 1.5 can be extended to a canonical form for quartics as a sum of fourth powers of linear forms. Note that  $x_n$  appears in each summand of (6.1), with, generally, a non-zero coefficient.

**Corollary 6.3.** *For general  $p \in H_4(\mathbb{C}^n)$ , there exist  $\ell_k \in H_1(\mathbb{C}^n)$  and  $q \in H_4(\mathbb{C}^{n-1})$  so that, with  $a(n) = \lfloor \frac{(n+1)^2}{4} \rfloor$ ,*

$$p(x_1, \dots, x_n) = \sum_{k=1}^{a(n)} \ell_k(x_1, \dots, x_n)^4 + q(x_1, \dots, x_{n-1}).$$

As a consequence, a general  $p \in H_4(\mathbb{C}^n)$  can be written as

$$p(x_1, \dots, x_n) = \sum_{m=0}^{\lfloor (n-1)/2 \rfloor} \sum_{r=1+2m}^n \sum_{k=1}^{r-2m} (t_{m,r,1+2m}^{\{k\}} x_{1+2m} + \dots + t_{m,r,r}^{\{k\}} x_r)^4.$$

*Proof.* By Corollary 1.3 and (6.1), for general  $p \in H_4(\mathbb{C}^n)$ , we can write

$$(6.11) \quad \begin{aligned} \frac{\partial p}{\partial x_n} &= \sum_{m=0}^{\lfloor (n-1)/2 \rfloor} \sum_{k=1}^{n-2m} (t_{m,1+2m}^{\{k\}} x_{1+2m} + \cdots + t_{m,n}^{\{k\}} x_n)^3 \\ &=: \sum_{m=0}^{\lfloor (n-1)/2 \rfloor} \sum_{k=1}^{n-2m} (\ell_m^{\{k\}}(x))^2. \end{aligned}$$

As before, if  $q = p - \sum_{k,m} \frac{1}{4t_{m,n}^{\{k\}}} \ell_{k,m}^4$ , then  $\frac{\partial q}{\partial x_n} = 0$ , so  $q = q(x_1, \dots, x_{n-1})$ . Repeat as before. There are  $N(n, 3)$  coefficients in (6.11), and since  $N(n, 3) + N(n-1, 4) = N(n, 4)$ , the count is correct for a canonical form.  $\square$

Note that there is no variable which appears in each linear form in (6.11), so the argument can't be extended to quintics. For the same reason, Theorem 1.4 does not extend to quartics. By combining Theorems 1.3 and 6.3, we obtain canonical forms as a sum of powers of linear forms in the four exceptional cases of Alexander-Hirschowitz, of course at the expense of the number of summands. With regards to ternary quartics and Theorem 1.10, Corollary 6.3 becomes the following canonical form for  $H_4(\mathbb{C}^3)$  as a sum of seven fourth powers.

$$\sum_{k=1}^3 (t_{k1}x_1 + t_{k2}x_2 + t_{k3}x_3)^4 + t_{10}x_3^4 + \sum_{\ell=1}^2 (u_{\ell 1}x_1 + u_{\ell 2}x_2)^4 + u_5x_1^4.$$

There is an arithmetic obstruction to a ‘‘Reichstein-type’’ canonical form for quartics; that is, one in which each linear form is allowed to involve each variable. If

$$(6.12) \quad p(x_1, \dots, x_n) = \sum_{k=1}^r (\alpha_{k1}x_1 + \cdots + \alpha_{kn}x_n)^4 + q(x_1, \dots, x_m).$$

were a canonical form for some  $n$ , then we would have  $N(n, 4) = rn + N(m, 4)$ . However, for  $n = 12$ , there does not exist  $m < 12$  so that  $12 \mid \binom{15}{4} - \binom{m+3}{4}$ , so no such canonical form can exist. More generally, let

$$(6.13) \quad A_d = \left\{ n : 0 \leq m < n \implies n \nmid \binom{n+d-1}{d} - \binom{m+d-1}{d} \right\}$$

denote the set of  $n$  for which this argument rules out Reichstein-type canonical forms. We present without proof a number of results about  $A_d$ . Note that there is no obstacle for (6.12) in prime degree, such as  $d = 2, 3$ .

**Proposition 6.4.**

- (i) If  $3 \nmid k$ , then  $n = 2^{2k} \cdot 3 \in A_4$ .
- (ii) If  $p \equiv 1 \pmod{144}$  is prime, then  $12p \in A_4$ .
- (iii) If  $p$  is prime, then  $p \mid \binom{n+p-1}{p} - \binom{n}{p}$ , hence  $A_p = \emptyset$  for prime  $p$ .
- (iv) The smallest elements of  $A_6, A_8, A_{10}, A_{12}, A_{14}$  and  $A_{15}$  are 10, 1792, 6, 242, 338 and 273 respectively. If  $A_9$  or  $A_{16}$  are non-empty, then their smallest elements are at least  $10^5$ .

## 7. SUBSPACE CANONICAL FORMS AND THE PROOF OF THEOREM 1.11

One natural generalization of the definition of canonical forms is to consider maps  $F : X \mapsto H_d(\mathbb{C}^n)$  where  $X \subset \mathbb{C}^M$  is an  $N(n, d)$ -dimensional subspace of  $\mathbb{C}^M$ . (Similar ideas can be found in Wakeford [46], though his approach is different from ours.) These can be analyzed in the simplest non-trivial case:  $M = 4, N(2, 2) = 3$ .

*Proof of Theorem 1.11.* Assume that some  $c_j \neq 0$ . Without loss of generality, we may assume that  $c_4 \neq 0$  and divide through by  $c_4$  so that the equation is  $t_4 = a_1 t_1 + a_2 t_2 + a_3 t_3$ , where  $a_i = -c_i/c_4$  for  $i = 1, 2, 3$ . Then (1.13) can be parameterized as a map from  $\mathbb{C}^3 \mapsto H_2(\mathbb{C}^2)$  as:

$$(7.1) \quad F(t; x) = (t_1 x + t_2 y)^2 + (t_3 x + (a_1 t_1 + a_2 t_2 + a_3 t_3) y)^2.$$

The partials with respect to the  $t_j$ 's are:

$$(7.2) \quad \begin{aligned} & 2x(t_1 x + t_2 y) + 2a_1 y(t_3 x + (a_1 t_1 + a_2 t_2 + a_3 t_3) y), \\ & 2y(t_1 x + t_2 y) + 2a_2 y(t_3 x + (a_1 t_1 + a_2 t_2 + a_3 t_3) y), \\ & 2(x + a_3 y)(t_3 x + (a_1 t_1 + a_2 t_2 + a_3 t_3) y). \end{aligned}$$

Now, (7.1) is a canonical form if and only if there exists a choice of  $t_i$  so that the three quadratics in (7.2) span  $H_2(\mathbb{C}^2)$ . A computation shows that the determinant of the forms in (7.2) with respect to the basis  $\{x^2, xy, y^2\}$  is the cubic

$$(7.3) \quad -8((a_1 a_2 - a_3)t_1 + (1 + a_2^2)t_2 + (a_2 a_3 + a_1)t_3)(a_1 t_1^2 + a_2 t_1 t_2 + a_3 t_1 t_3 - t_2 t_3).$$

The second factor in (7.3) always has the term  $-t_2 t_3$  and so never vanishes, hence this determinant is not identically zero (and (7.1) is a canonical form), unless

$$(7.4) \quad a_1 a_2 - a_3 = 1 + a_2^2 = a_2 a_3 + a_1 = 0.$$

In the exceptional case where (7.4) holds, then  $a_2 = \epsilon$ , where  $\epsilon = \pm i$ , and  $a_3 = \epsilon a_1$ . Evaluating (7.1) at  $(x, y) = (a_1, \epsilon)$  yields

$$\begin{aligned} & (a_1 t_1 + \epsilon t_2)^2 + (a_1 t_3 + \epsilon a_1 t_1 + \epsilon^2 t_2 + \epsilon^2 a_1 t_3)^2 \\ & = (a_1 t_1 + \epsilon t_2)^2 + ((1 + \epsilon^2)a_1 t_3 + \epsilon a_1 t_1 + \epsilon^2 t_2)^2 = (a_1 t_1 + \epsilon t_2)^2 + \epsilon^2 (a_1 t_1 + \epsilon t_2)^2 = 0, \end{aligned}$$

as claimed.  $\square$

It would be interesting to know how Theorem 1.11 generalizes to higher degrees.

Conjecture 1.12 is true for degree 2 by Theorem 1.11. We have verified it for even degrees up to eight by Corollary 2.3 applied to random choices for  $\alpha_j, \beta_j$  in (1.14). We hold some hope that generalizations such as Conjecture 1.12 will have applications in more than two variables as well.

## REFERENCES

- [1] J. Alexander and A. Hirschowitz, *Polynomial interpolation in several variables*, J. Algebraic Geom., **4** (1995), 201–222, MR1311347 (96f:14065).
- [2] D. W. Babbage, *Canonical forms for general polynomials – a note on a paper by Stefano Guazzone*, J. London Math. Soc., (2)**13** (1976), 564, MR0417206 (54 #5264).
- [3] E. Carlini, M. V. Catalisano and A. Geramita, *The solution to the Waring problem for monomials and the sum of pairwise coprime monomials*, J. Algebra, **370** (2012), 5–14, MR2966824.
- [4] M. D. Choi, T. Y. Lam and B. Reznick, *Sums of squares of real polynomials K-theory and algebraic geometry: connections with quadratic forms and division algebras* (Santa Barbara, CA, 1992), 103–126, Proc. Sympos. Pure Math., **58**, Part 2, Amer. Math. Soc., Providence, RI, 1995, MR1327293 (96f:11058).
- [5] A. Clebsch, *Ueber Curven vierter Ordnung*, J. Reine Angew. Math, **59** (1861), 125–145.
- [6] P. Comon and B. Mourrain, *Decomposition of quantics in sums of powers of linear forms*, Signal Processing, **53** (1996), 93–107.
- [7] D. Cox, J. Little and D. O’Shea, *Ideals, varieties, and algorithms. An introduction to computational algebraic geometry and commutative algebra, Third edition*, Undergraduate Texts in Mathematics. Springer, New York, 2007, MR2290010 (2007h:13036).
- [8] R. Ehrenborg and G.-C. Rota, *Apolarity and canonical forms for homogeneous polynomials*, European J. Combin., **14** (1993), 157–181, MR1215329 (94e:15062).
- [9] E. B. Elliott, *Algebra of Quantics*, 2nd ed., Chelsea, New York, 1964, originally published by Oxford at the Clarendon Press, 1913.
- [10] C. S. Fisher, *The last invariant theorists*, Archiv. europ. sociol., **VIII** (1967), 216–244.
- [11] R. Fröberg, G. Ottaviani and B. Shapiro, *On the Waring problem for polynomial rings*, Proc. Natl. Acad. Sci USA **109** (2012), 5600–5602, MR2935563.
- [12] A. Geramita, *Inverse systems of fat points: Waring’s problem, secant varieties of Veronese varieties and parameter spaces for Gorenstein ideals*, The Curves Seminar at Queen’s, Vol. X (Kingston, ON, 1995), 2–114, Queen’s Papers in Pure and Appl. Math., **102**, Queen’s Univ., Kingston, ON, 1996, MR1381732 (97h:13012).
- [13] J. H. Grace, *Edward Kingsley Wakeford*, Proc. London Math. Soc., **16** (1917/1918), liv–lvii.
- [14] S. Guazzone, *Canonical forms for general polynomials* J. London Math. Soc. (2)**10** (1975), 271–280, MR0382298 (52 #3183).
- [15] S. Gundelfinger, *Zur Theorie der binären Formen*, J. Reine Angew. Math., **100** (1887), 413–424.
- [16] D. Hilbert, *Über die Darstellung definiter Formen als Summe von Formenquadraten*, Math. Ann., **32** (1888), 342–350; see Ges. Abh. 2, 154–161, Springer, Berlin, 1933, reprinted by Chelsea, New York, 1981, MR1510517.
- [17] A. Iarrobino and V. Kanev, *Power Sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics, **1721** (1999), MR1735271 (2001d:14056).
- [18] J. P. S. Kung, *Gundelfinger’s theorem on binary forms*, Stud. Appl. Math., **75** (1986), 163–169, MR0859177 (87m:11020).
- [19] J. P. S. Kung, *Canonical forms for binary forms of even degree*, in *Invariant theory*, Lecture Notes in Mathematics, **1278**, 52–61, Springer, Berlin, 1987, MR0924165 (89h:15037).
- [20] J. P. S. Kung, *Canonical forms of binary forms: variations on a theme of Sylvester*, in *Invariant theory and tableaux (Minnesota, MN, 1988)*, 46–58, IMA Vol. Math. Appl., **19**, Springer, New York, 1990, MR1035488 (91b:11046).
- [21] J. P. S. Kung and G.-C. Rota, *The invariant theory of binary forms*, Bull. Amer. Math. Soc. (N. S.), **10** (1984), 27–85, MR0722856 (85g:05002).
- [22] J. M. Landsberg, *Tensors: geometry and applications*, Graduate Studies in Mathematics, **128**, Amer. Math. Society, Providence, 2012, MR2865915.

- [23] J. M. Landsberg and Z. Teitler, *On the ranks and border ranks of symmetric tensors*, *Found. Comput. Math.*, **10** (2010), 339–366, MR2628829 (2011d:14095).
- [24] E. Lasker, *Zur Theorie der kanonischen Formen*, *Math. Ann.* **58** (1904), 434–440, MR1511244.
- [25] Online Encyclopedia of Integer Sequences, <http://oeis.org>, accessed June 10, 2013.
- [26] A. Pfister and C. Scheiderer, *An elementary proof of Hilbert’s theorem on ternary quartics*, *J. Algebra* **371** (2012), 1–25, MR2975385.
- [27] D. Plaumann, B. Sturmfels and C. Vinzant, *Quartic curves and their bitangents*, *J. Symbolic Comput.* **46** (2011), 712–733, MR2781949 (2012e:14065).
- [28] V. Powers and B. Reznick, *Notes towards a constructive proof of Hilbert’s Theorem on ternary quartics*, *Proceedings, Quadratic forms and their applications, Dublin 1999* (A. Ranicki ed.), *Cont. Math.*, **272** (2000), 209–227, MR1803369 (2001h:11049).
- [29] V. Powers, B. Reznick, C. Scheiderer and F. Sottile, *A new approach to Hilbert’s theorem on ternary quartics*, *C. R. Math. Acad. Sci. Paris*, **339** (2004), 617–620, MR2103198 (2005i:11051).
- [30] K. Ranestad and F.-O. Schreyer, *Varieties of sums of powers*, *J. Reine Angew. Math.* **525** (2000), 147–181, MR1780430 (2001m:14009).
- [31] B. Reichstein, *On expressing a cubic form as a sum of cubes of linear forms*, *Linear Algebra Appl.*, **86** (1987), 91–122, MR0870934 (88e:11022).
- [32] B. Reznick, *Sums of even powers of real linear forms*, *Mem. Amer. Math. Soc.*, **96** (1992), no. 463, MR1096187 (93h:11043).
- [33] B. Reznick, *Sums of powers of complex linear forms*, Preprint, August 1992.
- [34] B. Reznick, *Homogeneous polynomial solutions to constant coefficient PDE’s*, *Adv. Math.*, **117** (1996), 179–192, MR1371648 (97a:12006).
- [35] B. Reznick, *Some concrete aspects of Hilbert’s 17th Problem*, *Contemp. Math.*, **253** (2000), 251–272, MR 1747589 (2001i:11042).
- [36] B. Reznick, *On the length of binary forms*, to appear in *Quadratic and Higher Degree Forms*, (K. Alladi, M. Bhargava, D. Savitt, P. Tiep, eds.), *Developments in Math.*, Springer, New York, arXiv:1007.5485.
- [37] B. Reznick, *Linear dependence among powers of quadratic forms*, in preparation.
- [38] H. W. Richmond, *On canonical forms*, *Quart. J. Pure Appl. Math.*, **33** (1902), 331–340.
- [39] J.J. Sylvester, *An Essay on Canonical Forms, Supplement to a Sketch of a Memoir on Elimination, Transformation and Canonical Forms*, originally published by George Bell, Fleet Street, London, 1851; pp. 203–216 in Paper 34 in *Mathematical Papers*, Vol. I, Chelsea, New York, 1973. Originally published by Cambridge University Press in 1904.
- [40] J. J. Sylvester, *On a remarkable discovery in the theory of canonical forms and of hyperdeterminants*, originally in *Philosophical Magazine*, vol. II, 1851; pp. 265–283 in Paper 41 in *Mathematical Papers*, Vol. 1, Chelsea, New York, 1973. Originally published by Cambridge University Press in 1904.
- [41] J. J. Sylvester, *Additional notes to Prof. Sylvester’s Exeter British Association Address*, originally published as an appendix to *The laws of verse* (Longmans, London, 1870), p. 714 in *Mathematical Papers*, Vol. II, Chelsea, New York, 1973. Originally published by Cambridge University Press in 1908.
- [42] H. W. Turnbull, *The theory of determinants, matrices and invariants*, 3rd ed. Dover Publications, Inc., New York 1960, MR0130257 (24 #A123).
- [43] A. Várilly-Alvarado, *Weak approximation on del Pezzo surfaces of degree 1*, *Adv. Math.*, **219** (2008), 2123–2145, MR2456278 (2009j:14045).
- [44] A. Várilly-Alvarado, *Density of rational points on isotrivial rational elliptic surfaces*, *Algebra Number Theory*, **5** (2011), 659–690, MR2889751.
- [45] E. K. Wakeford, *A canonical form of the binary sextic*, *Mess. of Mathematics*, **43** (1914), 25–28.

- [46] E. K. Wakeford, *On canonical forms*, Proc. London Math. Society, **(2)18**, (1920), 403-410, MR1576066.

DEPARTMENT OF MATHEMATICS AND CENTER FOR ADVANCED STUDY, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801

*E-mail address:* reznick@math.uiuc.edu