

Balloons, Cut-Edges, Matchings, and Total Domination in Regular Graphs of Odd Degree

Suil O*, Douglas B. West†

November 9, 2008; revised June 2, 2009

Abstract

A *balloon* in a graph G is a maximal 2-edge-connected subgraph incident to exactly one cut-edge of G . Let $b(G)$ be the number of balloons, let $c(G)$ be the number of cut-edges, and let $\alpha'(G)$ be the maximum size of a matching. Let $\mathcal{F}_{n,r}$ be the family of connected $(2r + 1)$ -regular graphs with n vertices, and let $b = \max\{b(G) : G \in \mathcal{F}_{n,r}\}$. For $G \in \mathcal{F}_{n,r}$, we prove the sharp inequalities $c(G) \leq \frac{r(n-2)-2}{2r^2+2r-1} - 1$ and $\alpha'(G) \geq \frac{n}{2} - \frac{rb}{2r+1}$. Using $b \leq \frac{(2r-1)n+2}{4r^2+4r-2}$, we obtain a simple proof of the bound $\alpha'(G) \geq \frac{n}{2} - \frac{r}{2} \frac{(2r-1)n+2}{(2r+1)(2r^2+2r-1)}$ proved by Henning and Yeo. For each of these bounds and each r , the approach using balloons allows us to determine the infinite family where equality holds. For the total domination number $\gamma_t(G)$ of a cubic graph, we prove $\gamma_t(G) \leq \frac{n}{2} - \frac{b(G)}{2}$ (except that $\gamma_t(G)$ may be $n/2 - 1$ when $b(G) = 3$ and the balloons cover all but one vertex). With $\alpha'(G) \geq \frac{n}{2} - \frac{b(G)}{3}$ for cubic graphs, this improves the known inequality $\gamma_t(G) \leq \alpha'(G)$.

1 Introduction

A graph is a *cubic graph* if every vertex has degree 3. In 1891, Petersen [12] proved that every cubic graph without cut-edges has a perfect matching. It is natural to ask how small $\alpha'(G)$ can be in a cubic graph G with n vertices, where $\alpha'(G)$ is the maximum size of a matching in G (called the *matching number* of G). Chartrand et al. [5] proved that $\alpha'(G) \geq n/2 - \lceil c(G)/3 \rceil$ when G is a cubic n -vertex graph, where $c(G)$ denotes the number of cut-edges in G .

*Department of Mathematics, University of Illinois, Urbana, IL, 61801, suilo2@math.uiuc.edu. research partially supported by the Korean Research Foundation (MOEHRD, Basic Research Promotion Fund), grant KRF-2005-C00003.

†Department of Mathematics, University of Illinois, Urbana, IL, 61801, west@math.uiuc.edu; research is partially supported by the National Security Agency under Award No. H98230-06-1-0065.

By this result, an upper bound on $c(G)$ yields a lower bound on $\alpha'(G)$. Let G be a connected cubic graph with n vertices. In Section 3, we prove that $c(G) \leq (n - 7)/3$ and that this is sharp. The result of [5] then yields $\alpha'(G) \geq (7n + 14)/18$, but this is not the best bound on $\alpha'(G)$. The smallest value of $\alpha'(G)$ is $\lceil (4n - 1)/9 \rceil$, proved first by Biedl et al. [4]. Henning and Yeo [9] generalized the result, proving that $\alpha'(G) \geq \frac{n}{2} - \frac{r}{2} \frac{(2r-1)n+2}{(2r+1)(2r^2+2r-1)}$ when G is a $(2r + 1)$ -regular n -vertex connected graph, which is sharp.

Although maximizing $c(G)$ in a cubic graph does not minimize $\alpha'(G)$, another concept does yield a simple proof of the sharp bound on $\alpha'(G)$. We define a *balloon* in a graph G to be a maximal 2-edge-connected subgraph of G incident to exactly one cut-edge of G . The term arises from viewing the cut-edge as a string tying the balloon to the rest of the graph; the vertex incident to the cut-edge is the *neck* of the balloon. A balloon may contain cut-vertices and thus consist of several blocks.

Maximal 2-edge-connected subgraphs are pairwise disjoint, since the union of two 2-edge-connected subgraphs sharing a vertex is also 2-edge-connected. Among these subgraphs, the balloons are those incident to precisely one cut-edge. Thus the number of balloons in G is well-defined; let $b(G)$ denote this number.

Let $\mathcal{F}_{n,r}$ be the family of connected $(2r + 1)$ -regular graphs with n vertices. Let $b = \max\{b(G) : G \in \mathcal{F}_{n,r}\}$. For $G \in \mathcal{F}_{n,r}$, we prove that $c(G) \leq \frac{r(n-2)-2}{2r^2+2r-1} - 1$ and $\alpha'(G) \geq \frac{n}{2} - \frac{rb}{2r+1}$. We obtain a lower bound on $\alpha'(G)$ by proving that $b \leq \frac{(2r-1)n+2}{4r^2+4r-2}$, and we use balloons to prove the upper bound on $c(G)$. In Section 2, we construct an infinite family \mathcal{H}_r showing that all these bounds are sharp; it contains the smaller families provided in [4] and [9] (graphs in \mathcal{H}_r exist when $n \equiv 4(r + 1)^2 \pmod{(8r^3 + 12r^2 - 2)}$). The bounds for $b(G)$ and $c(G)$ are sharp in a larger family \mathcal{H}'_r (occurring when $n \equiv (4r + 6) \pmod{(4r^2 + 4r - 2)}$). In Section 3, we prove the upper bounds on $b(G)$ and $c(G)$ and show that equality holds if and only if $G \in \mathcal{H}'_r$. In Section 4, we prove the lower bound on $\alpha'(G)$; in Section 5, we show that equality holds if and only if $G \in \mathcal{H}_r$.

The restriction to connected graphs is important; consider cubic graphs. For a connected cubic graph, $b(G) \leq (n+2)/6$ and $\alpha'(G) \geq (4n-1)/9$. However, if G consists of many disjoint copies of the unique 16-vertex cubic graph with no perfect matching, then $b(G) = 3n/16$ and $\alpha'(G) = 7n/16$; these values are more extreme than the bounds for graphs in $\mathcal{F}_{n,3}$.

Our results apply only for regular graphs of odd degree, since a connected regular graph of even degree is 2-edge-connected and hence has no balloons. In addition to solving the extremal problem for odd degree, Henning and Yeo [9] also determined the smallest matching number for connected $2r$ -regular graphs with n vertices. We will use a generalization of balloons to address this in a subsequent paper [11].

In Section 6, we apply balloons to total domination. A *total dominating set* in a graph G is a set S of vertices in G such that every vertex in G has a neighbor in S . The *total domination*

number, written $\gamma_t(G)$, is the least size of a total dominating set in G . For n -vertex graphs with minimum degree at least 3, $\gamma_t(G) \leq n/2$ ([2, 15], for 3-regular graphs in [6]). Henning, Kang, Shan, and Yeo [8] proved the stronger result that $\gamma_t(G) \leq \alpha'(G)$ whenever G is regular with degree at least 3. For degree at least 4, stronger bounds hold. Thomassé and Yeo [13] proved that $\gamma_t(G) \leq 3n/7$ for every n -vertex regular graph with degree at least 4. This upper bound is a smaller fraction of n than the lower bound on $\alpha'(G)$. Earlier, Henning and Yeo [9] observed that $\gamma_t(G) < \alpha'(G)$ when G is a regular graph with degree at least 4.

We use balloons to strengthen the bound for cubic graphs. We prove that $\gamma_t(G) \leq \frac{n}{2} - \frac{b(G)}{2}$ when G is cubic, except that $\gamma_t(G) = n/2 - 1$ is possible when $b(G) = 3$ and the balloons cover all but one vertex. Since $\alpha'(G) \geq \frac{n}{2} - \frac{b(G)}{3}$ for cubic graphs, we have large separation when the number of balloons is large, and $\gamma_t(G) = \alpha'(G)$ can happen in a cubic graph only when the number of balloons is 0 or when G consists of three balloons plus one vertex.

The condition of being connected is the same as being 1-edge-connected. A generalized version of balloons is useful for studying the restriction of these problems to k -edge-connected graphs. The extension of Petersen's result by Bähler [3] states that every $(2r+1)$ -regular $2r$ -edge-connected graph has a perfect matching. As the edge-connectivity rises, the lower bound on the matching number should also rise. In a subsequent paper [11], we will use generalized balloons to determine the smallest matching number for d -regular k -edge-connected graphs with n vertices, when $d \geq 4$ and $k \geq 2$. In addition, there we use balloons to determine the worst case of the Chinese Postman Problem for 3-regular graphs with n vertices.

2 The Construction

Biedl et al. [4] and Henning and Yeo [9] presented examples for sharpness in the lower bounds on $\alpha'(G)$ for connected 3-regular and $(2r+1)$ -regular graphs, respectively. We present a more general family that includes their examples.

Construction 2.1. Let B_r be the graph obtained from the complete graph K_{2r+3} by deleting a matching of size $r+1$ and one more edge incident to the remaining vertex. This is the smallest graph in which one vertex has degree $2r$ and the others have degree $(2r+1)$. Thus B_r is the smallest possible balloon in a $(2r+1)$ -regular graph. Note that deleting the vertex of degree $2r$ (the neck) from B_r leaves a subgraph having a perfect matching.

Let \mathcal{T}'_r be the family of trees such that every non-leaf vertex has degree $2r+1$. Let \mathcal{H}'_r be the family of $(2r+1)$ -regular graphs obtained from trees in \mathcal{T}'_r by identifying each leaf of such a tree with the neck in a copy of B_r . Let \mathcal{T}_r be the subfamily of \mathcal{T}'_r obtained by requiring all leaves to have the same color in a proper 2-coloring. Let \mathcal{H}_r be the subfamily of \mathcal{H}'_r arising from trees in \mathcal{T}_r by adding balloons at leaves. \square

To compute the matching number for n -vertex graphs in \mathcal{H}_r , we use standard concepts about matchings. The *deficiency* of a vertex set S in a graph G , written $\text{def}_G(S)$ or simply $\text{def}(S)$, is $o(G - S) - |S|$, where $o(H)$ is the number of components of H having an odd number of vertices. Every matching must leave at least $\text{def}(S)$ vertices unmatched, so for any S the quantity $\frac{1}{2}(n - \text{def}(S))$ is an upper bound on $\alpha'(G)$. Furthermore, if there is a matching that matches S into vertices of distinct odd components of $G - S$ and leaves at most one unmatched vertex in each odd component of $G - S$, then $\alpha'(G) = \frac{1}{2}(n - \text{def}(S))$.

Proposition 2.2. *Let $p_r = 2r^2 + 2r - 1$. For any n -vertex graph G in \mathcal{H}_r ,*

$$\begin{aligned} n &\equiv 4(r+1)^2 \pmod{(4r+2)p_r}, & b(G) &= \frac{(2r-1)n+2}{2p_r}, \\ \alpha'(G) &= \frac{1}{2} \left(n - \frac{r(2r-1)n+2r}{(2r+1)p_r} \right), & c(G) &= \frac{r(n-2)-2}{p_r} - 1. \end{aligned}$$

Furthermore, the formulas given for $b(G)$ and $c(G)$ also hold when $G \in \mathcal{H}'_r$.

Proof. We first compute $b(G)$ and $c(G)$ on \mathcal{H}'_r . The smallest tree in \mathcal{T}'_r has two vertices. The resulting graph in \mathcal{H}'_r has $4r + 6$ vertices, two balloons, and one cut-edge, and the formulas hold. For any larger tree T in \mathcal{T}'_r , the penultimate vertex of a longest path has $2r$ leaf neighbors, and deleting them yields a smaller tree T' in \mathcal{T}'_r . Let G and G' be the corresponding graphs in \mathcal{H}'_r . Compared to G' , in G there are $2r$ more cut-edges, $2r - 1$ more balloons, and $2r(2r + 3) - (2r + 2)$ more vertices. This last formula simplifies to $2p_r$, and hence the formulas for $b(G)$ and $c(G)$ in terms of n are established by induction on n .

Now consider the more restrictive families \mathcal{T}_r and \mathcal{H}_r . The smallest graph in \mathcal{T}_r is the star $K_{1,2r+1}$ with $2r + 1$ leaves. We claim that every other tree in \mathcal{T}_r arises from a smaller tree in \mathcal{T}_r by appending $2r$ edges at a leaf y and appending $2r$ additional edges at each new neighbor of y . This produces $(2r)^2$ leaves, which replace y in the set of leaves and are in the same partite set as y , so the larger graph lies in \mathcal{T}_r .

To prove that this generates all of \mathcal{T}_r , consider a longest path P in a tree $T \in \mathcal{T}_r$ such that T is not a star. Let y, z, w be the last three vertices on P , in order (w is the leaf). Since P is a longest path, all $2r$ neighbors of z other than y are leaves. Since leaves all lie in the same partite set, no neighbor of y is a leaf. Hence the $2r - 1$ neighbors of y not on P must all have $2r$ leaf neighbors (again since P is a longest path and non-leaves have degree $2r + 1$). Now T arises in the specified way from a smaller tree in \mathcal{T}_r having y as a leaf.

To compute $\alpha'(G)$ for $G \in \mathcal{H}_r$, let T be the corresponding tree in \mathcal{T}_r . Let X and Y be its partite sets, with Y containing the leaves. Let $S = X$. Now $o(G - S) = |Y|$, since each vertex of Y is an isolated vertex in $G - S$ or is the neck of a copy of B_r that is an odd component of $G - S$. Thus $\text{def}(S) = |Y| - |X|$. Root T at a vertex of X , and then match each vertex of S to one of its children, which is or lies in an odd component of $G - S$.

When that odd component is a copy of B_r , pair its remaining vertices in a matching. This produces a matching with exactly $\text{def}(S)$ uncovered vertices.

It therefore suffices to compare $\text{def}(S)$ and the formula for $\alpha'(G)$ inductively. When $T = K_{1,2r+1}$, we have $\text{def}(S) = 2r$. Adding the balloons yields $(2r + 3)(2r + 1) + 1$ (this equals $4(r + 1)^2$, giving the basis for the claim about n). The subtractive term in the formula for $\alpha'(G)$ is $\frac{r(2r-1)(4r^2+8r+4)+2r}{(2r+1)p_r}$, which equals $2r$.

For larger $G \in \mathcal{H}_r$, let T be the corresponding tree in \mathcal{T}_r , expanded from T' with corresponding graph $G' \in \mathcal{H}_r$. In the expansion, $|X|$ increases by $2r$ and $|Y|$ increases by $4r^2$, so $\text{def}(S)$ increases by $4r^2 - 2r$. Comparing G with G' , one balloon is lost and $4r^2$ are created; the number of vertices increases by $4r^2(2r + 3) + 2r - (2r + 2)$. The increase in n simplifies to $(4r + 2)p_r$ (completing the proof of the claim about n). The subtractive term in the formula for $\alpha'(G)$ thus increases by $r(2r - 1)2$, which equals the change in $\text{def}(S)$. \square

Corollary 2.3. *For n -vertex cubic graphs, the matching number of graphs in \mathcal{H}_1 is $\frac{4n-1}{9}$.*

3 Balloons and Cut-edges

Recall that $\mathcal{F}_{n,r}$ is the family of connected $(2r + 1)$ -regular graphs with n vertices. We begin by bounding the number of balloons for graphs in $\mathcal{F}_{n,r}$.

Every balloon in a $(2r + 1)$ -regular graph has at least $2r + 3$ vertices; it has at least $2r + 2$ vertices because it has a vertex of degree $2r + 1$, and equality cannot hold because then the degree-sum would be odd. Thus $b(G) \leq \frac{n}{2r+3}$. Surprisingly, this trivial upper bound can be improved only slightly; the optimal bound is $\frac{n+2\epsilon}{2r+3+\epsilon}$, where $\epsilon = 1/(2r - 1)$. Of course, $\epsilon = 1$ for cubic graphs. We use a counting argument; the bound can also be proved inductively.

Lemma 3.1. *If $G \in \mathcal{F}_{n,r}$, then $b(G) \leq \frac{(2r-1)n+2}{4r^2+4r-2}$, with equality if and only if $G \in \mathcal{H}'_r$.*

Proof. For $G \in \mathcal{F}_{n,r}$, let G' be the graph obtained from G by shrinking each balloon to a single vertex; G' is connected, and the balloons of G become vertices of degree 1 in G' . Let $n' = |V(G')|$ and $m' = |E(G')|$. Since G' is connected, $m' \geq n' - 1$, and the degree-sum formula yields $(2r + 1)n' - 2rb(G) = 2m' \geq 2n' - 2$. Thus $2rb(G) \leq (2r - 1)n' + 2$. Since each balloon has at least $2r + 3$ vertices, $n' \leq n - (2r + 2)b(G)$. Combining the inequalities yields $2rb(G) \leq (2r - 1)n + 2 - (2r - 1)(2r + 2)b(G)$, which simplifies to the desired bound.

Equality requires equality in each contributing inequality. Hence G' is a tree with non-leaf vertices having degree $2r + 1$. That is, $G' \in \mathcal{T}'_r$, and $G \in \mathcal{H}'_r$. \square

Corollary 3.2. *Every connected n -vertex cubic graph has at most $\frac{n+2}{6}$ balloons, and this is sharp for $n \equiv 4 \pmod{6}$.*

The bounds of Lemma 3.1 and Corollary 3.2 do not hold for disconnected graphs. An n -vertex graph consisting of disjoint copies of the smallest graph in \mathcal{H}_r has $\frac{2r+1}{6r+10}n$ balloons, which is more than the bound above.

Lemma 3.3. *The following hold for balloons and cut-edges in a graph G in $\mathcal{F}_{n,r}$.*

- (a) *Each component formed by deleting a cut-edge contains a balloon of G .*
- (b) *The number of vertices in a balloon may be any odd number that is at least $2r + 3$.*

Proof. (a) Let e be a cut-edge. Among the paths containing e , let P be a path containing the maximum number of cut-edges of G . The portion of P after the last cut-edge toward either end lies in a 2-edge-connected subgraph, and by the choice of P it is a balloon.

(b) In a balloon, the neck has degree $2r$, and other vertices have degree $2r + 1$. We form such a graph with $2k + 3$ vertices, where $k \geq r$. The complete graph K_{2k+3} decomposes into $k + 1$ spanning cycles. The union of r of these cycles plus a near-perfect matching from one of the remaining cycles is a 2-edge-connected graph with the desired degrees. \square

Theorem 3.4. *If $G \in \mathcal{F}_{n,r}$, then $c(G) \leq \frac{r(n-2)-2}{2r^2+2r-1} - 1$, with equality if and only if $G \in \mathcal{H}'_r$.*

Proof. We use induction on n . If $n \leq 4r + 6$, then the bound at most 1, with equality only when $n = 4r + 6$. Every graph having a cut-edge has at least two balloons and hence at least $4r + 6$ vertices, by Lemma 3.3. The graph with $4r + 6$ vertices consisting of two copies of B_r joined by an edge lies in \mathcal{H}'_r . Hence all claims hold for the basis.

For larger n , consider a cut-edge e in G . Let G_1 and G_2 be the components of $G - e$. Let G'_1 and G'_2 be the graphs obtained from G by replacing G_2 and G_1 , respectively, with B_r . The cut-edges of G consists of the cut-edges in G_1 and G_2 , plus e itself. Since e is a cut-edge in both G_1 and G_2 , and the added B_r contains no cut-edge, we have $c(G) = c(G'_1) + c(G'_2) - 1$. If neither G_1 nor G_2 equals B_r , then G'_1 and G'_2 have fewer vertices than G , and we can apply the induction hypothesis to both. Letting $n_i = |V(G'_i)|$, we have $n = n_1 + n_2 - (4r + 6)$. With $p_r = 2r^2 + 2r - 1$ (as in Proposition 2.2), we obtain the desired bound on $c(G)$:

$$\begin{aligned} c(G) &= c(G'_1) + c(G'_2) - 1 \leq \frac{r(n_1 - 2) - 2}{p_r} + \frac{r(n_2 - 2) - 2}{p_r} - 3 \\ &= \frac{r(n - 2) - 2}{p_r} + \frac{r(4r + 4) - 2}{p_r} - 3 = \frac{r(n - 2) - 2}{p_r} - 1. \end{aligned}$$

In the remaining case, every cut-edge in G is incident to a copy of B_r . Since each copy of B_r is incident to exactly one cut-edge, we obtain $c(G) = b(G)$ (note that $n > 4r + 6$). Let Q be the set of endpoints of cut-edges outside the balloons. If any two balloons have distinct nonadjacent neighbors in Q , then let G' be the graph obtained by deleting the two balloons and adding one edge to make their neighbors adjacent. The graph G' is connected

and $(2r+1)$ -regular and has $n - (4r+6)$ vertices. Crucially, G' has exactly $c(G) - 2$ cut-edges, because the only cut-edges in G are those incident to balloons. By the induction hypothesis,

$$c(G) \leq 2 + \frac{r(n - 4r - 8) - 2}{p_r} - 1 = \frac{r(n - 2) - 2}{p_r} - \frac{4r^2 + 6r}{p_r} + 1 < \frac{r(n - 2) - 2}{p_r} - 1.$$

Hence we may assume that the vertices of Q are pairwise adjacent. Let $q = |Q|$, and let S be the set of vertices outside both Q and the balloons. If $S = \emptyset$, then $c(G) = q(2r+2-q)$ and $n = (2r+3)c(G) + q$. Since $1 \leq q \leq 2r+1$, we obtain $n \geq (2r+3)(2r+1) + 1 = 2p_r + 4r + 6$. Since $c(G) = b(G)$, Lemma 3.1 yields $c(G) \leq \frac{(2r-1)n+2}{2p_r} = \frac{rn}{p_r} - \frac{n-2}{2p_r}$. It thus suffices to show that $\frac{n-2}{2p_r} \geq \frac{2r+2}{p_r} + 1$. This requires $n - 2 \geq 4r + 4 + 2p_r$, which we have proved for this case.

Finally, suppose that $S \neq \emptyset$. Each vertex of S has $2r+1$ neighbors outside the balloons, so $n \geq 2r+2 + (2r+3)c(G)$. If equality holds, then $S \cup Q$ induces a complete graph, $G = K_{2r+2}$, and $c(G) = 0$. Otherwise, $n \geq (2r+3)[c(G) + 1]$. Now $c(G) \leq \frac{n}{2r+3} - 1$, and we only need $\frac{n}{2r+3} \leq \frac{r(n-2)-2}{2r^2+2r-1}$. This simplifies to $n \geq 4r+6$, which holds when $c(G) > 0$.

For the characterization of equality, consider each case. When G has a cut-edge not incident to a balloon that is a copy of B_r , the induction hypothesis requires achieving equality for both G'_1 and G'_2 , which must therefore lie in \mathcal{H}'_r . The construction of G from G'_1 and G'_2 indeed puts G in \mathcal{H}'_r . When $c(G) = b(G)$ and two balloons have nonadjacent neighbors, we obtained strict inequality in the bound. When $c(G) = b(G)$ and $S = \emptyset$, equality requires $b(G)$ to meet its bound, which already requires $G \in \mathcal{H}'_r$ (indeed, it requires more, and equality is obtained only by putting copies of B_r at the leaves of the star $K_{1,2r+1}$). When $S \neq \emptyset$, equality requires $n = 4r+6$ and $c(G) = 1$, in which case G is the graph in \mathcal{H}'_r consisting of a cut-edge joining two copies of B_r . \square

Corollary 3.5. *Every n -vertex $(2r+1)$ -regular graph has at most $\frac{r(n-2)-2}{2r^2+2r-1} - 1$ cut-edges, which reduces to $\frac{n-7}{3}$ for cubic graphs.*

Proof. Since the contributions not linear in n are negative and we seek an upper bound, the bound holds also for disconnected n -vertex $(2r+1)$ -regular graphs. \square

4 Balloons and Matchings

Here we use balloons to prove the result of Henning and Yeo [9] minimizing the matching number for n -vertex $(2r+1)$ -regular connected graphs; in the next section we characterize the graphs where equality holds.

We use the Berge–Tutte Formula for the matching number. Recall that the deficiency $\text{def}(S)$ of a vertex set S in G is defined by $\text{def}(S) = o(G - S) - |S|$. Tutte [14] proved

that a graph G has a 1-factor if and only if $\text{def}(S) \leq 0$ for all $S \in V(G)$. The equivalent Berge–Tutte Formula (see Berge [3]) states that $\alpha'(G) = \min_{S \subseteq V(G)} \frac{1}{2}(n - \text{def}(S))$.

Lemma 4.1. *Let G be an n -vertex $(2r + 1)$ -regular graph, and let S be a subset of $V(G)$. If the number of edges from each odd component of $G - S$ to S is only 1 or is at least $2r + 1$, then $\text{def}(S) \leq \frac{2rb(G)}{2r+1}$.*

Proof. Let c_1 be the number of odd components of $G - S$ having one edge to S . By Lemma 3.3(a), each component of $G - S$ having one edge to S contains a balloon. Thus $c_1 \leq b(G)$. Counting the edges joining S to odd components of $G - S$ yields

$$(2r + 1)|S| \geq (2r + 1)o(G - S) - 2rc_1 \geq (2r + 1)o(G - S) - 2rb(G),$$

and hence $\text{def}(S) = o(G - S) - |S| \leq \frac{2rb(G)}{2r+1}$. □

Corollary 4.2. *If G is a connected cubic graph, then $\alpha'(G) \geq \frac{n}{2} - \left\lfloor \frac{b(G)}{3} \right\rfloor$.*

Proof. In a 3-regular graph, all edge-cuts between sets of odd size have odd size, which is 1 or at least 3. Hence Lemma 4.1 yields the claim (using the floor function in the second term is valid because $\alpha'(G)$ and $n/2$ are integers). □

If in a connected graph G some set of maximum deficiency satisfies the hypothesis of Lemma 4.1, then $\alpha'(G) \geq \frac{n}{2} - \frac{r}{2} \frac{(2r-1)n+2}{(2r+1)(2r^2+2r-1)}$, by the Berge–Tutte Formula and Lemma 3.1. We prove that this bound also holds for all connected odd-regular graphs. We already have a family \mathcal{H}_r where the bound is sharp; in the next section, we will show that these are the only graphs where equality holds.

Theorem 4.3. *If $G \in \mathcal{F}_{n,r}$, then $\alpha'(G) \geq \frac{n}{2} - \frac{r}{2} \frac{(2r-1)n+2}{(2r+1)(2r^2+2r-1)}$, with equality if $G \in \mathcal{H}_r$.*

Proof. In Proposition 2.2, we proved equality for graphs in \mathcal{H}_r ; here we prove the bound.

By the Berge–Tutte Formula, it suffices to show that every set $S \subseteq V(G)$ has deficiency at most $r \frac{(2r-1)n+2}{(2r+1)(2r^2+2r-1)}$. By Lemma 4.1, we may assume that there is an odd component of $G - S$ such that the number of edges from $G - S$ to S is between 3 and $2r - 1$; call such a component of $G - S$ a *bad subgraph*.

For each edge e joining S to a bad subgraph, replace e with a cut-edge incident to a copy of B_r at its end outside S . Also delete all vertices in bad subgraphs. Let G' denote the resulting graph; note that G' is $(2r + 1)$ -regular. Unfortunately, G' may be disconnected.

Let c be the number of bad subgraphs, and let x be the total number of vertices in them. Let y be the total number of edges in G joining S to bad subgraphs; y is the number of balloons added in forming G' .

Let p be the number vertices in some bad subgraph Q . If $p \leq 2r + 1$, then regularity forces each vertex of Q to have at least $2r + 2 - p$ neighbors in S . Hence the number of edges from S to $V(Q)$ is at least $p(2r + 2 - p)$, which is at least $2r + 1$, contradicting that Q is a bad subgraph. We conclude that $p \geq 2r + 3$, and hence $x \geq (2r + 3)c$.

The number of vertices in G' is $n - x + (2r + 3)y$. We also need the number of components of G' . Each time we pull an edge off a bad subgraph Q and make it incident to a copy of B_r , we increase the number of components by 0 or 1. Doing this with the last edge to Q (and deleting $V(Q)$) does not change the number of components. Since G is connected, we conclude that G' has at most $1 + y - c$ components.

The alteration from G to G' ensures that S satisfies the hypotheses of Lemma 4.1 for G' . Lemma 4.1 does not require connected graphs, so $\text{def}_{G'}(S) \leq \frac{2rb(G')}{2r+1}$. However, applying Lemma 3.1 to replace the number of balloons with upper bounds in terms of the number of vertices does require connected graphs. Therefore, we apply Lemma 3.1 to each component of G' . We obtain an additive constant 2 in the numerator for each component. Thus $b(G') \leq \frac{(2r-1)(n-x+(2r+3)y)+2(1+y-c)}{4r^2+4r-2}$. With $x \geq (2r + 3)c$, we have $b(G') \leq \frac{(2r-1)n+2}{4r^2+4r-2} + \frac{4r^2+4r-1}{4r^2+4r-2}(y-c)$.

Meanwhile, we must also relate $\text{def}_{G'}(S)$ to $\text{def}_G(S)$. We have replaced c odd components in $G - S$ with y odd components in $G' - S$. Thus

$$\begin{aligned} \text{def}_G(S) &= \text{def}_{G'}(S) - (y - c) \leq \frac{2rb(G')}{2r + 1} - (y - c) \\ &\leq \frac{r}{2r + 1} \frac{(2r - 1)n + 2}{2r^2 + 2r - 1} + \frac{2r}{2r + 1} \frac{4r^2 + 4r - 1}{4r^2 + 4r - 2} (y - c) - (y - c) \end{aligned}$$

Thus it suffices to show that $2r(4r^2 + 4r - 1) \leq (2r + 1)(4r^2 + 4r - 2)$. This inequality has the form $ab \leq (a + 1)(b - 1)$ with $a < b$, and hence it holds. \square

Corollary 4.4. *If G is a connected n -vertex cubic graph, then $\alpha'(G) \geq \frac{4n-1}{9}$, and this is sharp infinitely often.*

5 Characterization of Equality

Proposition 2.2 establishes equality in the bound of Theorem 4.3 for $G \in \mathcal{H}_r$. Now we show that these are the only graphs achieving equality. Recall that \mathcal{T}_r is the family of trees from which graphs in \mathcal{H}_r are formed by appending small balloons at leaves, and that the König–Egerváry Theorem states that if G is bipartite, then $\alpha'(G)$ equals the minimum number of vertices needed to “cover” the edges (by including at least one vertex from each edge).

Lemma 5.1. *If T is an n -vertex tree in which every non-leaf vertex has degree $2r + 1$, then $\alpha'(T) \geq \frac{n-1}{2r+1}$, with equality only when $T \in \mathcal{T}_r$.*

Proof. Since T has $n - 1$ edges and maximum degree $2r + 1$, the number of vertices needed to cover $E(T)$ is at least $\frac{n-1}{2r+1}$, and hence the König–Egerváry Theorem yields $\alpha'(T) \geq \frac{n-1}{2r+1}$.

If all leaves lie in the same partite set, then the other partite set is a vertex cover of size $\frac{n-1}{2r+1}$. Conversely, equality holding requires a vertex cover Q of size $\frac{n-1}{2r+1}$. No two vertices of Q can cover the same edge, so Q is an independent set. Also every vertex adjacent to a leaf must be in Q , since a leaf covers only one edge.

To show that all leaves are in the same partite set, let x and y be leaves, and let P be the path from x to y in T . The edges of P must be covered by vertices on P , so Q contains a vertex of each edge of P . Since Q is independent, the vertices of P alternate between Q and not- Q , with the neighbors of x and y being in Q . Hence the distance between x and y is even, and they are in the same partite set. \square

For a graph $G \in \mathcal{F}_{n,r}$ that achieves the minimum value of the matching number, we show that $G \in \mathcal{H}_r$ by showing that if we shrink each balloon to a single vertex, then the resulting graph is in \mathcal{T}_r .

Theorem 5.2. *If $G \in \mathcal{F}_{n,r}$ and $\alpha'(G) = \frac{n}{2} - \frac{r}{2} \frac{(2r-1)n+2}{(2r+1)(2r^2+2r-1)}$, then $G \in \mathcal{H}_r$.*

Proof. Equality in the bound requires equality in all the inequalities of Theorem 4.3. A set S with maximum deficiency must satisfy $\text{def}(S) = \frac{r}{2r+1} \frac{(2r-1)n+2}{2r^2+2r-1}$. Since the coefficient on $y - c$ in the final displayed inequality for Theorem 4.3 is negative, we must have $y = c$. This states that the total number of edges joining S to bad subgraphs equals the number of bad subgraphs, which implies that one edge goes to each bad subgraph, and therefore they are not bad. We conclude that $y = c = 0$, and the number of edges joining S to each odd component of $G - S$ is 1 or is at least $2r + 1$.

Now Lemma 4.1 applies and yields $\text{def}(S) \leq \frac{2rb(G)}{2r+1}$. From Lemma 3.1, we now have

$$\frac{r}{2r+1} \frac{(2r-1)n+2}{2r^2+2r-1} \leq \frac{2rb(G)}{2r+1} \leq \frac{r}{2r+1} \frac{(2r-1)n+2}{2r^2+2r-1},$$

so $b(G) = \frac{(2r-1)n+2}{4r^2+4r-2}$. From the proof of Lemma 3.1, equality in the bound requires each balloon to have exactly $2r + 3$ vertices.

Let G' be the graph obtained from G by shrinking each balloon to a single vertex. Let $n' = |V(G')|$ and $m' = |E(G')|$. Since each balloon has $2r + 3$ vertices, we have $n = n' + (2r + 2)b(G)$. Substituting this expression for n into the formula $b(G) = \frac{(2r-1)n+2}{4r^2+4r-2}$ and simplifying yields $2rb(G) = (2r - 1)n' + 2$.

Contraction does not disconnect, so G' is connected. To show that G' is a tree, we count the edges. By the degree sum formula,

$$2m' = (2r + 1)n' - 2rb(G) = (2r + 1)n' - (2r - 1)n' - 2 = 2n' - 2.$$

Finally, we show $G' \in \mathcal{T}_r$. By Lemma 5.1, it suffices to show that G' has a matching of size $\frac{n'-1}{2r+1}$. Note that $\alpha'(G') \geq \alpha'(G) - (r+1)b(G)$, and we are given $\alpha'(G) = \frac{n}{2} - \frac{rb(G)}{2r+1}$. Since $\frac{n}{2} - (r+1)b(G) = \frac{n'}{2}$ and $2rb(G) = (2r-1)n' + 2$, we conclude that $\alpha'(G') \geq \frac{n'-1}{2r+1}$. \square

6 Balloons and Total Domination

Balloons also help in proving bounds on the total domination number. The results are strongest for cubic graphs. We use a lemma proved by Henning that provides a useful upper bound in nearly regular graphs. Let $\Delta(G)$ and $\delta(G)$ denote the maximum and minimum vertex degrees in a graph G .

Lemma 6.1. (Henning's Lemma [7]) *If G is a graph with n vertices and m edges, where $n \geq 3$, then $\gamma_t(G) \leq n - \frac{m}{\Delta(G)}$.* \square

Lemma 6.2. *If B is a balloon with p vertices in a cubic graph G , then $\gamma_t(B) \leq \frac{p-1}{2}$. Furthermore, B has a dominating set of size $(p-1)/2$ that contains the neck of B and a neighbor of every vertex other than the neck.*

Proof. Let v be the neck of B . Recall that v has degree 2 in B , and the other vertices of B have degree 3 in B . By Henning's Lemma, $\gamma_t(B) \leq p - (3p-1)/6 = p/2 + 1/6$. Since p is odd and $\gamma_t(B)$ is an integer, $\gamma_t(B) \leq (p-1)/2$.

Let T be the set consisting of v and its two neighbors in B . The number of edges joining T and $V(B) - T$ is 2 or 4, depending on whether T induces a triangle. Note that $B - T$ has $p-3$ vertices and at least $\lceil 3(p-3)/2 \rceil$ edges. If $\Delta(B - T) = 3$, then Henning's Lemma yields $\gamma_t(B - T) \leq (p-3) - (3p-9-4)/6 = (p-3)/2 + 2/3$. Since p is odd and $\gamma_t(B - T)$ is an integer, $\gamma_t(B) \leq (p-3)/2$ in this case, and adding v to a smallest total dominating set of $B - T$ yields the desired set.

In the remaining case, $\Delta(B - T) < 3$. Since deleting T removes at most four edges incident to $V(B) - T$, this case requires $p \leq 7$. If $p = 7$, then $B - T = C_4$, and T is a total dominating set of size $(p-1)/2$ containing v . If $p = 5$, then B is the unique smallest balloon B_1 , and v with one of its neighbors forms a total dominating set of size $(p-1)/2$. \square

Example 6.3. When $|V(B)| = 7$, it may happen that B has no total dominating set of size $(p-1)/2$ containing its neck. This occurs in only one special balloon, which we denote by \hat{B} . This balloon consists a spanning cycle, say with vertices $v_0, v_1, v_2, v_3, v_4, v_5, v_6$ in order, where v_0 is the neck and the additional edges are v_1v_6, v_2v_4 , and v_3v_5 . Note that the neck induces a triangle with its neighbors, and this property determines \hat{B} completely. No total dominating set of size 3 contains the neck, because when the neck and a neighboring vertex lie in S , the undominated vertices induce a triangle having no neighbor in S ; thus adding a third vertex cannot complete a total dominating set.

In addition to small dominating sets, we also need large matchings in balloons.

Lemma 6.4. *Every balloon in a 3-regular graph has a matching that covers every vertex except its neck.*

Proof. Let v be the neck of a balloon B , with $N(v) = \{u, w\}$. Let B' consist of two disjoint copies of B plus a cut-edge joining their necks. Now B' is a 3-regular graph with one cut-edge, since B has no cut-edge.

Petersen proved that a 3-regular graphs with at most two cut-edges has a perfect matching. Since B has odd order, the cut-edge lies in every perfect matching. Deleting it leaves the desired matching in B . \square

Since $\alpha'(G) \geq \frac{n}{2} - \frac{b(G)}{3}$ when G is 3-regular and connected (Corollary 4.2), proving $\gamma_t(G) \leq \frac{n}{2} - \frac{b(G)}{2}$ would yield $\gamma_t(G) \leq \alpha'(G)$, with equality only when $b(G) = 0$. However, the desired upper bound may fail when G consists of three balloons plus one common neighbor.

The 2-edge-connected case (no balloons) has been well-studied. By Henning's Lemma, $\gamma_t(G) \leq n/2$. Equality may hold when G is 2-edge-connected; such graphs were characterized by Henning, Soleimanfallah, Thomassé, and Yeo [10]. The graphs achieving equality consist of two infinite families and one additional 16-vertex graph. In one family, the graph consists of two even cycles with vertex sets x_1, \dots, x_{2k} and y_1, \dots, y_{2k} , plus the edges $x_{2i-1}y_{2i}$ and $x_{2i}y_{2i-1}$ for $1 \leq i \leq k$. Being 2-edge-connected, these graphs also have perfect matchings, so here $\gamma_t(G) = \alpha'(G)$.

Hence we may confine our attention to graphs having balloons. Our strategy is to assemble a small total dominating set S using $(|V(B)| - 1)/2$ vertices in each balloon B and $|V(G')|/2$ vertices in the graph G' obtained by deleting the balloons. This gives the desired size. Vertices having neighbors in balloons have degree less than 3 in G' . Such a vertex in S does not need a neighbor in $S \cap V(G')$; Lemma 6.2 allows us to give it the neck of the balloon as a neighbor. This weakened restriction on S as a dominating set in G' motivates the following definition.

Definition 6.5. *A dominating set S in a graph G is a semitotal dominating set (abbreviated SD-set) if every vertex with maximum degree in G has a neighbor in S .*

In an SD-set, vertices of non-maximum degree can dominate themselves. The problem of finding an SD-set, like the problem of finding a total dominating set, can be modeled using hypergraphs. In the generalization of graphs to hypergraphs, any vertex set can form an edge; graphs are 2-uniform hypergraphs.

Definition 6.6. *A k -uniform hypergraph is a hypergraph in which every edge has size k . The transversal number $\tau(H)$ of a hypergraph H is the minimum size of a set of vertices that intersects every edge.*

For any graph, the total domination number equals the transversal number of the hypergraph on the same vertex set in which the edges are the vertex neighborhoods. An SD-set corresponds to a transversal when the edge of the hypergraph corresponding to a vertex v of non-maximum degree is its closed neighborhood (the neighborhood plus v itself). The theorem of Chvátal and McDiarmid on transversal number of k -uniform hypergraphs provides exactly what we need to find a sufficiently small SD-set in the graph obtained by deleting the balloons. (In [10], the Chvátal–McDiarmid result is used to explore the total domination numbers of regular graphs, noting that $\gamma_t(G) \leq n/2$ follows immediately for cubic graphs. Various special cases of the theorem are proved in [15] and [13], particularly for small k .)

Theorem 6.7. (Chvátal and McDiarmid [6]) *If H is a k -uniform hypergraph with n vertices and m edges, then $\tau(H) \leq \frac{\lfloor k/2 \rfloor m + n}{\lfloor 3k/2 \rfloor}$.*

We state the next two results for a graph G' because we will apply them when G' is the graph obtained from a 3-regular graph G by deleting the vertices in the balloons.

Corollary 6.8. *If G' is an n -vertex graph in which every vertex has degree $2r + 1$ or $2r$, then G' has an SD-set of size at most $\frac{(r+1)n}{3r+1}$.*

Proof. Form the hypergraph H with $V(H) = V(G')$ by letting the edges be the open neighborhoods of vertices with degree $2r + 1$ and the closed neighborhoods of vertices with degree $2r$. Thus H is a $(2r + 1)$ -uniform hypergraph with n vertices and n edges. By Theorem 6.7, $\tau(H) \leq \frac{(r+1)n}{3r+1}$. Every transversal of H is an SD-set in G' . \square

Using the plan we described above, Corollary 6.8 implies that $\gamma_t(G) \leq \frac{n}{2} - \frac{b(G)}{2}$ when $\Delta(G) = 3$ and no two balloons have a common neighbor. The remaining case will need special attention; here deleting the balloons leaves a vertex of degree 1.

Theorem 6.9. *If G' is a connected n -vertex graph with maximum degree at most 3, and $n > 1$, then G' has a dominating set S of size at most $n/2$ such that every vertex of degree 3 has a neighbor in S .*

Proof. The complement of any minimal dominating set is also dominating, so every n -vertex graph has a dominating set with at most $n/2$ vertices. Therefore, when $\Delta(G') < 3$ a smallest dominating set suffices. Hence we may assume that $\Delta(G') = 3$. The case $\Delta(G') < 3$ includes the basis step for induction on n .

If $\delta(G') \geq 2$, then Corollary 6.8 provides the desired SD-set. When G' has a vertex u of degree 1, let v be the neighbor of u . Let $F = G' - \{u, v\}$. If F has no isolated vertex, then we can apply the induction hypothesis to each component of F to obtain a set with the desired properties. Let T be the union of these sets; note that $|T| \leq (n - 2)/2$.

If v has degree 2, then F is connected, and $T \cup \{v\}$ is an SD-set in G' .

Suppose that v has degree 3. If v has no neighbor of degree 1 other than u , then F has no isolated vertices. Now $T \cup \{v\}$ is an SD-set in G' if T contains a neighbor of v , while otherwise $T \cup \{u\}$ is an SD-set.

In the remaining case, v has degree 3 and has another neighbor w of degree 1. In this case, let $F = G' - \{u, w\}$, and let T be the set in F guaranteed by the induction hypothesis (F is connected, since we only deleted vertices of degree 1). If $v \in T$, then $T \cup \{u\}$ is an SD-set in G' . Otherwise, T must contain the remaining neighbor of v to dominate v , and now $T \cup \{v\}$ is an SD-set in G' . \square

Theorem 6.10. *If G is a connected cubic graph with n vertices, then $\gamma_t(G) \leq \frac{n}{2} - \frac{b(G)}{2}$ (except that $\gamma_t(G) \leq n/2 - 1$ when $b(G) = 3$ and the three balloons have a common neighbor), and this is sharp for all even values of $b(G)$.*

Proof. Let G' be the graph obtained by deleting all vertices in balloons. If $G' = K_1$, then G consists of three balloons and their common neighbor. Lemma 6.2 yields a total dominating set in two of the balloons and a dominating set in the third that combine with one vertex of G' to yield $\gamma_t(G) \leq n/2 - 1$.

When G' has more than one vertex, we can apply Theorem 6.9 to obtain an SD-set S in G' . For each balloon B , let v be the neck. Use Lemma 6.2 to add a set S_B of size $(|V(B)| - 1)/2$. If the neighbor of v in $V(G')$ is in S , then choose S_B to be a set that contains v and contains a neighbor of every vertex in $V(B) - \{v\}$. If the neighbor of v in $V(G')$ is not in S , then simply choose S_B to be a total dominating set of B . After these contributions from all balloons, the size is at most $\frac{n}{2} - \frac{b(G)}{2}$.

If equality holds in the bound, then G' must have no SD-set of size less than $|V(G')|/2$. Let G' be formed from a cycle C_t by adding a pendant edge at each vertex. An SD-set in G' must use one vertex from each set consisting of a vertex of degree 1 and its neighbor.

We construct our example G by adding two 7-vertex balloons adjacent to each vertex of degree 1 in G' . Each such balloon is the special balloon \hat{B} in Example 6.3. The number of balloons is $2t$. Recall that \hat{B} has no total dominating set of size 3 that contains its neck. Therefore, if a total dominating set in G avoids some vertex u of degree 1 in G' , then the balloons adjacent to u contribute at least four vertices each, and the 16-vertex “wedge” containing them, u , and the neighbor of u in G' contributes at least eight vertices. Using u still requires it to contribute seven vertices, including three from each balloon. Thus we can save only 1 for each pair of balloons, and $\gamma_t(G) = \frac{n}{2} - \frac{b(G)}{2}$. \square

Corollary 4.2 and Theorem 6.10 together improve the inequality $\gamma_t(G) \leq \alpha'(G)$ for connected cubic graphs.

Corollary 6.11. *If G is a connected n -vertex cubic graph, then $\gamma_t(G) \leq \alpha'(G) - b(G)/6$, except when $b(G) = 3$ and there is exactly one vertex outside the balloons, in which case still $\gamma_t(G) \leq \alpha'(G)$.*

Proof. From the bounds in Corollary 4.2 and Theorem 6.10, it suffices to consider the exceptional case. Here $b(G) = 3$, and $\gamma_t(G) = n/2 - 1$ is possible. By Lemma 6.4, there are matchings in the balloon that cover all but the neck. One of the necks can be matched to their common neighbor, leaving only the two other necks as uncovered vertices. Hence $\alpha'(G) = n/2 - 1$ (equality holds, because deleting the vertex outside the balloons leaves three odd components). \square

The 3-regular case is the only case where the inequality between γ_t and α' is delicate. When more edges are added, α' tends to increase and γ_t tends to decrease, so the separation increases. For $(2r + 1)$ -regular graphs, applying the Chvátal–McDiarmid Theorem to the neighborhood hypergraph immediately yields $\gamma_t(G) \leq \frac{(r+1)n}{3r+1}$. On the other hand, $\alpha'(G) \geq \lfloor \frac{2}{n} \left(1 - \frac{2r-1}{2r+1} \frac{r}{2r^2+2r-1} \right) \rfloor$ (Theorem 4.3).

For larger r , balloons are not needed to separate $\gamma_t(G)$ and $\alpha'(G)$. Already when $r = 2$, the bound above yields $\gamma_t(G) \leq 3n/7 < 9n/22 < \alpha'(G)$. For large r , the results of Alon [1] on transversals of uniform hypergraphs yield the bound $\gamma_t(G) \leq n(1 + \ln k)/k$ when $\delta(G) \geq k$; refining this bound for k -regular graphs by using balloons does not seem worthwhile. Even for 4-regular graphs, the extremal problem for total domination is not completely solved. Yeo [16] conjectured that if G is a connected n -vertex graph with $\delta(G) \geq 4$ other than the bipartite complement of the Heawood graph, then $\gamma_t(G) \leq \frac{2}{5}n$.

Acknowledgment

We thank Alexandr Kostochka for pointing out a flaw in our original proof of Theorem 4.3. We thank Zoltán Füredi for pointing out the effectiveness of the Chvátal–McDiarmid Theorem in proving Corollary 6.8. We thank the referees for careful reading and comments.

References

- [1] N. Alon, Transversal numbers of uniform hypergraphs, *Graphs & Combin.* **6** (1990), 1–4.
- [2] D. Archdeacon, J. Ellis-Monaghan, D. Fischer, D. Fronček, P.C.B. Lam, S. Seager, B. Wei, and R. Yuster, Some remarks on domination. *J. Graph Theory* **46** (2004), 207–210.
- [3] F. Bäßler, Über die Zerlegung regulärer Streckenkomplexe ungerader Ordnung, *Comment. Math. Helvet.* **10** (1938), 275–287.

- [4] T. Biedl, E.D. Demaine, C.A. Duncan, R. Fleischer, and S.G. Kobourov, Tight bounds on maximal and maximum matchings, *Discrete Math* **285** (2004), 7–15.
- [5] G. Chartrand, S.F. Kapoor, L. Lesniak, and S. Schuster, Near 1-factors in graphs, *Proc. Fourteenth Southeastern Conference on Combinatorics, Graph Theory and Computing, Congress Numerantium* **41** (1984), 131–147.
- [6] V. Chvatal and C. McDiarmid, Small transversals in hypergraphs, *Combinatorica* **12** (1992), 19–26.
- [7] M.A. Henning, A linear Vizing-like relation relating the size and total domination number of a graph, *J. Graph Theory* **49** (2005), 285–290; erratum *Journal of Graph Theory* **54** (2007), 350–353.
- [8] M.A. Henning, L. Kang, E. Shan, and A. Yeo, On matching and total domination in graphs, *Discrete Math.* **308** (2008), 2313–2318.
- [9] M.A. Henning and A. Yeo, Tight lower bounds on the size of a maximum matching in a regular graph, *Graphs and Combinatorics* **23** (2007), 647–657.
- [10] M.A. Henning and A. Yeo, Hypergraphs with large transversal number and with edge sizes at least three, *J. Graph Theory* **59** (2008), 473–487.
- [11] Suil O and D.B. West, Generalized balloons and the Chinese Postman Problem in regular graphs. preprint (2008).
- [12] J. Petersen, Die Theorie der regulären Graphen, *Acta Math.* **15** (1891), 193–220.
- [13] S. Thomassé and A. Yeo, Total domination of graphs and small transversals of hypergraphs, *Combinatorica* **27** (2007), 473–487.
- [14] W.T. Tutte, The factorization of linear graphs, *J. Lond. Math. Soc.* **22** (1947), 107–111.
- [15] Z. Tuza, Covering all cliques of a graph, *Discrete Math.* **86** (1990), 117–126.
- [16] A. Yeo, Improved bound on the total domination in graphs with minimum degree four, manuscript (2005).