

Cycles in Color-Critical Graphs

Benjamin R. Moore*, Douglas B. West†

Revised January, 2021

Abstract

Tuza [1992] proved that a graph with no cycles of length congruent to 1 modulo k is k -colorable. We prove that if a graph G has an edge e such that $G - e$ is k -colorable and G is not, then for $2 \leq r \leq k$, the edge e lies in at least $\prod_{i=1}^{r-1} (k - i)$ cycles of length $1 \pmod r$ in G , and $G - e$ contains at least $\frac{1}{2} \prod_{i=1}^{r-1} (k - i)$ cycles of length $0 \pmod r$.

A (k, d) -coloring of G is a homomorphism from G to the graph $K_{k:d}$ with vertex set \mathbb{Z}_k defined by making i and j adjacent if $d \leq j - i \leq k - d$. When k and d are relatively prime, define s by $sd \equiv 1 \pmod k$. A result of Zhu [2002] implies that G is (k, d) -colorable when G has no cycle C with length congruent to is modulo k for any $i \in \{1, \dots, 2d - 1\}$. In fact, only d classes need be excluded: we prove that if $G - e$ is (k, d) -colorable and G is not, then e lies in at least one cycle with length congruent to $is \pmod k$ for some i in $\{1, \dots, d\}$. Furthermore, if this does not occur with $i \in \{1, \dots, d - 1\}$, then e lies in at least two cycles with length $1 \pmod k$ and $G - e$ contains a cycle of length $0 \pmod k$.

1 Introduction

One of the most fundamental results in graph theory is that graphs without odd cycles are 2-colorable. This has been generalized in many ways. Stong [12] proved that a graph is k -colorable if every vertex lies in fewer than $\binom{k}{2}$ odd cycles. Erdős and Hajnal [5] proved that a graph having no odd cycle of length more than $2j - 1$ is $2j$ -colorable. Tuza [14] proved that a graph having no cycle with length congruent to 1 modulo k is k -colorable. Tuza also strengthened Minty's Theorem [9] that a graph G is k -colorable if it has an orientation in which no cycle of G has more than $k - 1$ times as many forward edges as backward edges; Tuza showed that only cycles of length $1 \pmod k$ need be considered in that computation.

More recent work has provided guarantees not only for existence of cycles with certain lengths, but also lower bounds on the number of such cycles. A graph is $(k + 1)$ -critical

*Department of Combinatorics and Optimization, University of Waterloo, Waterloo, ON, Canada: brmoore@uwaterloo.ca. Research supported by the Natural Sciences and Engineering Research Council of Canada.

†Departments of Mathematics, Zhejiang Normal University, Jinhua, China, and University of Illinois, Urbana, IL: west@math.uiuc.edu. Research supported by National Natural Science Foundation of China grants NNSFC 11871439 and 11971439.

if it is not k -colorable but every proper subgraph is k -colorable. It is *doubly critical* if deleting the endpoints of any edge reduces the chromatic number by 2. In studying a weaker form of the Erdős–Lovász Tihany Conjecture that all doubly critical graphs are complete, Kawarabayashi, Pedersen, and Toft [8] showed that in a doubly critical graph with chromatic number $k + 1$, every edge lies in $\prod_{i=1}^{l-2} (k - i)$ cycles of length l . They used the technique of *generalized Kempe chains*, introduced as early as Neumann-Lara [10] and named in Toft [13]. Such a chain is a path following a particular list of colors.

Without the doubly-critical requirement, in this paper we obtain cycles in congruence classes rather than with specified lengths. We illustrate the generalized Kempe chain technique by first strengthening Tuza’s basic result, showing that for $2 \leq r \leq k$ every edge in a $(k + 1)$ -critical graph lies in at least $\prod_{i=1}^{r-1} (k - i)$ cycles of length congruent to 1 modulo r . The statement is more general.

Theorem 1. *For $2 \leq r \leq k$ and $e \in E(G)$, if $G - e$ is k -colorable and G is not, then e lies in at least $\prod_{i=1}^{r-1} (k - i)$ cycles of length congruent to 1 modulo r .*

Thus every graph with fewer than $(k - 1)!$ cycles of length congruent to 1 modulo k is k -colorable.

We also apply our technique to the more general problem of H -coloring. A *homomorphism* from a graph G to a graph H is a map $\phi: V(G) \rightarrow V(H)$ such that $uv \in E(G)$ implies $\phi(u)\phi(v) \in E(H)$. A homomorphism into H is also called an H -*coloring*. A proper k -coloring is simply a K_k -coloring, where K_k is the complete graph with k vertices. The H -coloring problem becomes more complicated when H is not complete because there are more ways for a coloring to violate an edge. An ordinary proper coloring requires colors on adjacent vertices to be distinct, but this is no longer enough.

The *circular clique* $K_{k;d}$ has vertex set \mathbb{Z}_k and edge set $\{ij: d \leq j - i \leq k - d\}$. The complete graph K_k is simply the circular clique $K_{k;1}$. Homomorphism into $K_{k;d}$ is called (k, d) -*coloring*, and a graph having a (k, d) -coloring is (k, d) -*colorable*. When G has an edge, (k, d) -coloring requires $k \geq 2d$. The *circular chromatic number* χ_c of a graph G is the least k/d such that G is (k, d) -colorable. In particular, if $k'/d' \leq k/d$, then $K_{k';d'}$ is $K_{k;d}$ -colorable. See Zhu [15, 17] for surveys on this topic.

Zhu [16] extended Tuza’s result to circular coloring. Given an orientation of a graph G , and given a cycle C in G viewed in a consistent direction, let C^- denote the set of edges in C oriented oppositely to their orientation of G . Zhu proved that if G has an orientation such that $|E(C)|/|C^-| \leq k/d$ for every cycle C (in each direction) such that $d|E(C)|$ is congruent modulo k to some value in $\{1, \dots, 2d - 1\}$, then G is (k, d) -colorable.

When d and k are relatively prime, let s be the congruence class such that $sd \equiv 1 \pmod{k}$. It follows from Zhu’s result that if G has no cycle with length congruent to is modulo k for any i with $1 \leq i \leq 2d - 1$, then G is (k, d) -colorable. That is, (k, d) -colorability holds when

$2d - 1$ congruence classes of cycle lengths modulo k are forbidden. Our result implies that it suffices to exclude $i \in \{1, \dots, d\}$. In particular, if $G - e$ is $K_{k:d}$ -colorable and G is not, then e lies in a cycle with length congruent to $is \pmod k$ for some i in $\{1, \dots, d\}$. Note that $ds \equiv 1 \pmod k$. We further show that if G has no cycle through e with length $is \pmod k$ when $i \in \{1, \dots, d - 1\}$, then e lies in at least two cycles with length $1 \pmod k$ and $G - e$ contains a cycle of length $0 \pmod k$. (Note that (k, d) -coloring forbids loops.)

The special case of $K_{(2d+1):d}$ -coloring shows that our result is sharp. Here $-2d \equiv 1 \pmod{(2d+1)}$, so $s = -2$, and we seek a cycle length congruent to $-2i$ for some $i \in \{1, \dots, k\}$. These lengths are the odd values from $2d - 1$ to 1 . The graph $K_{(2d+1):d}$ is isomorphic to the odd cycle C_{2d+1} . Thus $\chi_c(G) \leq 2 + 1/d$ for a C_{2d+1} -colorable graph G . All shorter odd cycles are critical non- C_{2d+1} -colorable graphs (we can also view a single vertex with a loop as a degenerate such example). This means that although possibly only one cycle length among the listed classes of lengths occurs, we cannot omit any of those classes from the list.

Returning to the topic of proper k -coloring, there are further questions to ask. An edge in the complete graph K_{k+1} lies in exactly $(k - 1)!$ cycles of length $1 \pmod k$. When r is at most $k/2$, the guarantee of $\prod_{i=1}^{r-1} (k - i)$ cycles of length $1 \pmod r$ is not sharp in K_{k+1} , because cycles of length $r + 1$ and cycles of length $2r + 1$ both count.

Question 1. *Can the guarantee of $\prod_{i=1}^{r-1} (k - i)$ cycles of length $1 \pmod r$ through each edge be sharp for non-complete $(k + 1)$ -critical graphs when $k/2 < r < k$?*

We can also consider other congruence classes. In fact, K_{k+1} has cycles in all congruence classes modulo r for $2 \leq r \leq k$ except for the class of $2 \pmod k$. In a paper on cycle lengths in directed graphs, Chen, Ma, and Zang [2] proved that for integers l and k with $1 \leq l \leq k$ and $k \geq 2$, a graph containing no cycle of length congruent to $l \pmod k$ is k -colorable if $l \neq 2$, and is $(k + 1)$ -colorable if $l = 2$. This proved a strong form of a conjecture of Diwan, Kenkre, and Vishwanathan [4] and answered a question asked by Tuza [14].

A recent paper of Gao, Huo, Lui, and Ma [6] resolves almost all the existence questions in a very strong way. Stated in our language, they proved that a non- k -colorable graph contains $k - 1$ cycles of consecutive lengths. This covers all but one congruence class modulo k and all congruence classes for smaller moduli. Non- k -colorable graphs have $(k + 1)$ -critical subgraphs, which have minimum degree at least k ; they proved also that 3-connected nonbipartite graphs with minimum degree at least k have cycles with $k - 1$ consecutive lengths. Also, minimum degree at least k guarantees cycles of all even lengths modulo $k - 1$, extending to all lengths modulo $k - 1$ in the 2-connected nonbipartite case. Finally, for $k \geq 3$ every k -connected graph has a cycle whose length is a multiple of k . These results were variously conjectured by Sudakov and Verstraëte, by Bondy and Vince, by Thomassen, and by Dean. In [6] they are proved by a unified approach, but the argument is quite long.

A subsequent paper by Gao, Huo, and Ma [7] resolved the remaining question about $2 \bmod k$, proving that when $k \geq 6$ every non- k -colorable graph not having K_{k+1} as a block contains k cycles of consecutive lengths. Hence it has cycles in all congruence classes with modulus at most k . The case $2 \bmod k$ when $k = 3$ follows from a combination of results of Saito [11] and Dean, Kaneko, Ota, and Toft [3]. For $k \in \{4, 5\}$, the authors of [7] state that their method works but yields a proof that was too long to include. Their result culminates a long series of conjectures and theorems on cycle lengths in color-critical graphs by many researchers; see [7] for the history and further references.

One may also wonder whether the cycle length guarantees follow from weaker hypotheses. Since $(k + 1)$ -critical graphs are k -edge-connected, one may wonder whether being k -chromatic and critically k -edge-connected is enough. Already this fails when $r = k$ and we ask for just one cycle. The Petersen graph is 3-colorable (but not 3-critical) and is critically 3-edge-connected. However, it has no 4-cycle, no 7-cycle, and no 10-cycle, so it has no cycle of length congruent to 1 modulo 3.

2 Proper Coloring

We consider the cycles forced when deletion of an edge reduces the chromatic number.

Theorem 2. *Fix $r, k \in \mathbb{N}$ with $2 \leq r \leq k$, and let e be an edge in a graph G . If $G - e$ is k -colorable and G is not, then e belongs to at least $\prod_{i=1}^{r-1} (k - i)$ cycles in G having lengths congruent to 1 modulo r .*

Proof. Let $[k] = \{1, \dots, k\}$. Fix a proper k -coloring ϕ of $G - e$ with colors in $[k]$. Let x and y be the endpoints of e . We obtain a cycle through e for each cyclic list of r members of $[k]$ containing $\phi(x)$. Starting with $\phi(x)$, the cyclic list σ can be formed in $\prod_{i=1}^{r-1} (k - i)$ ways.

Given such σ , define the σ -subdigraph of G generated by ϕ to be the digraph D_σ with vertex set $V(G)$ such that uv is an edge in D_σ if and only if $uv \in E(G)$ and $\sigma(\phi(u)) = \phi(v)$. Let F be the subdigraph of D_σ induced by all vertices reachable from x by paths in D_σ .

Define a recoloring ϕ' of $G - e$ by $\phi'(u) = \sigma(\phi(u))$ for $u \in V(F)$ and $\phi'(u) = \phi(u)$ for $u \in V(G) - V(F)$. An edge is improperly colored by ϕ' only if the color of one endpoint remains fixed and the other changes into it, but then the oriented version of the edge lies in F and both endpoints change color. Thus ϕ' is a proper k -coloring of $G - e$.

Since G is not k -colorable, $\phi(x) = \phi(y)$. Also $\phi'(x) \neq \phi(x)$. If $y \notin V(F)$, then we have $\phi'(x) \neq \phi(x) = \phi(y) = \phi'(y)$, and ϕ' is a proper coloring of G , which by hypothesis does not exist. Hence $y \in V(F)$, meaning that y is reachable from x via a path in F .

Since paths in F follow colors according to σ , and $\phi(y) = \phi(x)$, the length of any x, y -path in F is a multiple of r , and the cycle in G completed by adding the edge yx has length congruent to 1 modulo r . Furthermore, since the coloring ϕ is fixed, the resulting x, y -paths in G are distinct for distinct choices of σ . Hence we obtain $\prod_{i=1}^{r-1} (k - i)$ cycles through e . \square

Our result was motivated by a similar quantitative argument due to Brewster, McGuinness, Moore, and Noel [1], which we describe in our language to generalize it. For $k > 2$, they showed that if G is not k -colorable but $G - xy$ is k -colorable, then $G - xy$ contains at least $(k - 1)!/2$ cycles with lengths divisible by k . The early paper of Tuza [14] notes that Toft and Tuza had observed for $k > 2$ that every non- k -colorable graph contains a cycle whose length is divisible by k .

Theorem 3. *For $3 \leq r \leq k$, if a graph G is not k -colorable but $G - e$ is k -colorable, where $e \in E(G)$, then $G - e$ contains at least $\frac{1}{2} \prod_{i=1}^{r-1} (k - i)$ cycles whose lengths are divisible by r , none of which contain e .*

Proof. Let x and y be the endpoints of e . Again fix a proper k -coloring ϕ of $G - e$ and a cyclic permutation σ of a set of r colors containing $\phi(x)$. Define the digraph D_σ as above. Note again that $\phi(x) = \phi(y)$, since G is not k -colorable.

If D_σ is acyclic, then we recolor G by again changing the color on v from $\phi(v)$ to $\sigma(\phi(v))$, but this time we perform the change *one vertex at a time*, always changing the color at a sink of the unchanged subgraph. At each step we have a proper k -coloring of $G - e$. If at some point the color on x or y changes, then we have produced a proper k -coloring of G . Since G has no such coloring, D_σ must contain a cycle. Since $\phi(x) = \phi(y)$, the edge e does not appear in D_σ , so such cycles do not contain e .

The length of any cycle in D_σ is a multiple of r . However, a cyclic permutation and its reverse will select the same cycle in G , because the corresponding digraphs are obtained from each other by reversing all the edges. Hence we are in fact guaranteed $\frac{1}{2} \prod_{i=1}^{r-1} (k - i)$ cycles whose lengths are multiples of r , and none of these cycles contain e . \square

We have guaranteed $(k - 1)!$ cycles of length $1 \pmod k$ in a $(k + 1)$ -critical graph. We next present a probabilistic argument that guarantees more, suggested by a referee. Although $k!/2$ is greater than $(k - 1)!$, the probabilistic argument does not yield $k!/2$ such cycles *through every edge* in a $(k + 1)$ -critical graph, so it does not directly imply that conclusion.

Theorem 4. *A graph that is not k -colorable has at least $k!/2$ cycles with length congruent to 1 modulo k , with equality only when those cycles all have length exactly $k + 1$.*

Proof. Randomly order the vertices and orient each edge toward its later endpoint in the order. Now, bound the probability that some cycle of length $1 \pmod k$ (or its reverse) has more than $k - 1$ times as many forward edges as backward edges. If this probability is at most $2/k!$ and there are fewer than $k!/2$ such cycles, then some orientation has no cycle of length $1 \pmod k$ with more than $k - 1$ times as many forward edges as backward edges. By Tuza's strengthening of Minty's Theorem, G is then k -colorable.

A cycle C of length $qk + 1$ is bad if following the vertices along C involves at most q backward steps and more than $(k - 1)q$ forward steps in the ordering. Since the vertices

outside the cycle are irrelevant, it suffices to show that at most $(qk + 1)!/k!$ of the $(qk + 1)!$ orderings of v_1, \dots, v_{qk+1} have at most q instances of v_i preceding v_{i-1} .

From one backward step to the next is an increasing run. Hence to form an ordering with at most q backward steps we assign the vertices to bins 1 through q and place the vertices within a bin in increasing order. Such assignments include the orderings where there are fewer backwards steps. Since we are following a cycle, we can start at any of the $qk + 1$ vertices in the resulting linear order. The probability is thus bounded by $q^{qk+1}/(qk)!$. Multiplying by 2 accounts for following the cycle in the opposite order.

The bound is at most $2/k!$, with equality only when $q = 1$. This yields the additional observation that if a non- k -colorable graph has only $k!/2$ cycles of length congruent to 1 modulo k , then those cycles all must have length exactly $k + 1$. This occurs for example in K_{k+1} , so the result is sharp. \square

3 Circular Coloring

In this section we consider the analogous problem for (k, d) -coloring. We will only use color cycles of the form $(0, d, 2d, \dots, -d)$ and their reverse, so we get existence results rather than quantitative results. Nevertheless, they are sharp in terms of the number of classes allowed, as discussed in the introduction.

The proof may require many steps of recoloring to find a desired cycle. This is inherently necessary, because a C_{2d+1} -coloring of $C_{2d-1} - e$ may alternate 0 and d along the path.

Theorem 5. *Given k and d relatively prime with $k > 2d$, let s be the element of \mathbb{Z}_k such that $sd \equiv 1 \pmod k$. For an edge e in a graph G , if $G - e$ is $K_{k,d}$ -colorable and G is not, then e lies in a cycle in G of length congruent to $is \pmod k$ for some i in $\{1, \dots, d\}$.*

Proof. Fix a $K_{k,d}$ -coloring ϕ of $G - e$. Let x and y be the endpoints of e . By cyclic symmetry, we may assume $\phi(y) = 0$. Since G is not $K_{k,d}$ -colorable, $\phi(x) \in \{0, \pm 1, \dots, \pm(d-1)\}$. Let σ be the cyclic permutation $(0, d, 2d, \dots, -d)$ of colors. Define the digraph D_σ as in Theorem 2, and let F be the subdigraph of D_σ induced by all vertices reachable from x in D_σ .

Given ϕ , define ϕ' on $G - e$ by letting $\phi'(v) = \phi(v) + 1$ for $v \in V(F)$ and $\phi'(v) = \phi(v)$ for $v \notin V(F)$. We claim that ϕ' is a $K_{k,d}$ -coloring of $G - e$. First, edges within F or in $G - V(F)$ remain properly colored. When $v \in V(F)$, the exploration of D_σ extends along the edge vw if $\phi(w) - \phi(v) = d$. Since $\phi(w) - \phi(v) \in \{d, d+1, \dots, k-d\}$ for $vw \in E(G - e)$, having $v \in V(F)$ and $w \notin V(F)$ requires $\phi(w) - \phi(v) \in \{d+1, \dots, k-d\}$. Now $\phi'(w) - \phi'(v) \in \{d, \dots, k-d-1\}$, so such edges are also properly colored in ϕ' .

We will consider cases where $\phi(x) = j$, for $0 \leq j \leq d-1$. For the case $\phi(x) = -j$ with $1 \leq j \leq d-1$, add j to the color at each vertex to obtain $\phi(x) = 0$ and $\phi(y) = j$, and then interchange the roles of x and y and apply the argument below.

When $\phi(x) = j$ and $\phi(y) = 0$, we claim that G has a cycle through xy with length congruent to is modulo k for some i in $\{1, \dots, d - j\}$. Note first that if F has an x, y -path of length r , then $rd \equiv -j \pmod{k}$, since each edge increases the color value by d . Multiplying by s yields $r \equiv -js \pmod{k}$. Since $sd \equiv 1$, adding 1 to each side to compute the length of the cycle yields $r + 1 \equiv -js + ds \equiv (d - j)s \pmod{k}$.

We now prove the claim by induction on $d - j$. First consider $j = d - 1$. If $y \notin V(F)$, then ϕ' is a $K_{k:d}$ -coloring of G , since $\phi'(x) = d$ and $\phi'(y) = 0$. Hence $y \in V(F)$. Now by the computation above we have a cycle through yx with length congruent to $1s \pmod{k}$.

Now suppose $j < d - 1$. If $y \in V(F)$, then the computation yields a cycle through yx with length congruent to $(d - j)s \pmod{k}$. Hence we may assume $y \notin V(F)$. Now ϕ' is a $K_{k:d}$ -coloring of $G - xy$ with $\phi'(x) = j + 1$ and $\phi'(y) = 0$. The induction hypothesis, applied to ϕ' with $\phi'(x) = j + 1$, now implies that G has a cycle through xy with length congruent to $is \pmod{k}$ for some i in $\{1, \dots, d - j - 1\}$. Including $d - j$ in the set thus covers all cases to complete the induction step. \square

Proposition 6. *If in the setting of Theorem 5, e does not lie in a cycle with length congruent to $is \pmod{k}$ for some i in $\{1, \dots, d - 1\}$, then e lies in at least two cycles of length $1 \pmod{k}$ and $G - e$ contains a cycle of length $0 \pmod{k}$.*

Proof. With the cases $1 \leq i \leq d - 1$ excluded in the argument of Theorem 5, the remaining case is $j = 0$ and $y \in V(F)$. To reach y from x along steps of value $+d$, the number of steps must be a multiple of k , since $\phi(x) = \phi(y)$, and adding e completes a cycle. Under σ^{-1} , using the same (k, d) -coloring ϕ of $G - e$ and starting again from x yields a second cycle of length $1 \pmod{k}$ through xy .

Furthermore, if $G - e$ has no cycle of length $0 \pmod{k}$, then D_σ is acyclic. Working backward from sinks, we can add 1 to the color of each reached vertex, *one vertex at a time*, always maintaining a (k, d) -coloring of $G - e$, until x or y changes color. This reduces the problem to the case $j > 0$. Since in this case G has no cycle of length is with $i \in \{1, \dots, d - 1\}$, the previous arguments produce a (k, d) -coloring of G , which by hypothesis does not exist. Therefore, in fact $G - e$ also contains a cycle of length $0 \pmod{k}$. \square

In these arguments, we have not used cycles in $K_{k:d}$ other than that generated by d or $-d$. When $k = 2d + 1$, these two are the only permutations yielding cycles in the host graph, and that is why our sharpness examples in the introduction are for C_{2d+1} -coloring. In that case $s = -2$. The set of cycle lengths that cannot be avoided are the congruence classes $-2i \pmod{(2d + 1)}$ for $1 \leq i \leq d$, and if there are no cycles through e in the classes with $1 \leq i \leq d - 1$, then we obtain two cycles with lengths $1 \pmod{(2d + 1)}$.

Other cycles in the host graph can yield other sets of forced cycles, but the key is designing a recoloring that produces another H -coloring of $G - e$.

References

- [1] R. C. Brewster, S. McGuinness, B. Moore, and J. A. Noel, A dichotomy theorem for circular colouring reconfiguration. *Theoret. Comput. Sci.* 639 (2016), 1–13.
- [2] Z. Chen, J. Ma, and W. Zang, Colouring digraphs with forbidden cycles. *J. Combin. Theory Ser. B* 115 (2015), 210–223.
- [3] N. Dean, A. Kaneko, K. Ota and B. Toft, Cycles modulo 3. *DIMACS Technical Report* 91-32 (1991).
- [4] A. A. Diwan, S. Kenkre, and S. Vishwanathan, Circumference, chromatic number and online coloring. *Combinatorica* 33 (2013) 319–334.
- [5] P. Erdős and A. Hajnal, On chromatic number of graphs and set-systems. *Acta Math. Acad. Sci. Hungar.* 17 (1966), 61–99.
- [6] J. Gao, Q. Huo, C. Liu, and J. Ma, A unified proof of conjectures on cycle lengths in graphs. *arXiv:1904.08126* 2019, *Intl. Math. Res. Notices* (to appear).
- [7] J. Gao, Q. Huo, and J. Ma, A strengthening on odd cycles in graphs of given chromatic number. *arXiv:2012.10624* 2020.
- [8] K. Kawarabayashi, A. Pedersen, and B. Toft, Double-critical graphs and complete minors, *Electron J. Combin.* 17 (2010), Paper #R87.
- [9] G. J. Minty, A theorem on n -coloring the points of a linear graph. *Amer. Math. Monthly* 69 (1962), no. 7, 623–624.
- [10] V. Neumann-Lara, The dichromatic number of a digraph. *J. Combin. Theory Ser. B* 33 (1982), no. 3, 265–270.
- [11] A. Saito, Cycles of length 2 modulo 3 in graphs. *Discrete Mathematics* 101 (1992), no. 1–3, 285–289.
- [12] R. Stong, Solution to problem 11086. *Amer. Math. Monthly* 113 (2006), 372. (Proposed in Volume 111 (2004), 440.)
- [13] B. Toft, Colouring, stable sets and perfect graphs. In *Handbook of Combinatorics, Vol. 1* (Elsevier, 1995), 233–288.
- [14] Zs. Tuza, Graph coloring in linear time. *J. Combin. Theory Ser. B* 55 (1992), no. 2, 236–243.
- [15] X. Zhu, Circular chromatic number: a survey. In “Combinatorics, graph theory, algorithms and applications,” *Discrete Math.* 229 (2001), no. 1-3, 371–410.
- [16] X. Zhu, Circular colouring and orientation of graphs. *J. Combin. Theory Ser. B* 86 (2002), no. 1, 109–113.
- [17] X. Zhu, Recent developments in circular colouring of graphs. In “Topics in discrete mathematics,” *Algorithms Combin.* 26, (Springer, 2006), 497–550.