

The vulnerability of the diameter of enhanced hypercubes

Meijie Ma^{1,4} Douglas B. West^{1,2} Jun-Ming Xu³

Abstract

For an interconnection network G , the ω -wide diameter $d_\omega(G)$ is the least ℓ such that any two vertices are joined by ω internally-disjoint paths of length at most ℓ , and the $(\omega - 1)$ -fault diameter $D_\omega(G)$ is the maximum diameter of a subgraph obtained by deleting fewer than ω vertices of G .

The enhanced hypercube $Q_{n,k}$ is a variant of the well-known hypercube. Yang, Chang, Pai, and Chan gave an upper bound for $d_{n+1}(Q_{n,k})$ and $D_{n+1}(Q_{n,k})$ and posed the problem of finding the wide diameters and fault diameters of $Q_{n,k}$. By constructing internally disjoint paths between any two vertices in the enhanced hypercube, for $n \geq 3$ and $2 \leq k \leq n$ we prove

$$D_\omega(Q_{n,k}) = d_\omega(Q_{n,k}) = \begin{cases} d(Q_{n,k}) & \text{for } 1 \leq \omega < n - \lfloor \frac{k}{2} \rfloor; \\ d(Q_{n,k}) + 1 & \text{for } n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1. \end{cases}$$

where $d(Q_{n,k})$ is the diameter of $Q_{n,k}$. These results mean that interconnection networks modelled by enhanced hypercubes are extremely robust.

Keywords: interconnection network; enhanced hypercube; wide diameter; fault diameter.

1 Introduction

An interconnection network is conveniently represented by an undirected graph. The vertices (or edges) of the graph represent the nodes (or links) of the network. Throughout this paper, vertex and node, edge and link, graph and network are used interchangeably. Reliability and efficiency are important criteria in the design of interconnection networks. In the study of fault-tolerance and transmission delay of networks, wide diameter and fault diameter are important parameters that have been studied by many researchers. They combine connectivity with diameter to measure simultaneously the fault-tolerance and efficiency of parallel processing computer networks.

¹Department of Mathematics, Zhejiang Normal University, Jinhua 321004, China.

²Department of Mathematics, Univ. of Illinois, Urbana, IL 61801, USA; dwest@math.uiuc.edu.

³School of Mathematical Sciences, University of Science and Technology of China, Wentsun Wu Key Laboratory of CAS, Hefei 230026, China.

⁴Corresponding author: mameij@mail.ustc.edu.cn.

These parameters were studied by several authors for some Cartesian product graphs [2, 3, 4, 28, 29] and for the hypercube and its variants [6, 7, 8, 10, 11, 13, 20, 21, 22, 23].

Let u and v be two vertices in a network G . A u, v -path is a path with endpoints u and v . The *distance* between u and v , denoted by $d(u, v)$, is the minimum length (number of edges) of a u, v -path. The *diameter* of G , denoted by $d(G)$, is the maximum distance between vertices. The *connectivity* $\kappa(G)$ is the minimum number of vertices whose removal results in a disconnected or 1-vertex network. We say that G is k -connected when $0 < k \leq \kappa(G)$. By Menger's Theorem [18], in a k -connected network there exist k internally disjoint paths joining any two vertices (*internally disjoint* means that the only shared vertices are the endpoints).

Given a k -connected graph G , fix ω with $1 \leq \omega \leq k$. The ω -wide diameter of G , denoted by $d_\omega(G)$, is the least ℓ such that for any $u, v \in V(G)$ there exist ω internally disjoint u, v -paths of length at most ℓ . Throughout this paper, we abuse terminology by writing "disjoint paths" to mean "internally disjoint paths". Note that $d_1(G)$ is just the diameter $d(G)$ of G . From the definition,

$$d(G) = d_1(G) \leq d_2(G) \leq \dots \leq d_{k-1}(G) \leq d_k(G).$$

Failures are inevitable when a network is put in use. Therefore, it is important to consider faulty networks. The $(\omega - 1)$ -fault diameter of a graph G , denoted by $D_\omega(G)$, is the maximum diameter among subgraphs obtained from G by deleting fewer than ω vertices; it measures the worst-case effect on the diameter when vertex faults occur. Note that $D_\omega(G)$ is well-defined if and only if G is ω -connected, moreover,

$$d(G) = D_1(G) \leq D_2(G) \leq \dots \leq D_{k-1}(G) \leq D_k(G).$$

From the definitions, it follows that $D_\omega(G) \leq d_\omega(G)$ when G is k -connected and $1 \leq \omega \leq k$ [15]. Equality holds for some well-known networks [9, 14].

As a topology for an interconnection network of a multiprocessor system, the hypercube is a widely used and well-known model, since it possesses many attractive properties such as regularity, symmetry, logarithmic diameter, high connectivity, recursive construction, ease of bisection, and relatively low link complexity [12, 19, 27]. We study an important variant of the hypercube Q_n , the enhanced hypercube $Q_{n,k}$ proposed by Tzeng and Wei [24]; its properties have been studied in [5, 16, 25, 26, 30]. We give the definition and basic properties of $Q_{n,k}$ in Section 2.

It was shown by Liu [16] that $\kappa(Q_{n,k}) = n + 1$. Thus, the wide diameter $d_\omega(Q_{n,k})$ and the fault diameter $D_\omega(Q_{n,k})$ are well-defined when $\omega \leq n + 1$. Yang, Chang, Pai, and Chan [30] gave an upper bound for $d_{n+1}(Q_{n,k})$ and $D_{n+1}(Q_{n,k})$, and they posed the problem of finding the wide diameter and fault diameter of $Q_{n,k}$. In this paper, for $n \geq 3$ and $2 \leq k \leq n$, we prove

$$D_\omega(Q_{n,k}) = d_\omega(Q_{n,k}) = \begin{cases} d(Q_{n,k}) & \text{for } 1 \leq \omega < n - \lfloor \frac{k}{2} \rfloor; \\ d(Q_{n,k}) + 1 & \text{for } n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1. \end{cases}$$

The special case $k = n$ (folded hypercube) was obtain earlier by Simó and Yebra [21], along with the same values for edge deletions. For enhanced hypercubes also, our arguments yield the same values for edge deletions as for vertex deletions.

2 Properties of $Q_{n,k}$

Let $x_n \cdots x_1$ be an n -bit binary string. We call the rightmost bit the first bit and the leftmost bit the n th bit. For simplicity we use a^i to mean that the bit a is repeated i times; for example, $01^30^2 = 011100$. The *Hamming distance* between strings u and v , denoted by $H(u, v)$, is the number of positions where the two strings differ.

The n -dimensional hypercube Q_n is the graph whose vertices are the n -bit binary strings and whose edges are the pairs of vertices differing in exactly one position. An edge of Q_n is a j -dimensional edge if the two endpoints differ in the j th position. For $1 \leq j \leq n$, let E_j denote the set of j -dimensional edges in Q_n .

As a variant of the hypercube, the n -dimensional folded hypercube FQ_n , proposed first by El-Amawy and Latifi [1], is obtained from the hypercube Q_n by making each vertex u adjacent to its complementary vertex, denoted \bar{u} and obtained from u by subtracting each bit from 1. Such an edge is often called a *complementary edge*.

For $2 \leq k \leq n$, the n -dimensional enhanced hypercube $Q_{n,k}$ is obtained from the hypercube Q_n by adding the edge uv whenever u and v are related by $u = x_n \cdots x_1$ and $v = x_n \cdots x_{k+1} \bar{x}_k \bar{x}_{k-1} \cdots \bar{x}_1$; that is, the first k bits are complemented. Such an edge is called a k -complementary edge. For convenience, we use E_0 to denote the set of k -complementary edges. Thus $E(Q_{n,k}) = E(Q_n) \cup E_0$. When $k = n$, we have $Q_{n,n} = FQ_n$; hence the enhanced hypercube is a generalization of the folded hypercube. The graphs shown in Fig. 1 are $Q_{3,3}$ and $Q_{4,3}$, where the hypercube edges and 3-complementary edges are represented by solid lines and dashed lines, respectively.




Figure 1: Enhanced hypercubes $Q_{3,3}$ and $Q_{4,3}$

A graph G is *vertex-transitive* if for any $u, v \in V(G)$ there is some $\sigma \in \text{Aut}(G)$, the automorphism group of G , such that $\sigma(u) = v$; it is *edge-transitive* if for any $xy, uv \in E(G)$ there is some $\sigma \in \text{Aut}(G)$ such that $\{\sigma(x), \sigma(y)\} = \{u, v\}$. The hypercube Q_n and folded hypercube FQ_n are vertex-transitive and edge-transitive, and the enhanced hypercube $Q_{n,k}$ is vertex-transitive but not edge-transitive when $k < n$ [17, 27, 30].

The *Cartesian product* $G \square H$ of graphs G and H is the graph with vertex set $V(G) \times V(H)$, in which vertices (u, v) and (u', v') are adjacent whenever $uu' \in E(G)$ and $v = v'$, or $u = u'$ and $vv' \in E(H)$. By the definition of $Q_{n,k}$, we have $Q_{n,k} = Q_{n-k} \square FQ_k$. Although $Q_{n,k}$ is not edge-transitive when $2 \leq k < n$, we have the following property.

Proposition 1. *Permuting the first k positions and/or permuting the last $n - k$ positions in the names of the vertices of $Q_{n,k}$ does not change the graph.*

Proof. Exchanging the i th bit and j th bit among the first k or among the last $n - k$ preserves the adjacency relation, since the number of coordinates in which two vertices differ is not changed by exchanging such coordinates. An arbitrary permutation is obtained by a succession of such exchanges. \square

Proposition 2. *Given $u, v \in V(Q_{n,k})$, let r and s be the numbers of positions in which u and v differ among the first k and last $n - k$ positions, respectively. The distance between u and v is computed by $d(u, v) = s + \min\{r, k - r + 1\}$.*

Proof. The distance is the minimum number of steps to change u into v . All steps change one bit, except that a k -complementary edge changes the first k bits. If no k -complementary edge is used, then the number of steps is at least the Hamming distance, and this suffices. For this reason, a shortest path uses at most one k -complementary edge. If a k -complementary edge is used, then the $k - r$ positions in which u and v agree among the first k must be changed individually. \square

Proposition 2 immediately yields $d(Q_{n,k}) = (n - k) + \lceil \frac{k}{2} \rceil$, which equals $n - \lfloor \frac{k}{2} \rfloor$. This was observed by Tzeng and Wei [24], along with an algorithm for finding shortest paths joining vertices. Note that if u and v differ in more than $\lceil \frac{k}{2} \rceil$ positions among the first k , then every shortest u, v -path contains exactly one k -complementary edge.

3 Construction of Paths

In this section we will prove our main results by constructing disjoint paths of bounded length joining any two vertices in $Q_{n,k}$.

Let P be a path $u_0 \rightarrow u_1 \rightarrow \dots \rightarrow u_{\ell-1} \rightarrow u_\ell$ from the vertex u_0 to a vertex u_ℓ in $Q_{n,k}$. The path P traverses the edges $u_0u_1, u_1u_2, \dots, u_{\ell-1}u_\ell$. Since the edge $u_{j-1}u_j$ is in E_{d_j} for some $d_j \in \{0, 1, \dots, n\}$, and every vertex is incident with exactly one edge in E_{d_j} , we can represent the path P from u_0 by the list (d_1, \dots, d_ℓ) , where d_j indicates the type of the edge joining u_{j-1} and u_j . For example, in $Q_{5,3}$ the path originating from 00000 determined by $(2, 0, 5)$ is $00000 \xrightarrow{2} 00010 \xrightarrow{0} 00101 \xrightarrow{5} 10101$. Note that the length of a path determined by a list I is the number of elements in I .

We use the following two lemmas to construct disjoint paths. A *proper segment* of a list is a string of consecutive elements that is not the full list.

Lemma 1. *No two cyclic permutations of a list of distinct elements have a common proper initial segment.*

Proof. Let $I_1 = (d_1, d_2, \dots, d_\ell)$ and $I_j = (d_j, \dots, d_\ell, d_1, \dots, d_{j-1})$ for $2 \leq j \leq \ell$.

Consider j and j' with $1 \leq j' < j \leq \ell$. A proper initial segment of I_j for $j \geq 2$ contains d_j and not d_{j-1} , but this is not true for any initial segment of $I_{j'}$, since when d_j is not the first element, d_{j-1} also occurs in any initial segment containing d_j . Therefore, I_j and $I_{j'}$ have no common proper initial segment. \square

Lemma 2. *Let S be a set of ℓ distinct elements of $\{0, 1, \dots, n\}$. Let I_1, \dots, I_m be orderings of S such that no two have a common proper initial segment. If $\{0, 1, \dots, k\}$ are not all in S , then for any vertex u in $Q_{n,k}$, the lists I_1, \dots, I_m determine disjoint paths to a single vertex.*

Proof. Let $S = \{d_1, \dots, d_\ell\}$ and $[k] = \{1, \dots, k\}$. If $0 \notin S$, then each path reaches the vertex v that differs from u in the positions of S . If $0 \in S$, then each path reaches the vertex v that differs from u in the positions of $([k] - S) \cup (S - \{0\} - [k])$.

If two subsets of S both contain 0 or both omit 0, then they produce paths to the same vertex only if they are the same set. If $0 \in T \subseteq S$ and $0 \notin T' \subseteq S$, then T and T' produce paths to the same vertex only if they agree outside $[k]$ and intersect $[k]$ in complementary subsets. By the hypothesis that $0, 1, \dots, k$ are not all present in S , this cannot occur.

Therefore, the paths from u determined by I_j and $I_{j'}$ have a common internal vertex if and only if they have a common proper initial segment. \square

Lemma 3. *Let $I = (d_1, \dots, d_\ell)$ with all d_i distinct and in $\{0, 1, \dots, n\}$. If $\{0, 1, \dots, k\}$ are not all in I , then for any vertex u in $Q_{n,k}$, the ℓ cyclic permutations of I determine disjoint paths to a single vertex.*

Proof. The conclusion follows immediately from Lemmas 1 and 2. \square

When G is the complete graph K_n with $n \geq 3$, we have $D_\omega(G) = 1$ but $d_\omega(G) = 2$ for $2 \leq \omega \leq n - 1$. Since $Q_{2,2} = K_4$, we consider $Q_{n,k}$ with $n \geq 3$ and $2 \leq k \leq n$.

Theorem 1. *For any two distinct vertices u and v in $Q_{n,k}$ with $n \geq 3$ and $2 \leq k \leq n$, there exist $n + 1$ disjoint u, v -paths of length at most $d(Q_{n,k}) + 1$, such that at least $n - \lfloor \frac{k}{2} \rfloor - 1$ of the paths have length at most $d(Q_{n,k})$.*

Proof. The vertex transitivity of $Q_{n,k}$ and Proposition 1 allow us to assume $u = 0^n$ and $v = 0^{n-k-j}1^j0^{k-i}1^i$, where $0 \leq i \leq k$ and $0 \leq j \leq n - k$. We will construct the desired paths. Let $r = \min\{i + j, k - i + j + 1\}$. Recall that $d(Q_{n,k}) = n - \lfloor \frac{k}{2} \rfloor$. We consider two cases according to the relationship between r and $d(Q_{n,k})$.

Case 1: $r < d(Q_{n,k})$. We first specify a list I of length r , in two cases. Note that $r = i + j$ when $i \leq k - i$ and $r = k - i + j + 1$ when $i > k - i$.

Case	Condition	I
A	$i \leq k - i$	$(1, \dots, i, k + 1, \dots, k + j)$
B	$i > k - i$	$(0, i + 1, \dots, k, k + 1, \dots, k + j)$

Not all of $0, 1, \dots, k$ appear in I , since having 0 requires $i > k/2$ (Case B), and then also having 1 requires $i = 0$, a contradiction. Hence Lemma 3 applies, so the r cyclic permutations of I determine r disjoint u, v -paths of length r .

The remaining $n - r + 1$ paths, with length $r + 2$, are specified by adding one of $\{0, 1, \dots, n\} - I$ at both the beginning and the end of I . Let I' be the list obtained by adding h , and let P be the path from u determined by I' .

If $h > k$ or if $h \in [k]$ and $0 \notin I$, then u and v agree in position h , and P is the only path in the constructed set containing vertices that differ from them in position h . Furthermore, all internal vertices of P differ from u and v in position h .

If $h = 0$, then $i \leq k - i$ (Case A). All internal vertices of P differ from u and v in positions k and $k - 1$, and no other path has any such vertices.

If $h \in [k]$ and $0 \in I$, then $1 \leq h \leq i$ (Case B). The first vertex of P after u differs from u only in position h . The next vertex disagrees with u on all of positions $1, \dots, i$ except h , and this remains true of all other internal vertices of P , because I contains no element of $\{1, \dots, i\}$. All the other paths in the construction have no vertices satisfying either of these conditions.

Since $r < d(Q_{n,k})$, all the paths have length at most $d(Q_{n,k}) + 1$; in fact, all have length at most $d(Q_{n,k})$ unless $r = d(Q_{n,k}) - 1$. In this case there are r paths of length r , which suffices since $r = d(Q_{n,k}) - 1 = n - \lfloor \frac{k}{2} \rfloor - 1$.

Case 2: $r \geq d(Q_{n,k})$. Let $s = d(Q_{n,k})$. Since $r = \min\{i + j, k - i + j + 1\}$, we have $i + j \geq s$ and $k - i + j + 1 \geq s$, so $k + 2j + 1 \geq 2s$. If $j \leq n - k - 1$, then $2n - k - 1 \geq 2s = 2n - 2\lfloor \frac{k}{2} \rfloor$, which is impossible. Hence $j = n - k$. With $j = n - k$, we have $i \geq \lceil \frac{k}{2} \rceil$ and $k + 1 - i \geq \lceil \frac{k}{2} \rceil$, so $\lceil \frac{k}{2} \rceil \leq i \leq \lfloor \frac{k}{2} \rfloor + 1$. When $i = \lceil \frac{k}{2} \rceil$, we have $r = i + j$; when $i = \lfloor \frac{k}{2} \rfloor + 1$, we have $r = k - i + j + 1$. Both cases apply when k is odd. In either case, $r = n - \lfloor \frac{k}{2} \rfloor = d(Q_{n,k})$, and we define three lists.

Case	i	I	J	I'
A	$\lceil \frac{k}{2} \rceil$	$(1, \dots, i, k + 1, \dots, n)$	$(k + 1, \dots, n)$	$(0, i + 1, \dots, k)$
B	$\lfloor \frac{k}{2} \rfloor + 1$	$(0, i + 1, \dots, k, k + 1, \dots, n)$	$(k + 1, \dots, n)$	$(1, \dots, i)$

Since $j = n - k$, in each case I has length r . By Lemma 3, the cyclic permutations of I yield r disjoint u, v -paths of length r . Since $r = n - \lfloor \frac{k}{2} \rfloor = d(Q_{n,k})$, this yields enough paths of length at most $d(Q_{n,k})$. Let $T = \{i + 1, \dots, k\}$ in Case A, $T = \{1, \dots, i\}$ in Case B. Every vertex in each of these paths is constant in the positions of T (all-0 or all-1), and in fact all-0 in Case A.

Since $r = n - \lfloor \frac{k}{2} \rfloor$, we only need to find $\lfloor \frac{k}{2} \rfloor + 1$ more paths of length at most $d(Q_{n,k}) + 1$. Note that I' has length $\lfloor \frac{k}{2} \rfloor + 1$. Form $\lfloor \frac{k}{2} \rfloor + 1$ lists by inserting J after the first element of each cyclic permutation of I' . The first and last lists are $(0, J, i + 1, \dots, k)$ and $(k, J, 0, i + 1, \dots, k - 1)$ in Case A, $(1, J, 2, \dots, i)$ and $(i, J, 1, \dots, i - 1)$ in Case B. Each of these lists has length $n - \lceil \frac{k}{2} \rceil + 1$, which is at most $d(Q_{n,k}) + 1$.

Each of these lists is an ordering of a single set of elements. By an argument like that of Lemma 1, they have no common proper initial segments. Since they also do not contain all of $\{0, 1, \dots, k\}$, by Lemma 2 these paths are disjoint.

In Case A, each internal vertex on each of these paths is not all 0 in the positions of T . In Case B, each internal vertex on each of these paths has between 1 and $|T| - 1$ nonzero positions in T . Hence these paths are disjoint from the earlier paths. \square

4 Consequences

From Theorem 1 and the definition of wide diameter, we immediately obtain an upper bound on $d_\omega(Q_{n,k})$.

Corollary 1. *If $n \geq 3$ and $2 \leq k \leq n$, then*

$$d_\omega(Q_{n,k}) \leq \begin{cases} d(Q_{n,k}) & \text{for } 1 \leq \omega < n - \lfloor \frac{k}{2} \rfloor, \\ d(Q_{n,k}) + 1 & \text{for } n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1. \end{cases}$$

Proof. When $\omega < n - \lfloor \frac{k}{2} \rfloor$, Theorem 1 provides at least ω disjoint paths with length at most $d(Q_{n,k})$ joining any two vertices in $Q_{n,k}$. When $\omega \leq n + 1$, it provides at least ω such paths with length at most $d(Q_{n,k}) + 1$. \square

We next give a lower bound on the fault diameter $D_\omega(Q_{n,k})$.

Lemma 4. *Fix $n \geq 3$. If $2 \leq k \leq n$ and $n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1$, then*

$$D_\omega(Q_{n,k}) \geq d(Q_{n,k}) + 1.$$

Proof. Since $D_\omega(G)$ is nondecreasing in ω , proving $D_{n - \lfloor \frac{k}{2} \rfloor}(Q_{n,k}) \geq d(Q_{n,k}) + 1$ is sufficient. Let $u = 0^n$ and $v = 1^{n-k}0^{k-i}1^i$, where $i = \lceil \frac{k}{2} \rceil - 1$. Note that v has 1s in $n - \lfloor \frac{k}{2} \rfloor - 1$ positions. Let W be the set of neighbors of u whose single 1 occurs in a position where v has a 1, so $|W| = n - \lfloor \frac{k}{2} \rfloor - 1$. On any u, v -path in $Q_{n,k} - W$, the neighbor of u has a single 1 in a position among $i + 1, \dots, k$ or is $0^{n-k}1^k$.

By Proposition 2, the distance between v and a neighbor u' of u not in W is $n - k + \min\{i + 1, k - i\}$. Since $i = \lceil \frac{k}{2} \rceil - 1$, the distance is $n - \lfloor \frac{k}{2} \rfloor$, which equals $d(Q_{n,k})$. Hence every u, v -path in $Q_{n,k} - W$ has length at least $d(Q_{n,k}) + 1$. \square

Theorem 2. *For $3 \leq n$ and $2 \leq k \leq n$,*

$$D_\omega(Q_{n,k}) = d_\omega(Q_{n,k}) = \begin{cases} d(Q_{n,k}) & \text{for } 1 \leq \omega < n - \lfloor \frac{k}{2} \rfloor; \\ d(Q_{n,k}) + 1 & \text{for } n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1. \end{cases}$$

Proof. Since $d(Q_{n,k}) \leq D_\omega(Q_{n,k}) \leq d_\omega(Q_{n,k})$ for $1 \leq \omega \leq n + 1$, Corollary 1 yields $D_\omega(Q_{n,k}) = d_\omega(Q_{n,k}) = d(Q_{n,k})$ for $1 \leq \omega < n - \lfloor \frac{k}{2} \rfloor$.

For $n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1$, Corollary 1 and Lemma 4 yield $D_\omega(Q_{n,k}) = d_\omega(Q_{n,k}) = d(Q_{n,k}) + 1$. \square

Theorem 2 shows that the fault diameter $D_\omega(Q_{n,k})$ equals the wide diameter $d_\omega(Q_{n,k})$ for the enhanced hypercubes $Q_{n,k}$. More importantly, they equal the traditional diameter when $1 \leq \omega < n - \lfloor \frac{k}{2} \rfloor$, and they exceed it only by 1 when $n - \lfloor \frac{k}{2} \rfloor \leq \omega \leq n + 1$. Thus, the resilience of enhanced hypercubes is similar to that of hypercubes, which increases the appeal of enhanced hypercubes.

Acknowledgements

This work was supported partially by NSFC (Nos. 11101378, 11571319, 11571044), and ZJNSF (No. LY14A010009). Research of D.B. West was supported by 1000 Talent Plan, State Administration of Foreign Experts Affairs, China.

References

- [1] A. El-Amawy and S. Latifi, Properties and performance of folded hypercubes, *IEEE Trans. Parallel Distrib. Syst.* 2 (1991) 31–42.
- [2] I. Banič, R. Erveš, J. Žerovnik, The edge fault-diameter of Cartesian graph bundles, *European J. Combin.* 30 (2009) 1054–1061.
- [3] I. Banič, J. Žerovnik, The fault-diameter of Cartesian products, *Adv. Appl. Math.* 40 (2008) 98–106.
- [4] I. Banič, J. Žerovnik, Wide diameter of Cartesian graph bundles, *Discrete Math.* 310 (2010) 1697–1701.
- [5] C.-H. Chang, C.-K. Lin, J.J.M. Tan, H.-M. Huang, L.-H. Hsu, The super spanning connectivity and super spanning laceability of the enhanced hypercubes, *J. Supercomput.* 48 (2009) 66–87.
- [6] C.-P. Chang, C.-C. Wu, Conditional fault diameter of crossed cubes, *J. Parallel Distr. Comput.* 69 (2009) 91–99.
- [7] X.-B. Chen, Edge-fault-tolerant diameter and bipanconnectivity of hypercubes, *Inform. Proc. Lett.* 110 (2010) 1088–1092.
- [8] K. Day, A. E. Al-Ayyoub, Fault diameter of k -ary n -cube networks, *IEEE Trans. Parallel Distrib. Syst.* 8 (1997) 903–907.
- [9] D.Z. Du, D.F. Hsu, Y.D. Lyuu, On the diameter vulnerability of Kautz digraphs, *Discrete Math.* 151 (1996) 81–85.
- [10] T.-L. Kung, C.-K. Lin, T. Liang, L.-Y. Hsu, J.J.M. Tan, Fault diameter of hypercubes with hybrid node and link faults, *J. Inter. Networks* 10 (3) (2009) 233–242.
- [11] S. Latifi, Combinatorial analysis of the fault-diameter of the n -cube, *IEEE Trans. Comput.* 42 (1993) 27–33.
- [12] F.T. Leighton, *Introduction to Parallel Algorithms and Architectures: Arrays, Trees, and Hypercubes*, Morgan Kaufmann, 1992.
- [13] S.C. Liaw, G. J. Chang, Generalized diameters and Rabin numbers of networks, *J. Comb. Optim.* 4 (1999) 371–384.
- [14] S.C. Liaw, G.J. Chang, Wide diameters of butterfly networks, *Taiwan. J. Math.* 3 (1) (1999) 83–88.
- [15] S.C. Liaw, G.J. Chang, F. Cao, D.F. Hsu, Fault-tolerant routing in circulant networks and cycle prefix networks, *Ann. Combin.* 2 (1998) 165–172.
- [16] H. Liu, Properties of enhanced hypercube networks, *J. Syst. Sci. Inform.* 6 (2008) 251–216.

- [17] M. Ma, J.-M. Xu, Algebraic properties and panconnectivity of folded hypercubes, *Ars Combin.* 95 (2010) 179–186.
- [18] K. Menger, Zur allgemeinen Kurventheorie, *Fund. Math.* 10 (1927) 95–115.
- [19] Y. Saad and M.H. Schultz, Topological properties of hypercubes, *IEEE Trans. Comput.* 37 (1988) 867–872.
- [20] T. Shi, M. Lu, Fault-tolerant diameter for three family interconnection networks, *J. Combin. Optim.* 23 (2012) 471–482.
- [21] E. Sinó, J.L.A. Yebra, The vulnerability of the diameter of folded n -cubes, *Discrete Math.* 174 (1997) 317–322.
- [22] X. Tan, J. H. Park, Some topological properties of folded Petersen graph, *Discrete Math.* 308 (2008) 4786–4790.
- [23] T.-H. Tsai, Y.-C. Chen, J.J.M. Tan, Topological properties on the wide and fault diameters of exchanged hypercubes, *IEEE Trans. Parallel Distrib. Syst.* 25 (2014) 3317–3327.
- [24] N.-F. Tzeng, S. Wei, Enhanced hypercubes, *IEEE Trans. Comput.* 40 (1991) 284–294.
- [25] D. Wang, Diagnosability of enhanced hypercubes, *IEEE Trans. Comput.* 43 (1994) 1054–1061.
- [26] D. Wang, Diagnosability of hypercubes and enhanced hypercubes under the comparison diagnosis model, *IEEE Trans. Comput.* 48 (1999) 1369–1374.
- [27] J.-M. Xu, *Topological Structure and Analysis of Interconnection Networks*, Kluwer Academic Publishers, Dordrecht/Boston/London, 2001.
- [28] J.-M. Xu, Wide-diameter of Cartesian product graphs and digraphs, *J. Combin. Optim.* 8 (2) (2004), 171–181.
- [29] M. Xu, J.-M. Xu, X.-M. Hou, Fault diameter of Cartesian product graphs, *Inform. Proc. Lett.* 93 (2005) 245–248.
- [30] J.-S. Yang, J.-M. Chang, K.-J. Pai, H.-C. Chan, Parallel construction of independent spanning trees on enhanced hypercubes, *IEEE Trans. Parallel Distrib. Syst.* 26 (2015) 3090–3098.