

Covering a Triangle with Positive and Negative Homothetic Copies

Zoltán Füredi

Department of Mathematics, University of Illinois at
Urbana-Champaign,
Urbana, IL 61801, USA
z-furedi@math.uiuc.edu

and

Rényi Institute of Mathematics, Hungarian Academy of Sciences,
P.O. Box 127, Budapest, H-1364 Hungary
furedi@renyi.hu

Abstract Let Δ_0 be a triangle and let $\mathcal{H} = \{\Delta_1, \dots, \Delta_n\}$ be a set of homothetic copies of Δ_0 , $\Delta_i = x_i \Delta_0$, $x_1 \geq x_2 \geq \dots \geq x_n \geq 0$. We prove that $\sum x_i^2 \geq 1 + x_2$ implies that there are positive and negative signs $\varepsilon_1, \dots, \varepsilon_n = \pm 1$ and there exist translates of $\varepsilon_1 \Delta_1, \dots, \varepsilon_n \Delta_n$ that cover Δ_0 .

This result is used to answer a problem of Xu, Yuan and Ding [16]. We show that if Δ_0 is the isosceles right triangle and if the total area of \mathcal{H} is at least $(1 + \sqrt{2})/2 = 1.207\dots$ times the area of Δ_0 , then there exist rotations ρ_i , each of them multiples of 45° , and translates of $\rho_1 \Delta_1, \dots, \rho_n \Delta_n$ that cover Δ_0 .

1. Translation Coverings

Let C be a *disk*, i.e., convex, compact set on the Euclidean plane with interior points. Let $\mathcal{H} = \{C_1, \dots, C_i, \dots\}$ be a finite sequence of disks. We say that \mathcal{H} permits

Discrete Comput. Geometry 2007

masfe18.tex October 25, 2006

This research was supported in part by the Hungarian National Science Foundation under Grants OTKA 062321 and 060427 and by the National Science Foundation under Grant NFS DMS 06-00303.

a **translation covering** of C if there exist translations τ_i such that $C \subseteq \cup_i \tau_i(C_i)$. Moser and Moon [12] showed that if Q is the unit square and \mathcal{H} is a set of squares of sizes x_1, x_2, \dots with total area $\sum_i x_i^2 \geq 3$ and with sides parallel (or orthogonal) to Q , then \mathcal{H} permits a translation covering of Q . This is the best possible bound as one can see from the example $x_1 = x_2 = x_3 = 1 - \varepsilon$, $x_4 = \dots = 0$.

L. Fejes Tóth proposed the following more general question. Suppose that each C_i is a (positive) homothetic copy of C . How large must the sum of areas of the C_i 's be, so that C can be covered by translates of the C_i 's? Denote the ratio of this minimum (infimum) and the area of C by $f(C)$. The above cited theorem of Moon and Moser states $f(Q) = 3$. This easily implies $f(C) \leq 12$ for every disk (Bezdek and Bezdek [3]) and it was recently improved to $f(C) \leq 6.5$ by Januszewski [10]. One can observe that for any C one has $f(C) \geq 2$ (two copies of size $1 - \varepsilon$ can't cover a diameter of C). Bezdek and Bezdek [3] conjectured that this is achievable for any triangle Δ . This has been recently established by the present author [7] as

$$f(\Delta) = 2. \quad (1)$$

Small Sets. In fact, in the case of the unit square, Q , Moon and Moser showed that for $x_1 \geq \dots \geq x_n$ the total area

$$\sum x_i^2 \geq 1 + 2x_1 \quad (2)$$

ensures a translation covering. Analyzing their proof one can see that $1 + x_1 + x_2$ is sufficient. This was generalized in [7] as follows.

Denote a convex body by $C \subset \mathbf{R}^2$, the infimum of the densities of coverings \mathbf{R}^2 by translates of C by $\vartheta_T(C)$, and the family of all copies of C whose positive homothety ratios are at most δ by $\mathcal{F}_\delta(C)$. For every $\varepsilon > 0$ there exists a positive $\delta = \delta(\varepsilon, C)$ such that if the total area of sets from any family $\mathcal{H}(C) \subset \mathcal{F}_\delta(C)$ is at least

$$(\vartheta_T(C) + \varepsilon)\text{Area}(C), \quad (3)$$

then there are translates of the sets from $\mathcal{H}(C)$ which cover C .

Large Sets. It is known that $\vartheta_T(C) \leq 3/2$ for every planar disk (this upper bound is due to Besicovitch [2] although usually attributed to Fáy [5]) and this density can be obtained by a hexagonal lattice arrangement. See, e.g., the excellent monograph by Pach and Agarwal [13]. It is also conjectured that $\vartheta_T(\Delta) = 3/2$ (see, e.g., L. Fejes-Tóth [6]). Obviously, $\vartheta_T(Q) = 1$. The theorem mentioned in (3) says that if the copies of C in \mathcal{H} are small, then good coverings can be constructed, so, ironically, $f(\Delta)$ and $f(Q)$ are large (2 and 3, resp.) because \mathcal{H} can contain large copies. Apparently, large copies mean large waste. This happens also in the coverings discussed below.

2. Translations and Rotations

Suppose that \mathcal{H} consists of positive **and negative** homothetic copies of a triangle Δ_0 , then a total area at least $4\text{Area}(\Delta_0)$ ensures a translation covering, and here the constant 4 is the best possible. This was conjectured by K. Böröczky and proved by Januszewski [9]. The more special case of \mathcal{H} being a finite sequence purely of homothetic copies of $-\Delta_0$ was proved earlier by Vászárhelyi [14]. She also considered translation coverings of the triangle Δ_0 when \mathcal{H} consists of homothetic copies each of them **rotated** by a certain angle φ , see [15]. Here we consider a more general problem.

Definition 1 *Let C be a disk, let $\mathcal{H} = \{C_1, \dots, C_i, \dots\}$ be a finite sequence of disks, and let Γ be a set of planar motions (usually a group of rotations). Let τ_i be a translation and $\gamma_i \in \Gamma$, then the system $\{\tau_1(\gamma_1 C_1), \tau_2(\gamma_2 C_2), \dots\}$ is called a Γ -translation of \mathcal{H} . We say that \mathcal{H} permits a Γ -covering of C if there exist translations τ_i and motions $\gamma_i \in \Gamma$ such that $C \subseteq \cup_i \tau_i(\gamma_i C_i)$.*

Suppose that each C_i is a (positive) homothetic copy of C . How large must the sum of areas of the C_i 's be, so that C can be covered by Γ -translates of the C_i 's? Denote the ratio of this minimum (infimum) and the area of C by $f_\Gamma(C)$. Denote the infimum of the densities of coverings of the plane by Γ -translated *congruent* copies of C by $\vartheta_\Gamma(C)$.

Theorem 1 *For every $\varepsilon > 0$ there exists a $\delta = \delta(\varepsilon, C, \Gamma) > 0$ such that the following holds. If \mathcal{H} is a set of (positive) homothetic copies of C and each member of \mathcal{H} is smaller than δC and for the total area we have*

$$\sum_{H \in \mathcal{H}} \text{Area}(H) \geq (\vartheta_\Gamma(C) + \varepsilon) \text{Area}(C),$$

then there exist Γ -translates of the members of \mathcal{H} that cover C .

The proof is a bit technical but straightforward, one can follow the argument in [7], we omit it. Naturally, a similar statement holds for every (finite) dimension.

ε and ρ -Coverings. The cases $\Gamma = \{\text{rotations of integer multiples of } \pi\}$ and $\Gamma = \{\text{rotations of integer multiples of } \pi/4\}$ are called ε - and ρ -coverings, respectively.

Theorem 2 *Let Δ_0 be a triangle and let $\mathcal{H} = \{\Delta_1, \dots, \Delta_n\}$ be a set of homothetic copies of Δ_0 , $\Delta_i = x_i \Delta_0$, $x_1 \geq x_2 \geq \dots \geq x_n > 0$. Suppose that*

$$\sum x_i^2 \geq 1 + x_2. \quad (4)$$

Then there are positive and negative signs $\varepsilon_1, \dots, \varepsilon_n \in \{1, -1\}$ and there exist translates of $\varepsilon_1 \Delta_1, \dots, \varepsilon_n \Delta_n$ that cover Δ_0 .

Unlike in the Moon–Moser theorem (2) here the second-order term depends on x_2 . This theorem implies $f_\varepsilon(\Delta_0) = 2$, but we have already known this from (1), since $f_\varepsilon \leq f$. Using Theorem 2 we answer a problem of Xu, Yuan and Ding [16]. They investigated $f_\rho(R_0)$ where R_0 is a right isosceles triangle. Let $\mathcal{H} = \{R_1, \dots, R_n\}$, $R_i = x_i R_0$ and $x_1 \geq x_2 \geq \dots \geq x_n > 0$. They showed that if

$$\sum_i x_i^2 \geq \frac{8 + \sqrt{8}}{7} + \sqrt{2}x_1 \sim 1.56 + 1.41x_1$$

then \mathcal{H} permits a ρ -cover. Theorem 2 clearly supersedes this. Here we also determine $f_\rho(R_0)$, it is $c := (1 + \sqrt{2})/2 = 1.2071\dots$

Theorem 3 *Let R_0 be the triangle with vertices $(0, 0)$, $(a, 0)$, and $(0, a)$ and let $\mathcal{H} = \{R_1, \dots, R_n\}$ be a set of homothetic copies of R_0 . Suppose that*

$$\sum_{i \geq 1} \text{Area}(R_i) \geq \frac{1 + \sqrt{2}}{2} \text{Area}(R_0). \quad (5)$$

Then there are triangles R'_i congruent to R_i and with sides parallel or orthogonal to some sides of R_0 such that $R_0 \subseteq \cup_{i \geq 1} R'_i$.

On the other hand, to establish the lower bound for $f_\rho(R)$ one needs, for every $\eta > 0$, a sequence x_1, \dots, x_n with $\sum x_i^2 > (c - \eta)a^2$ such that $\{x_1 R_1, \dots, x_n R_n\}$ cannot ρ -cover R_0 . It is not difficult to see that one can take

$$x_{k+1} := (1 - \eta/2)(\sqrt{2} - 1)^k, \quad \text{where } k = 0, 1, 2, \dots$$

In Sections 3–5 we show how to ε -cover Δ_0 , i.e., Theorem 2. In Sections 6–8 we prove Theorem 3 showing $f_\rho(R_0) \leq c$. The uninteresting proof of $f_\rho(R_0) \geq c$ is omitted.

3. Covering a Strip of Δ_0

The aim of this and the following two sections is to prove Theorem 2. We will define a translation and a rotation (of 0° or 180°) of each member of \mathcal{H} . The procedure also supplies an algorithm with linear running time after ordering the triangles by their sizes. As ε -translation coverings are affine invariant, we may suppose that Δ_0 (and all members of \mathcal{H}) are isosceles, right triangles. Suppose that the vertices of Δ_0 are $(0, 0)$, $(a, 0)$, and $(0, a)$.

First, we prove the following lemma (in the second term there is an x_1 not x_2 as in Theorem 2). Let $\mathcal{H} = \{\Delta_1, \dots, \Delta_n\}$ be a set of homothetic copies of Δ_0 with side lengths

$$a \geq x_1 \geq x_2 \geq \dots \geq x_n \geq 0. \quad (6)$$

Suppose that

$$x_1 + x_2 + \dots + x_n \geq 2a. \quad (7)$$

Lemma 4 *There exists a trapezoid T of height h on the bottom of Δ_0 , its vertices are $(0, 0)$, $(a, 0)$, $(0, h)$ and $(a - h, h)$, and there exists an $s \geq 1$ such that the following two properties hold:*

- *An ε -translation of the largest $2s + 1$ triangles $\{\Delta_1, \dots, \Delta_{2s+1}\}$ covers T .*
- *In case of $h < a$ the covering of T is economical, that is*

$$2\text{Area}(T) \geq \sum_{1 \leq i \leq 2s+1} 2\text{Area}(\Delta_i) - ax_1 + (a - h)x_{2s+1}. \quad (8)$$

Proof: If $x_1 = a$ or $x_1 = 0$, then there is nothing to prove so we may suppose

$$a > x_1 > 0.$$

Define s as the smallest integer with

$$a \leq x_1 + \sum_{1 \leq i \leq s} (x_{2i} + x_{2i+1} - x_{2s+1}). \quad (9)$$

We claim that such an s exists and $1 < 2s + 1 \leq n$. Obviously, s must be at least 1. On the other hand, the opposite of (9) and monotonicity (6) imply that

$$a - x_1 > x_2 + x_4 + \dots + x_{2s} \geq x_3 + x_5 + \dots + x_{2s+1},$$

so $2a - x_1 > \sum_{i < 2s+1} x_i$. Thus $2a > \sum_{i < 2s+2} x_i$ and (7) implies $2s + 3 \leq n$. So two more x_i 's can be joined to $\{x_1, \dots, x_{2s+1}\}$ and in finitely many steps we reach an s satisfying (9).

Let \mathcal{R} denote the set of the largest $2s + 1$ triangles from \mathcal{H} . Geometrically, (9) means that one can translate $\Delta_1, -\Delta_2, \dots, (-1)^{i+1}\Delta_i, \dots, \Delta_{2s+1}$ such that they cover a trapezoid of height x_{2s+1} and base exceeding a , see Figure 1.




Figure 1: $\Delta_1, -\Delta_2, \dots, -\Delta_{2s}, \Delta_{2s+1}$ cover a trapezoid of height x_{2s+1} .

By definition

$$a \geq x_1 + \sum_{1 \leq i \leq s-1} (x_{2i} + x_{2i+1} - x_{2s-1}). \quad (10)$$

We will use the following consequence of (6) and (10)

$$x_1 + x_2 + \dots + x_s \leq a. \quad (11)$$

Now we are ready to define the trapezoid T and its ε -translation covering by \mathcal{R} . The only thing needed to define T is its height h . It is obtained from the following equation

$$x_1 + \sum_{1 \leq i \leq s} (x_{2i} + x_{2i+1} - h) = a. \quad (12)$$

Comparing (9) and (11) we obtain that

$$x_{2s+1} \leq h. \quad (13)$$

Inequalities (6) and (11) imply $\sum_{1 \leq i \leq 2s+1} x_i \leq (2s + 1)a/s$. This and (12) give

$$h \leq \frac{s+1}{s^2}a. \quad (14)$$

We claim that \mathcal{R} can ε -cover T . Place the apex vertex of Δ_1 (the vertex with the right angle) to the point $(a - x_1, 0)$, its other two vertices are $(a - x_1, x_1)$ and $(a, 0)$.

For $1 \leq i \leq s$ the triangles $-\Delta_{2i}$ and Δ_{2i+1} can form a rectangle of height h and base length $(x_{2i} + x_{2i+1} - h)$. In the case $i = s$ this base might be negative, but as the sum of the base lengths is $a - x_1$ by (12) the s rectangles can be put next to each other to cover a rectangle of size $(a - x_1) \times h$, see Figure 2.




Figure 2: The trapezoid T of height h is covered by $\Delta_1, \dots, \Delta_{2s+1}$.

Informally Figure 2 is obtained from Figure 1 by pushing the triangles horizontally closer to each other so they can cover a strip wider than x_{2s+1} . More precisely, for $0 \leq i \leq s$ place Δ_{2i+1} such that its apex vertex lies on the line $y = 0$, it is at the point

$$\left(\sum_{i < j \leq s} (x_{2j} + x_{2j+1} - h), 0 \right),$$

and the triangles $-\Delta_{2i}$ are put below the line $y = h$, their apex vertices are at

$$\left(\sum_{i \leq j \leq s} (x_{2j} + x_{2j+1} - h), h \right).$$

We obtained that $\bigcup_{\Delta_i \in \mathcal{R}} \varepsilon_i \Delta \supseteq T$, where $\varepsilon_i = (-1)^{i+1}$.

In the case $h \geq a$ (by (14) this could happen only if $s = 1$) \mathcal{R} ε -covers the whole Δ_0 , so our procedure stops. To finish the proof of the Lemma in case of

$$h \leq a \tag{15}$$

we have to verify inequality (8). The proof of this inequality is purely algebraic, it does not use any geometry, therefore it is postponed to the next section as Lemma 5.

4. Proof of an Inequality

The following lemma implies (8).

Lemma 5 *Suppose that $a \geq x_1 \geq x_2 \geq \cdots \geq x_n \geq 0$ and $a \geq h \geq x_{2s+1}$ are real numbers and $s \geq 1$ is an integer satisfying (10) and (12), i.e.,*

$$\begin{aligned} x_1 + \sum_{1 \leq i \leq s-1} (x_{2i} + x_{2i+1} - x_{2s-1}) &\leq a, \\ x_1 + \sum_{1 \leq i \leq s} (x_{2i} + x_{2i+1} - h) &= a. \end{aligned}$$

Then

$$-\left(\sum_{i=1}^{2s+1} x_i^2\right) + ax_1 + 2ah - h^2 - ax_{2s+1} + hx_{2s+1} \geq 0.$$

Proof. It is high school algebra, but because the domain of F has many faces we have to distinguish many cases. First, as we have seen above, elementary calculation shows that the inequalities (6), (10), (12), (13), and (15) (appearing in the above lemma) imply that (9), (11), and (14) must hold, too.

Apply (6), (15), and (13), and then (12). We obtain

$$\begin{aligned} F(a, x_1, \dots, x_{2s+1}, h) &:= -\left(\sum x_i^2\right) + ax_1 + ah + (a-h)(h-x_{2s+1}) \\ &\geq -x_1\left(\sum x_i\right) + ax_1 + ah \\ &= -x_1(a+sh) + ax_1 + ah = h(a-sx_1). \end{aligned} \tag{16}$$

This implies that $F \geq 0$ for $sx_1 \leq a$, which is always true if $s = 1$.

Suppose that F can take negative values. Since it is homogeneous of degree 2, it takes a negative value with all variables at most 1. Since F is continuous on a compact part of the $(2s+3)$ -dimensional hypercube $[0, 1]^{2s+3}$, it takes its minimum, say at the point $(a, x_1, \dots, x_{2s+1}, h)$. We claim that for these values

$$a > x_1 > 0, \quad a > h > 0, \quad sx_1 > a, \quad s \geq 2. \tag{17}$$

We only have to show that these inequalities are strict. Indeed, $F < 0$ and (16) imply $h > 0$ and $sx_1 > a$, then (6) gives $s \geq 2$. In the case of $a = x_1$ (11) gives $x_2 = \cdots = x_{2s+1} = 0$, so $F = 2ah - h^2 \geq 0$. For $x_1 = 0$, inequality (6) again gives $x_2 = \cdots = x_{2s+1} = 0$ and $F = 2ah - h^2$. Finally, $a = h$ contradicts (14).

Define k as the largest integer with $x_1 = x_2 = \cdots = x_k$. Then (11) and $sx_1 > a$ from (17) give $k < s$. We claim that

$$a - kx_1 + kh > 0. \quad (18)$$

Using the monotonicity and that $k < s$, then (11), and finally the positivity of h , one gets the equivalent form

$$kx_1 = x_1 + \cdots + x_k \leq x_1 + \cdots + x_s \leq a < a + kh.$$

Now let $\eta > 0$ be sufficiently small and define

$$\begin{aligned} a' &= a - k\eta \\ x'_i &= \begin{cases} x_i - \eta & \text{for } 1 \leq i \leq k \\ x_i & \text{for } k < i \leq 2s + 1 \end{cases} \\ h' &= h \quad \text{unchanged.} \end{aligned}$$

One can see that $k < 2s - 1$ and (17) imply that the inequalities in Lemma 5 (i.e., (6), (10), (12), (13) and (15)) still hold for $(a', x'_1, \dots, x'_{2s+1}, h')$, and it is a feasible solution. However,

$$\begin{aligned} F(a, x_1, \dots, x_{2s+1}, h) - F(a', x'_1, \dots, x'_{2s+1}, h') \\ = \eta k(h - x_{2s+1}) + \eta(a - kx_1 + kh - 2k\eta) > 0. \end{aligned}$$

Here the first term is nonnegative by (13), and the second term is strictly positive by (18) if η is sufficiently small and positive. This final contradiction completes the proof of Lemma 5 (and thus the proof of Lemma 4, too). \square

5. Area Estimates

Here we finish the proof of Theorem 2. We may suppose that the vertices of Δ_0 are $(0, 0)$, $(0, 1)$, and $(1, 0)$ and \mathcal{H} consists of isosceles right triangles of sizes $x_1 \geq \cdots \geq x_n > 0$. Suppose that the integer N is large enough, namely

$$x_n > x_1/N,$$

and replace Δ_1 by N^2 small triangles of sizes x_1/N . The system of $n - 1 + N^2$ triangles obtained is denoted by \mathcal{H}^N . First, we define an ε -covering of Δ_0 by \mathcal{H}^N , and then we return to the case of \mathcal{H} itself.

We form disjoint subgroups $\mathcal{H}_1, \mathcal{H}_2, \dots, \mathcal{H}_m$ of \mathcal{H}^N , and at the same time we define right angled trapezoids T_1, T_2, \dots, T_m such that some ε -translations of the members of \mathcal{H}_k cover T_k for $k = 1, 2, \dots, m$. (T_m is a degenerate trapezoid, i.e., a triangle.) These trapezoids are obtained by cutting Δ_0 by horizontal lines, they are packed on each other, and T_1 lies on the bottom (i.e., adjacent to the x -axis). The base lengths of T_k are denoted by a_k and a_{k+1} , $a_1 = 1 > a_2 > \dots > a_m > a_{m+1} = 0$, its height is h_k , $h_k = a_k - a_{k+1}$ and side lengths are h_k and $\sqrt{2}h_k$, with vertices $(0, 1 - a_k)$, $(0, 1 - a_{k+1})$, $(a_k, 1 - a_k)$, and $(a_{k+1}, 1 - a_{k+1})$. Finally, the groups \mathcal{H}_k consist of smaller and smaller triangles, i.e.,

$$\min_{\Delta \in \mathcal{H}_k} x(\Delta) \geq \max_{\Delta \in \mathcal{H}_{k+1}} x(\Delta), \quad (19)$$

where $x(\Delta)$ stands for the side length of Δ . However, the heights of trapezoids do not necessarily form a monotone sequence. We also maintain

$$\sum_{\Delta \in \mathcal{H}^N \setminus (\mathcal{H}_1 \cup \dots \cup \mathcal{H}_{k-1})} 2\text{Area}(\Delta) \geq a_k^2 + a_k \max_{\Delta \in \mathcal{H}^N \setminus (\mathcal{H}_1 \cup \dots \cup \mathcal{H}_{k-1})} x(\Delta) \quad (20)$$

This condition is very similar to (4); x_1 stands for x_2 .

Our starting case is $k = 1$. Then $\max x(\Delta_i) = x_2$, and $\sum x_i^2 = \sum_{\Delta \in \mathcal{H}} 2\text{Area}(\Delta) \geq a_1^2 + a_1 x_2$. This last inequality implies that $\sum x_i \geq (a_1^2 + a_1 x_2)/x_2 \geq 2a_1$ so conditions (6) and (7) hold, and Lemma 4 can be applied to \mathcal{H}^N and Δ_0 . Then either the lemma supplies an ε -cover of Δ_0 (in the case of $h \geq a_1$), or one obtains a set of triangles $\mathcal{H}_1 \subset \mathcal{H}^N$ and a trapezoid T_1 with horizontal sides $a_1 (= 1)$ and a_2 , and height $h_1 := h$ such that \mathcal{H}_1 ε -cover T_1 and the induction hypotheses (19) and (20)

$$\sum_{i > 2s+1} 2\text{Area}(\Delta_i) \geq (a_1 - h)^2 + (a_1 - h)x_{2s+2}$$

are preserved for the rest of Δ_0 with the rest of \mathcal{H}^N . In general, if $\mathcal{H}_1, \dots, \mathcal{H}_{k-1}$ are already defined, then Lemma 4 either finishes the procedure by completing an ε -cover of Δ_0 , and then $m = k$, or supplies \mathcal{H}_k and T_k satisfying the induction hypotheses for $\mathcal{H}^N \setminus (\mathcal{H}_1 \cup \dots \cup \mathcal{H}_k)$ and $\Delta_0 \setminus (T_1 \cup \dots \cup T_k)$. Since \mathcal{H}^N has finitely many members, our induction procedure results in a full ε -translation covering of Δ_0 .

Since the smallest N^2 triangles are used at last, (if they are used at all, in the above ε -cover) they are contained in the subfamilies $\mathcal{H}_\ell, \mathcal{H}_{\ell+1}, \dots, \mathcal{H}_m$. Among these, $\mathcal{H}_{\ell+1}, \dots, \mathcal{H}_m$ consist purely of triangles of sizes x_1/N . Hence $T_{\ell+1} \cup \dots \cup T_m$ is contained in a triangle Δ'_1 of size x_1 , defined by the vertices $(0, 1)$, $(0, 1 - x_1)$, and

$(x_1, 1 - x_1)$. It is easy to see, that the x_1/N -sized members of \mathcal{H}_ℓ , apart from one or two exceptions, are all in the $2x_1/N$ neighborhood of Δ'_1 .

Consider the coverings by \mathcal{H}^N , \mathcal{H}^{N+1} , etc. It is not difficult to see that the positions of $\Delta_2, \dots, \Delta_n$ converge to a limit. The above considerations show that in that limit position they ε -cover $\Delta_0 \setminus \Delta'_1$. Place Δ_1 onto Δ'_1 to obtain an ε -translation covering of Δ_0 by \mathcal{H} . \square

6. ρ -Covering of the Isosceles Right Triangle, Starting the Induction

We prove Theorem 3 by induction on n . Suppose that $R_i = x_i R_0$, $x_1 \geq \dots \geq x_n > 0$. In the case of $n = 1$, (5) implies that $x_1 \geq \sqrt{c} > 1$ so it certainly can ρ -cover R_0 . If $x_2 \leq (c - 1) = 0.2071\dots$ then Theorem 2 gives an ε -cover and we are done. To avoid using many subscripts, we sometimes use x , y and z for x_1 , x_2 and x_3 , respectively. From now on, we suppose that $y := x_2 \geq (c - 1) = 0.2071\dots$, and we may also suppose that the size of R_0 is 1, that is $a = 1$.

In this section we deal with two cases, using induction if $x > 0.547$ (Step 1), and if $0.6 > x > 0.378$ (Step 2). In the next Section we eliminate the cases $0.38 \geq x > 0.336$ (Step 3) and $x > 0.234$ (Step 5). Here we used computers. Finally, in Section 8, with a different method we finish the induction by investigating the case $x_1 \leq 1/4$.

Step 1. $x \geq 0.547$

Put R_1 into R_0 such that a leg of R_1 lies on the hypotenuse of R_0 and splits it into two segments of lengths x and $\sqrt{2} - x$ (see Figure 3). The set $R_0 \setminus R_1$ can be covered by an isosceles right triangle R' of size $\sqrt{2} - x$. Hence one can use the induction hypothesis for R' and for the remaining $(n - 1)$ triangles $\{R_2, \dots, R_n\}$ if

$$\sum_{i \geq 2} x_i^2 \geq c \left(\sqrt{2} - x \right)^2. \quad (21)$$

Here the left hand side is at least $c - x^2$, so (21) holds if $c - x^2 \geq c(\sqrt{2} - x)^2$. Thus it holds for $1 \geq x \geq (1 + \sqrt{8})/7 = 0.5469\dots$




Figure 3: The cases $x > 0.547$ and $0.6 > x > 0.378$.

Step 2. $0.6 > x \geq 0.378$

Put R_1 into R_0 in homothetic position sharing a vertex other than the apex (see again Figure 3). Then $R_0 \setminus R_1$ can be covered by two triangles R' and R'' of sizes u and $2 - x - u\sqrt{2}$, respectively, for any choice of u .

Select a subset of triangles $\mathcal{H}' \subset \{R_2, \dots, R_n\}$ and let \mathcal{H}'' be the rest of the triangles. Define u such that $\sum_{R \in \mathcal{H}'} 2\text{Area}(R) =: cu^2$. Then the induction hypothesis implies that \mathcal{H}' can ρ -cover R' . If the rest of the triangles have sufficiently large total areas (at least $c\text{Area}(R'')$), then induction can be applied to \mathcal{H}'' and R'' and we are done. A sufficient condition for this is that

$$c - x^2 - cu^2 \geq c(2 - x - u\sqrt{2})^2. \quad (22)$$

This holds for $u_1 \leq u \leq u_2$, where

$$u_{1,2} = \frac{1}{3} \left(\sqrt{2}(2 - x) \mp \sqrt{-(2 - x)^2 + 3(1 - x^2/c)} \right).$$

Suppose now that u_1 and u_2 exists. We would like to select \mathcal{H}' such that u gets into the above range, i.e.,

$$cu_1^2 \leq \sum_{R_i \in \mathcal{H}'} x_i^2 \leq cu_2^2. \quad (23)$$

Select the triangles into \mathcal{H}' one by one but in a fixed order, say, x_2, x_3, \dots . If the difference of the left-hand side and the right-hand side of (23) is at least x^2 , then the sequence

$$0, x_2^2, x_2^2 + x_3^2, x_2^2 + x_3^2 + x_4^2, \dots, \sum_{2 \leq i \leq n} x_i^2 \quad (\geq c - x^2 \geq cu_1^2)$$

cannot jump the gap $[cu_1^2, cu_2^2]$, consequently, a partition $\mathcal{H}', \mathcal{H}''$ satisfying (22) exists.

Thus, we can use induction if $cu_2^2 - cu_1^2 \geq x^2$. Rearranging, this is equivalent to

$$c^2 \frac{32}{81} (2-x)^2 \left(-(2-x)^2 + 3 \left(1 - \frac{x^2}{c} \right) \right) - x^4 \geq 0.$$

One can easily see (e.g., by using Maple V as the author did) that this fourth-degree polynomial is indeed positive for $0.378 \leq x < 0.6$.

7. Gluing R_1, R_2 and R_3

In this section we continue the proof of Theorem 3. The method of Step 2 can be summarized in the following Lemma. Note that (like in Figure 3) R_0 can always be ρ -covered by three triangles $R', R'',$ and R''' of sizes $u, 2-v-u\sqrt{2},$ and $v,$ respectively.

Lemma 6 *Let $\mathcal{H}''' \subset \mathcal{H}$ be a set of triangles with total area $A/2,$ and suppose that \mathcal{H}''' can ρ -cover a triangle R''' of size $v.$ Let $\max\{x(R) : R \in \mathcal{H} \setminus \mathcal{H}'''\} \leq w.$ Suppose that*

$$c^2 \frac{32}{81} (2-v)^2 \left(-(2-v)^2 + 3 \left(1 - \frac{A}{c} \right) \right) - w^4 \geq 0. \quad (24)$$

Then there exists a partition $\mathcal{H}' \cup \mathcal{H}''$ of $\mathcal{H} \setminus \mathcal{H}'''$ and a real number $u,$ such that $\sum_{R \in \mathcal{H}'} 2\text{Area}(R) = cu^2$ and $\sum_{R \in \mathcal{H}''} 2\text{Area}(R) \geq c(2-v-u\sqrt{2})^2.$

Therefore, the induction hypothesis can be applied to \mathcal{H}' and R' and to \mathcal{H}'' and $R''.$ These together with \mathcal{H}''' yield a ρ -cover of $R_0.$

Step 3. $0.38 > x \geq 0.336$

In Step 2 we used Lemma 6 for $\mathcal{H}''' = \{R_1\}, v = x$ and $A = x^2.$ This time $\mathcal{H}''' = \{R_1, R_2\},$ i.e., we glue R_1 and R_2 together so that they can cover an isosceles right triangle R''' of size $v := (x+y)/\sqrt{2}.$ Then put R''' into R_0 in homothetic position sharing a vertex other than the apex. We can use induction applying Lemma 6 with $A := x^2 + y^2$ and $w := y^2.$ Substituting to the condition (24) we get

$$c^2 \frac{32}{81} \left(2 - \frac{x+y}{\sqrt{2}} \right)^2 \left(- \left(2 - \frac{x+y}{\sqrt{2}} \right)^2 + 3 \left(1 - \frac{x^2 + y^2}{c} \right) \right) - y^4 \geq 0.$$

One can easily see (e.g., by using Maple V), that this fourth degree polynomial is indeed positive for $(x, y) \in D,$ where D is a quadrilateral defined by the vertices

$(0.262, 0.262)$, $(0.5, 0.5)$, $(0.5, c - 1)$, and $(0.336, c - 1)$. Especially, from now on, we may suppose that

$$(c - 1) \leq y \leq x \leq 0.336, \quad y \leq 0.262. \quad (25)$$




Figure 4: Gluing R_1 and R_2 , and gluing R_1 , R_2 and R_3 .

Step 4. The case of small z .

Replace R_2 by another copy of R_3 . Then the system obtained ε -covers R_0 if Theorem 2 can be applied. This Theorem applies if the new system still has a total area at least $(1 + z)\text{Area}(R_0)$, i.e., if $1 + z \leq c - y^2 + z^2$. So from now on, we may suppose that the opposite holds:

$$(1 - c) + z - z^2 + y^2 \geq 0. \quad (26)$$

Step 5. $0.234 < x \leq 0.336$

Let $C \subset \mathbf{R}^3$ the set of those (x, y, z) points in 3-space which satisfy (25), (26) and $.234 \leq x$. In this section we apply induction if $(x_1, x_2, x_3) \in C$.

Since $y \leq 0.262$ we get from (26) that $z > 0.167$, so the sizes of R_1 , R_2 and R_3 are relatively close to each other. As before, attach the apex vertices of R_2 and R_3 so that a leg of R_3 is part of a leg of R_2 . Then they can cover an isosceles right triangle R^* of size $(y + z)/\sqrt{2}$. Then again glue R^* and R_1 together so that they can cover an isosceles right triangle R''' of size $x/\sqrt{2} + (y + z)/2$ (see Figure 4).

Apply Lemma 6 with $\mathcal{H}''' := \{R_1, R_2, R_3\}$, $v := 2 - x/\sqrt{2} - (y + z)/2$, $A := x^2 + y^2 + z^2$, and $w := z^4$. Substitute these values to (24):

$$c^2 \frac{32}{81} \left(2 - \frac{x}{\sqrt{2}} - \frac{y+z}{2} \right)^2 \left(- \left(2 - \frac{x}{\sqrt{2}} - \frac{y+z}{2} \right)^2 + 3 \left(1 - \frac{x^2 + y^2 + z^2}{c} \right) \right) - z^4. \quad (27)$$

One can easily see (e.g., by using Maple V) that this fourth-degree polynomial is indeed positive for $(x, y, z) \in C$. The calculations can be reduced to two variables, because if we start in any feasible point $(x, y, z) \in C$ and move in direction $(1, -\sqrt{2}, 0)$

(or its opposite direction) then (27) decreases. Hence the minimum is taken on the boundary of C .

Especially, from now on, we may suppose that $x \leq 0.234$.

8. ρ -Covering a Strip, the End of the Induction

In this section we finish the induction proof of Theorem 3. We need a lemma.

Lemma 7 *Suppose that P is a (finite) set of positive reals, $c, q > 0$ such that*

$$\sum_{p \in P} p^2 \geq c, \quad \max P = p_1, \quad 2p_1 \leq c(1 - q).$$

Then there exists a subset $S \subset P$ with

$$\sum_{p \in S} p \geq 2 \quad \text{and} \quad \frac{\min_{p \in S} p}{\max_{p \in S} p} \geq q.$$

Proof. Here p_1 , c , and q can be any reals, but we will use it for $p_1 = 0.234$, $c = (\sqrt{2} + 1)/2 = 1.207\dots$ and $q = 0.61$.

Consider the infinite region C on the positive quadrant of the plane defined as the union of rectangles of dimensions $2 \times p_1 q^i$, $i = 0, 1, 2, \dots$, and placed on the segment $(2i, 0), (2i + 2, 0)$. The area of C is $2p_1/(1 - q)$.

Next, order the members of P , $p_1 \geq p_2 \geq \dots$ and place squares of these sizes next to each other on the x -axis in the positive quadrant, i.e., let D be the union of squares with base segments defined by the vertices $(p_1 + \dots + p_{i-1}, 0)$ and $(p_1 + \dots + p_{i-1} + p_i, 0)$.

Since $\text{Area}(C) \leq \text{Area}(D)$, there is a segment $(2i, 0), (2i + 2, 0)$ which meets a square of size exceeding $p_1 q^i$. Suppose i is the smallest with this property. Here $i \geq 1$. Then S can consist of the sizes of those squares in D having a common point with the segment $(2i - 2, 0), (2i, 0)$. \square

Step 6. $x \leq 0.234$

Now we return to the proof of Theorem 3. Apply Lemma 7 for the set $P = \{x_1, x_2, \dots, x_n\}$ with $q = 0.61$. We obtain a subset $S = \{y_1, y_2, \dots, y_m\} \subset P$ such that

$$0.234 > y := y_1 \geq y_2 \geq \dots \geq y_m > 0.61y \tag{28}$$

and

$$y_1 + \cdots + y_m \geq 2. \quad (29)$$

The rest of the procedure is similar to the one in Section 3. Our aim is, like in Lemma 4, to define a trapezoid T with vertices $(0, 0)$, $(0, h)$, $(1, 0)$, and $(1 - h, h)$ and to find an $s \geq 1$ such that

(P1) the triangles of sizes y_1, \dots, y_{2s+1} can ε -cover T (here $2s + 1 \leq m$), and

(P2) $c\text{Area}(T) \geq \frac{1}{2} \sum_{1 \leq i \leq 2s+1} y_i^2$.

Then one can apply induction, and the rest of the triangles can ρ -cover $R_0 \setminus T$.

Let $g := 0.61y$. Consider the triangles of sizes y_{2i} and y_{2i+1} , turn them so that they touch each other at their hypotenuse and together they cover a rectangle Q_i of height g and base $y_{2i} + y_{2i+1} - g$. Note that the base of Q_i is at least $y_{2i} \geq g$ and at most $2y - g = 1.39y$.

Define s as the smallest integer satisfying

$$1 \leq y_1 + \sum_{1 \leq i \leq s} (y_{2i} + y_{2i+1} - g). \quad (30)$$

We claim that such an s exists, $7 \leq 2s + 1 \leq m$. Indeed, we get $1 \leq (1 + 1.39s)y \leq (1 + 1.39s)0.234$, this implies $s \geq 3$. On the other hand, by definition

$$1 > y_1 + \sum_{1 \leq i \leq s-1} (y_{2i} + y_{2i+1} - g). \quad (31)$$

This implies that

$$1 - y_1 > y_2 + y_4 + \cdots + y_{2s-2} \geq y_3 + y_5 + \cdots + y_{2s-1},$$

so $2 - y_1 > \sum_{i < 2s-1} y_i$. Thus at least two more y_i 's are needed to increase this sum to at least 2. Then (29) guarantees that these two positive y_i 's exist, and thus the sum in (30) will reach 1 after finitely many steps.

Similarly as in Section 3, we define the height h of the trapezoid T by the equation

$$1 = y_1 + \sum_{1 \leq i \leq s} (y_{2i} + y_{2i+1} - h). \quad (32)$$

Then the triangles of sizes y_1, \dots, y_{2s+1} ε -cover T . Comparing (30) and (32) we obtain that

$$y_{2s+1} \leq h. \quad (33)$$

We have to show that these inequalities imply **(P2)**, i.e.,

$$ch(2-h) \geq \sum_{1 \leq i \leq 2s+1} y_i^2. \quad (34)$$

Divide by c , add $((c-1)/c) \sum y_i^2$, rearrange and apply (32). We get that the above inequality is equivalent to

$$\begin{aligned} L &:= \frac{c-1}{c} \sum_{1 \leq i \leq 2s+1} y_i^2 \geq -h(2-h) + \sum y_i^2 \\ &= -h \left(2 \sum y_i - 2sh - h \right) + \sum y_i^2 = \sum_{1 \leq i \leq 2s+1} (y_i - h)^2. \end{aligned}$$

Define the function $p(t)$ as $\sum_{1 \leq i \leq 2s+1} (y_i - t)^2$. Then

$$L \geq p(t) \quad \text{for} \quad (2 - \sqrt{2})y_1 \leq t \leq \sqrt{2}y_{2s+1}. \quad (35)$$

Indeed, $L \geq p(t)$ if $((c-1)/c)y_i^2 \geq (y_i - t)^2$ for each i . Since $(c-1)/c = (\sqrt{2}-1)^2$, this holds for $(\sqrt{2}-1)y_i \geq |t - y_i|$, i.e., if $(2 - \sqrt{2})y_i \leq t \leq \sqrt{2}y_i$.

We have the lower bound $(2 - \sqrt{2})y_1 < g \leq h$ by (33). Then (35) implies that $L \geq p(h)$ certainly holds for

$$h \leq y_{2s+1}\sqrt{2}. \quad (36)$$

Subtract (31) from (32), and again use the monotonicity. We obtain

$$0 < y_{2s} + y_{2s+1} - sh + (s-1)g \leq y - sh + sy_{2s+1},$$

yielding $h < y_{2s+1} + (y/s)$. Since $y \leq y_{2s+1}/0.61$ we get $h \leq \sqrt{2}y_{2s+1}$ for $s \geq 4$. Therefore (36) holds for $s \geq 4$ and we are done.

The last remaining case is $s = 3$. We have $L \geq p(g)$ by (35), so our proof is complete if we show that $p(g) \geq p(h)$.

It is easy to see, that the second degree polynomial $p(t)$ takes its minimum at $\bar{y} := (\sum_i y_i)/(2s+1)$. We have $g \leq \bar{y}$, so $p(t) \leq p(g)$ for all $t \in [g, 2\bar{y} - g]$. So we are done if $h \leq 2\bar{y} - g$. Using (32) this is equivalent to

$$h + (2s+1)g \leq 2.$$

Our last task is to prove that this always holds for the case $s = 3$, $y < 1/4$. Then (32) gives $h \leq (7y-1) < 1/4$, so $h + 7g < 2$ follows from $g < y < 1/4$. \square

9. Conclusion, Remarks

Note that our theorem holds for **infinite** sets of triangles satisfying (5).

Moon and Moser's result was extended by Groemer [8] and Bezdek and Bezdek [3] to **higher dimensions** proving that $f(Q^{(d)}) = 2^d - 1$, where now $Q^{(d)}$ is the d -dimensional cube.

An algorithm for packing or covering a given set K with a sequence of sets $\{C_i\}$ is an **on-line** method if the sets C_i are given in sequence, and C_{i+1} is presented only after C_i has been put in place, without the option of changing the placement afterward. Januszewski, Lassak, Rote, and Woeginger [11] proved that in Euclidean d -space, every sequence of cubes of total volume greater than or equal to $2^d + 3$ can cover the unit cube in the on-line manner. This volume bound is astoundingly good, considering the best possible bound of $2^d - 1$ for the analogous off-line problem.

A recent study on **square coverings** is by Abbott and Kathchalski [1].

L. Fejes-Tóth **conjectured** that $2 \leq f(C) \leq 3$ for every planar disk, and $f(D) = 9/4$ for the **circular disks** (and for ellipses).

Concerning our computer aided proof of Theorem 3, with a little work Step 6 can be extended to cover all cases $x_1 \leq .25$, but this does not simplify the previous steps.

In [16] it was claimed that $f_\rho(R) \geq c = (1 + \sqrt{2})/2$ where R is the right isosceles triangle, but their argument seems to be rather incomplete. We **conjecture** more, that **any** placement with any rotations of the triangles of sizes

$$(1 - \eta)(\sqrt{2} - 1)^k, \quad \text{where } k = 0, 1, 2, \dots$$

cannot cover R .

There are many covering results where only a few number of smaller pieces can be used, see, e.g., some recent works of M. Lassak et al. These are obviously related to Hadwiger's conjecture and Borsuk's problem, see the recent problem book by Pach, Moser, and Brass [4].

Acknowledgments

The author is greatly thankful to I. Bárány and M. Simonovits for helpful conversations.

References

- [1] H. L. ABBOTT AND M. KATCHALSKI: Covering squares with squares. *Discrete Comput. Geom.* **24** (2000), 151–169.

- [2] A. S. BESICOVITCH: Measure of asymmetry of convex curves. *J. London Math. Soc.* **23** (1948), 237–240.
- [3] A. BEZDEK AND K. BEZDEK: Eine hinreichende Bedingung für die Überdeckung des Einheitswürfels durch homothetische Exemplare im n -dimensionalen euklidischen Raum. (German) [A sufficient condition for the covering of the unit cube by homothetic copies in the n -dimensional Euclidean space.] *Beiträge Algebra Geom.* **17** (1984), 5–21.
- [4] P. BRASS, W. MOSER, AND J. PACH: *Research Problems in Discrete Geometry*. Springer-Verlag, New York, 2005.
- [5] I. FÁRY: Sur la densité des réseaux de domaines convexes. (French) *Bull. Soc. Math. France* **78** (1950), 152–161.
- [6] L. FEJES-TÓTH: *Lagerungen in der Ebene auf der Kugel und im Raum*. (German) Springer-Verlag, Berlin, 1972.
- [7] Z. FÜREDI: Covering a triangle with homothetic copies. *Discrete Geometry*, pp. 435–445. *Monogr. Textbooks Pure Appl. Math.*, **253**, Dekker, New York, 2003.
- [8] H. GROEMER: Covering and packing properties of bounded sequences of convex sets. *Mathematika* **29** (1982), 18–31.
- [9] J. JANUSZEWSKI: Covering a triangle with sequences of its homothetic copies. *Period. Math. Hungar.* **36** (1998), 183–189.
- [10] J. JANUSZEWSKI: Translative covering a convex body by its homothetic copies. *Studia Sci. Math. Hungar.* **40** (2003), 341–348.
- [11] J. JANUSZEWSKI, M. LASSAK, G. ROTE, AND G. WOEGINGER: On-line q -adic covering by the method of the n th segment and its application to on-line covering by cubes. *Beiträge Algebra Geom.* **37** (1996), 51–65.
- [12] J. W. MOON AND L. MOSER: Some packing and covering theorems. *Colloq. Math.* **17** (1967), 103–110.
- [13] J. PACH AND P. K. AGARWAL: *Combinatorial Geometry*. Wiley, New York, 1995.
- [14] É. VÁSÁRHELYI: Über eine Überdeckung mit homothetischen Dreiecken. (German) [On a covering with homothetic triangles.] *Beiträge Algebra Geom.* **17** (1984), 61–70.
- [15] É. VÁSÁRHELYI: Covering of a triangle by homothetic triangles. *Studia Sci. Math. Hungar.* **28** (1993), 163–172.
- [16] CHANGQING XU, LIPING YUAN, AND REN DING: Covering isosceles right triangles with isosceles right triangles. *Geombinatorics* **14** (2005), 194–199.